

HUMAN IL-23 ANTIGEN BINDING PROTEINS

PATENT NUMBER	DOCUMENT ID	DATE PUBLISHED
20210024626	US 20210024626 A1	2021-01-28

INVENTOR INFORMATION

NAME	CITY	STATE	ZIP CODE	COUNTRY
Towne; Jennifer E.	Seattle	WA	N/A	US
Cheng; Janet D.	Montclair	CA	N/A	US
O'Neill; Jason C.	Brier	WA	N/A	US
Zhang; Yu	Shoreline	WA	N/A	US
Sun; Yu	Seattle	WA	N/A	US
Cerne; Heather	Seattle	WA	N/A	US
Piper; Derek E.	Santa Clara	CA	N/A	US
Ketchem; Randal R.	Snohomish	WA	N/A	US

APPLICATION NO	DATE FILED
17/060905	2020-10-01

DOMESTIC PRIORITY (CONTINUITY DATA)

parent US continuation 16793610 20200218 ABANDONED child US 17060905

parent US continuation 16453482 20190626 ABANDONED child US 16793610

parent US continuation 16194896 20181119 ABANDONED child US 16453482

parent US continuation 15922675 20180315 ABANDONED child US 16194896

parent US continuation 15271315 20160921 parent-grant-document US 9951129 child US 15922675

parent US continuation 13504449 20120831 parent-grant-document US 8722033 WO continuation PCT/US10/54148 20101026 child US 14228556

parent US division 14228556 20140328 parent-grant-document US 9487580 child US 15271315

us-provisional-application US 61381287 20100909

us-provisional-application US 61254982 20091026

US CLASS CURRENT:

[1/1](#)

CPC CURRENT

TYPE	CPC	DATE
CPCI	A 61 K 47/6845	2017-08-01
CPCI	A 61 K 39/395	2013-01-01
CPCI	A 61 P 37/00	2018-01-01
CPCI	C 07 K 16/24	2013-01-01
CPCI	C 07 K 16/244	2013-01-01

CPCA	C 07 K 2317/622	2013-01-01
CPCA	Y 02 A 50/30	2018-01-01
CPCA	C 07 K 2317/55	2013-01-01
CPCA	C 07 K 2317/626	2013-01-01
CPCA	C 07 K 2317/21	2013-01-01
CPCA	C 07 K 2317/565	2013-01-01
CPCA	C 07 K 2317/76	2013-01-01
CPCA	A 61 K 2039/505	2013-01-01
CPCA	C 07 K 2317/24	2013-01-01
CPCA	C 07 K 2317/31	2013-01-01
CPCA	C 07 K 2317/54	2013-01-01
CPCA	C 07 K 2317/92	2013-01-01

KWIC Hits

Abstract

Antigen binding proteins that bind to human IL-23 protein are provided. Nucleic acids encoding the antigen binding protein, vectors, and cells encoding the same as well as use of IL-23 antigen binding proteins for diagnostic and therapeutic purposes are also provided.

Background/Summary

CROSS-REFERENCE TO RELATED APPLICATIONS

[0001] This application claims the benefit under 35 U.S.C. 119(e) of U.S. patent application No. 61/254,982, filed Oct. 26, 2009 and U.S. patent application No. 61/381,287, filed Sep. 9, 2010, which are incorporated herein by reference.

BACKGROUND

[0002] Interleukin 23 (IL-23), a heterodimeric cytokine, is a potent inducer of pro-inflammatory cytokines. IL-23 is related to the heterodimeric cytokine Interleukin 12 (IL-12) both sharing a common p40 subunit. In IL-23, a unique p19 subunit is covalently bound to the p40 subunit. In IL-12, the unique subunit is p35 (Oppmann et al., *Immunity*, 2000, 13: 713-715). The IL-23 heterodimeric protein is secreted. Like IL-12, IL-23 is expressed by antigen presenting cells (such as dendritic cells and macrophages) in response to activation stimuli such as CD40 ligation, Toll-like receptor agonists and pathogens. IL-23 binds a heterodimeric receptor comprising an IL-12R β 1 subunit (which is shared with the IL-12 receptor) and a unique receptor subunit, IL-23R. The IL-12 receptor consists of IL-12W and IL-12R β 2. IL-23 binds its heterodimeric receptor and signals through JAK2 and Tyk2 to activate STAT1, 3, 4 and 5 (Parham et al., *J. Immunol.* 2002, 168:5699-708). The subunits of the receptor are predominantly co-expressed on activated or memory T cells and natural killer cells and also at lower levels on dendritic cells, monocytes, macrophages, microglia, keratinocytes and synovial fibroblasts. IL-23 and IL-12 act on different T cell subsets and play substantially different roles in vivo.

[0003] IL-23 acts on activated and memory T cells and promotes survival and expansion of the T cell subset, Th17. Th17 cells produce proinflammatory cytokines including IL-6, IL-17, TNF α , IL-22 and GM-CSF. IL-23 also acts on natural killer cells, dendritic cells and macrophages to induce pro-inflammatory cytokine expression. Unlike IL-23, IL-12 induces the differentiation of naïve CD4+T cells into mature Th1 IFN γ -producing effector cells, and induces NK and cytotoxic T cell function by stimulating IFN γ production. Th1 cells driven by IL-12 were previously thought to be the pathogenic T cell subset in many autoimmune diseases, however, more recent animal studies in models of inflammatory bowel disease, psoriasis, inflammatory arthritis and multiple sclerosis, in which the individual contributions of IL-12 versus IL-23 were evaluated have firmly established that IL-23, not IL-

12, is the key driver in autoimmune/inflammatory disease (Ahern et al., Immun. Rev. 2008 226:147-159; Cua et al., Nature 2003 421:744-748; Yago et al., Arthritis Res and Ther. 2007 9(5): R96). It is believed that IL-12 plays a critical role in the development of protective innate and adaptive immune responses to many intracellular pathogens and viruses and in tumor immune surveillance. See Kastelein, et al., Annual Review of Immunology, 2007, 25: 221-42; Liu, et al., Rheumatology, 2007, 46(8): 1266-73; Bowman et al., Current Opinion in Infectious Diseases, 2006 19:245-52; Fieschi and Casanova, Eur. J. Immunol. 2003 33:1461-4; Meeran et al., Mol. Cancer Ther. 2006 5: 825-32; Langowski et al., Nature 2006 442: 461-5. As such, IL-23 specific inhibition (sparing IL-12 or the shared p40 subunit) should have a potentially superior safety profile compared to dual inhibition of IL-12 and IL-23.

[0004] Therefore, use of IL-23 specific antagonists that inhibit human IL-23 (such as antibodies that bind at least the unique p19 subunit or bind both the p19 and p40 subunits of IL-23) that spare IL-12 should provide efficacy equal to or greater than IL-12 antagonists or p40 antagonists without the potential risks associated with inhibition of IL-12. Murine, humanized and phage display antibodies selected for inhibition of recombinant IL-23 have been described; see for example U.S. Pat. No. 7,491,391, WIPO Publications WO1999/05280, WO2007/0244846, WO2007/027714, WO 2007/076524, WO2007/147019, WO2008/103473, WO 2008/103432, WO2009/043933 and WO2009/082624. However, there is a need for fully human therapeutic agents that are able to inhibit native human IL-23. Such therapeutics are highly specific for the target, particularly in vivo. Complete inhibition of the in vivo target can result in lower dose formulations, less frequent and/or more effective dosing which in turn results in reduced cost and increased efficiency. The present invention provides such IL-23 antagonists.

SUMMARY

[0005] Antigen binding proteins that bind IL-23, particularly native human IL-23, are provided. The human IL-23 antigen binding proteins can reduce, inhibit, interfere with, and/or modulate at least one of the biological responses related to IL-23, and as such, are useful for ameliorating the effects of IL-23 related diseases or disorders. IL-23 antigen binding proteins can be used, for example, to reduce, inhibit, interfere with and/or modulate IL-23 signaling, IL-23 activation of Th17 cells, IL-23 activation of NK cells, or inducing production of proinflammatory cytokines.

[0006] Also provided are expression systems, including cell lines, for the production of IL-23 antigen binding proteins and methods of diagnosing and treating diseases related to human IL-23.

[0007] Some of the antigen binding proteins that bind IL-23 that are provided comprise at least one heavy chain variable region comprising a CDRH1, a CDRH2 and a CDRH3 selected from the group consisting of: a CDRH1 that differs by no more than one amino acid substitution, insertion or deletion from a CDRH1 as shown in TABLE 3; a CDRH2 that differs by no more than three, two or one amino acid substitutions, insertions and/or deletions from a CDRH2 as shown in TABLE 3; a CDRH3 that differs by no more than three, two or one amino acid substitutions, insertions and/or deletions from a CDRH3 as shown in TABLE 3; and comprising at least one light chain variable region comprising a CDRL1, a CDRL2 and a CDRL3 selected from the group consisting of: a CDRL1 that differs by no more than three, two or one amino acid substitutions, insertions and/or deletions from a CDRL1 as shown in TABLE 3; a CDRL2 that differs by no more than one amino acid substitution, insertion or deletion from a CDRL2 as shown in TABLE 3; a CDRL3 that differs by no more than one amino acid substitution, insertion or deletion from a CDRL3 as shown in TABLE 3. In one embodiment is provided isolated antigen binding proteins comprising: a CDRH1 selected from the group consisting of SEQ ID NO: 91, 94, 97, 100, and 103; a CDRH2 selected from the group consisting of SEQ ID NO:92, 95, 98, 101, 104, 107, and 110; a CDRH3 selected from the group consisting of SEQ ID NO: 93, 96, 99, 102, and 105; a CDRL1 selected from the group consisting of SEQ ID NO: 62, 65, 68, 71, and 74; a CDRL2 selected from the group consisting of SEQ ID NO:63, 66, 69, 72, 75, and 78; and a CDRL3 selected from the group consisting of SEQ ID NO:64, 67, 70 and 73. In another embodiment is provided isolated antigen bindings protein of comprising: a CDRH1 selected from the group consisting of SEQ ID NO: 91, 106, 109, 112, and 115; a CDRH2 selected from the group consisting of SEQ ID NO: 113, 116, 118, 120, 121, and 122; a CDRH3 selected from the group consisting of SEQ ID NO: 108, 111,

114, 117, and 119; a CDRL1 selected from the group consisting of SEQ ID NO: 77, 80, 83, 85, 86, 87, 88, 89 and 90; a CDRL2 is SEQ ID NO: 81; and a CDRL3 selected from the group consisting of SEQ ID NO: 76, 79, 82 and 84. In another embodiment is provided an isolated antigen-binding protein of that comprises at least one heavy chain variable region and at least one light chain variable region. In yet another embodiment is provided an isolated antigen-binding protein as described above that comprise at least two heavy chain variable regions and at least two light chain variable regions. In yet another embodiment is provided an isolated antigen binding protein wherein the antigen binding protein is coupled to a labeling group.

[0008] Also provided are isolated antigen binding proteins that bind IL-23 selected from the group consisting of a) an antigen binding protein having CDRH1 of SEQ ID NO:129, CDRH2 of SEQ ID NO:132, CDRH3 of SEQ ID NO:136, and CDRL1 of SEQ ID NO:123, CDRL2 of SEQ ID NO:81, and CDRL3 of SEQ ID NO: 76; b) an antigen binding protein having CDRH1 of SEQ ID NO:131, CDRH2 of SEQ ID NO: 134, CDRH3 of SEQ ID NO:137 and CDRL1 of SEQ ID NO:124, CDRL2 of SEQ ID NO:126 and CDRL3 of SEQ ID NO:128; c) an antigen binding protein having CDRH1 of SEQ ID NO:130, CDRH2 of SEQ ID NO:133, CDRH3 of SEQ ID NO:99 and CDRL1 of SEQ ID NO:68, CDRL2 of SEQ ID NO:69, and CDRL3 of SEQ ID NO:67; and d) an antigen binding protein having CDRH1 SEQ ID NO:91, CDRH2 SEQ ID NO: 135, CDRH3 SEQ ID NO:138 and CDRL1 SEQ ID NO:125, CDRL2 SEQ ID NO:127, and CDRL3 SEQ ID NO:64.

[0009] Also provided are isolated antigen binding proteins that bind IL-23 comprising at least one heavy chain variable region and at least one light chain variable region, selected from the group consisting of: a heavy chain variable region comprising amino acid residues 31-35, 50-65 and 99-113 of SEQ ID NO:31; and a light chain variable region comprising amino acid residues 23-36, 52-58 and 91-101 of SEQ ID NO:1; a heavy chain variable region comprising amino acid residues 31-35, 50-65 and 99-110 of SEQ ID NO:34 and heavy chain variable region comprising amino acid residues 31-35, 50-66 and 99-110 of SEQ ID NO:36; and a light chain variable region comprising amino acid residues 23-36, 52-62 and 97-105 of SEQ ID NO:4; a heavy chain variable region comprising amino acid residues 31-35, 50-66 and 99-114 of SEQ ID NO:38; and a light chain variable region comprising amino acid residues 23-34, 50-61 and 94-106 of SEQ ID NO:7; a heavy chain variable region comprising amino acid residues 31-35, 50-66 and 99-114 of SEQ ID NO:40; and a light chain variable region comprising amino acid residues 24-34, 50-56 and 94-106 of SEQ ID NO:9; a heavy chain variable region comprising amino acid residues 31-35, 50-66 and 99-114 of SEQ ID NO:42; and a light chain variable region comprising amino acid residues 23-34, 50-61 and 94-106 of SEQ ID NO:11; a heavy chain variable region comprising amino acid residues 31-35, 50-65 and 98-107 of SEQ ID NO:44; and a light chain variable region comprising amino acid residues 24-34, 50-56 and 89-97 of SEQ ID NO:13; a heavy chain variable region comprising amino acid residues 31-37, 52-67 and 100-109 of SEQ ID NO:46 or SEQ ID NO:153; and a light chain variable region comprising amino acid residues 24-34, 50-56 and 89-97 of SEQ ID NO:15; a heavy chain variable region comprising amino acid residues 31-37, 52-67 and 100-109 of SEQ ID NO:48; and a light chain variable region comprising amino acid residues 24-34, 50-56 and 89-97 of SEQ ID NO:17; a heavy chain variable region comprising amino acid residues 31-37, 52-67 and 101-109 of SEQ ID NO:50; and a light chain variable region comprising amino acid residues 24-34, 50-56 and 89-97 of SEQ ID NO:19; a heavy chain variable region comprising amino acid residues 31-35, 50-65 and 98-107 of SEQ ID NO: 52; and a light chain variable region comprising amino acid residues 24-34, 50-56 and 98-107 of SEQ ID NO:21; a heavy chain variable region comprising amino acid residues 31-37, 52-67 and 100-109 of SEQ ID NO:54; and a light chain variable region comprising amino acid residues 24-34, 50-56 and 89-97 of SEQ ID NO:23; a heavy chain variable region comprising amino acid residues 31-37, 52-67 and 100-109 of SEQ ID NO:56; and a light chain variable region comprising amino acid residues 24-34, 50-56 and 89-97 of SEQ ID NO:25; and a heavy chain variable region comprising amino acid residues 31-37, 52-57 and 100-109 of SEQ ID NO:58; and a light chain variable region comprising amino acid residues 24-34, 50-56 and 89-97 of SEQ ID NO:27.

[0010] Provided herein is an isolated antigen binding protein that binds IL-23 comprising a heavy chain variable region and a light chain variable region, wherein the heavy chain variable region sequence differs by no more than 13, 12, 11, 10, 9, 8, 7, 6, 5, 4, 3, 2 or 1 amino acid substitutions, additions and/or deletions from a heavy chain variable region sequence as shown in TABLE 2; and wherein the

light chain variable region sequence differs by no more than 13, 12, 11, 10, 9, 8, 7, 6, 5, 4, 3, 2 or 1 amino acid substitutions, additions and/or deletions from a light chain variable region sequence as shown in TABLE 1.

[0011] Also provided is an isolated antigen binding protein that binds IL-23 selected from the group consisting of a) a heavy chain variable region of SEQ ID NO:140 and a light chain variable region of SEQ ID NO: 30; b) a heavy chain variable region of SEQ ID NO:141 and a light chain variable region of SEQ ID NO:61; c) a heavy chain variable region of SEQ ID NO:142 and a light chain variable region of SEQ ID NO:4; and d) a heavy chain variable region of SEQ ID NO:143 and a light chain variable region of SEQ ID NO:139.

[0012] Also provided is an isolated antigen binding protein comprising a heavy chain variable region comprising of an amino acid sequence having at least 90%, 95%, 96%, 97%, 98% or 99% sequence identity to SEQ ID NO:31, 34, 36, 38, 40, 42, 44, 46, 48, 50, 52, 54, 56 and 58; and a light chain variable region comprising an amino acid sequence having at least 90% sequence identity to SEQ ID NO: 1, 4, 7, 9, 11, 13, 15, 17, 19, 21, 23, 25 and 27. In another embodiment is an isolated antigen binding protein comprising a heavy chain variable region selected from the group consisting of SEQ ID NO: 44, 46, 48, 50, 52, 54, 56, 58 and 153, and a light chain variable region selected from the group consisting of SEQ ID NO:13, 15, 17, 19, 21, 23, 25, and 27. In yet another embodiment is an isolated antigen binding protein comprising a heavy chain variable region selected from the group consisting of SEQ ID NO: 31, 34, 36, 38, 40 and 42, and a light chain variable region selected from the group consisting of SEQ ID NO: 1, 4, 7, 9 and 11.

[0013] Also provided is an isolated antigen binding protein that binds IL-23 comprising a heavy chain variable region and a light chain variable region selected from the group consisting of: a) a heavy chain variable region of SEQ ID NO:31 and a light chain variable region of SEQ ID NO:1; b) a heavy chain variable region of SEQ ID NO:34 or 36 and a light chain variable region of SEQ ID NO:4; c) a heavy chain variable region of SEQ ID NO:38 and a light chain variable region of SEQ ID NO: 7; d) a heavy chain variable region of SEQ ID NO:40 and a light chain variable region of SEQ ID NO:9; e) a heavy chain variable region of SEQ ID NO:42 and a light chain variable region of SEQ ID NO: 11; f) a heavy chain variable region of SEQ ID NO:44 and a light chain variable region of SEQ ID NO:13; g) a heavy chain variable region of SEQ ID NO:46 or SEQ ID NO:153 and a light chain variable region of SEQ ID NO:15; h) a heavy chain variable region of SEQ ID NO:48 and a light chain variable region of SEQ ID NO:17; i) a heavy chain variable region of SEQ ID NO:50 and a light chain variable region of SEQ ID NO: 19; j) a heavy chain variable region of SEQ ID NO:52 and a light chain variable region of SEQ ID NO:21; k) a heavy chain variable region of SEQ ID NO:54 and a light chain variable region of SEQ ID NO:23; l) a heavy chain variable region of SEQ ID NO:56 and a light chain variable region of SEQ ID NO:25; and m) a heavy chain variable region of SEQ ID NO:58 and a light chain variable region of SEQ ID NO:27.

[0014] Also provided is an isolated antigen binding protein that binds human IL-23, wherein the covered patch formed when the antigen binding protein is bound to human IL-23 comprises residue contacts 30, 31, 32, 49, 50, 52, 53, 56, 92 and 94 of SEQ ID NO:15, wherein the residue contacts have a difference value of greater than or equal to 10 Å^{sup.2} as determined by solvent exposed surface area. Within one embodiment the residue contacts comprise residues 31-35, 54, 58-60, 66, and 101-105 of SEQ ID NO:46.

[0015] Also provided is an isolated antigen binding protein that binds human IL-23, wherein the covered patch formed when the antigen binding protein is bound to human IL-23 comprises residue contacts 31-34, 51, 52, 55, 68, 93 and 98 of SEQ ID NO:1, wherein the residue contacts have a difference value of greater than or equal to 10 Å^{sup.2} as determined by solvent exposed surface area. Within one embodiment the residue contacts comprise residues 1, 26, 28, 31, 32, 52, 53, 59, 76, 101, 102 and 104-108 of SEQ ID NO:31.

[0016] Also provided is an isolated antigen binding protein that binds human IL-23, wherein when the antigen binding protein is bound to human IL-23, the antigen binding protein is 5 Å or less from residues 32-35, 54, 58-60, 66 and 101-105 of SEQ ID NO:46, as determined by X-ray crystallography.

In one embodiment the antigen binding protein is 5 Å or less from residues 31-35, 54, 56, 58-60, 66 and 101-105 of SEQ ID NO:46.

[0017] Also provided is an isolated antigen binding protein that binds human IL-23, wherein when the antigen binding protein is bound to human IL-23, the antigen binding protein is 5 Å or less from residues 30-32, 49, 52, 53, 91-94 and 96 of SEQ ID NO:15, as determined by X-ray crystallography. In one embodiment the antigen binding protein is 5 Å or less from residues 30-32, 49, 50, 52, 53, 56, 91-94 and 96 of SEQ ID NO:15.

[0018] Also provided is an isolated antigen binding protein that binds human IL-23, wherein when the antigen binding protein is bound to human IL-23, the antigen binding protein is 5 Å or less from residues 26-28, 31, 53, 59, 102 and 104-108 of SEQ ID NO:31, as determined by X-ray crystallography. In one embodiment the antigen binding protein is 5 Å or less from residues 1, 26-28, 30-32, 52, 53, 59, 100, and 102-108 of SEQ ID NO:31.

[0019] Also provided is an isolated antigen binding protein that binds human IL-23, wherein when said antigen binding protein is bound to human IL-23, said antigen binding protein is 5 Å or less from residues 31-34, 51, 52, 55, 68 and 93 of SEQ ID NO:1 as determined by X-ray crystallography. In one embodiment the antigen binding protein is 5 Å or less from residues 29, 31-34, 51, 52, 55, 68, 93 and 100 of SEQ ID NO:1.

[0020] Also provided is an isolated antigen binding protein as described above, wherein the antigen binding protein is an antibody. In one embodiment is provided an isolated antigen binding protein wherein the antibody is a monoclonal antibody, a recombinant antibody, a human antibody, a humanized antibody, a chimeric antibody, a multispecific antibody, or an antibody fragment thereof. In another embodiment is provided an isolated antigen binding protein wherein the antibody fragment is a Fab fragment, a Fab' fragment, a F(ab')₂ fragment, a Fv fragment, a diabody, or a single chain antibody molecule. In yet another embodiment is provided an isolated antigen binding protein wherein the antigen binding protein is a human antibody. In still another embodiment is provided an isolated antigen binding protein wherein the antigen binding protein is a monoclonal antibody. In another embodiment is provided an isolated antigen binding protein wherein the antigen binding protein is of the IgG1-, IgG2- IgG3- or IgG4-type. In yet another embodiment is provided an isolated antigen binding protein wherein the antigen binding protein is of the IgG1- or IgG2-type.

[0021] An isolated nucleic acid molecule encoding an antigen binding protein as described above, is also provided. In one embodiment is provided an isolated nucleic acid molecule wherein at least one heavy chain variable region is encoded by an isolated nucleic acid molecule selected from the group consisting of SEQ ID NOs:32, 35, 37, 39, 41, 43, 45, 47, 49, 51, 53, 55, 57, 59 and 152 and at least one light chain variable region is encoded by an isolated nucleic acid molecule selected from the group consisting of SEQ ID NOs:2, 5, 6, 8, 10, 12, 14, 16, 18, 20, 22, 24, 26, and 28. In another embodiment is provided a nucleic acid molecule wherein the nucleic acid molecule is operably linked to a control sequence. In another embodiment is provided a vector comprising a nucleic acid molecule as described above. In yet another embodiment is provided a host cell comprising the nucleic acid molecule as described above. In another embodiment is provided a host cell comprising the vector described above. In yet another embodiment is provided an isolated polynucleotide sufficient for use as a hybridization probe, PCR primer or sequencing primer that is a fragment of the nucleic acid molecule as described above or its complement.

[0022] Also provided is a method of making the antigen binding protein as described above, comprising the step of preparing said antigen binding protein from a host cell that secretes said antigen binding protein.

[0023] Also provided is an isolated antigen binding protein that binds human IL-23, wherein the covered patch formed when the antigen binding protein is bound to human IL-23 comprises a residue contact within residues 46-58, a residue contact within residues 112-120, and a residue contact within residues 155-163 of the human IL-23p19 subunit as described in SEQ ID NO:145, wherein the residue contact has a difference value greater than or equal to 10 Å^{sup.2} as determined by solvent exposed

surface area. In one embodiment is provided wherein the covered patch formed when the antigen binding protein is bound to human IL-23 comprises one, two, three, four, five, six, seven, eight, nine, ten, eleven, twelve or thirteen residue contacts within residues 46-58, one, two, three, four, five, six, seven, eight, nine or ten residue contacts within residues 112-120, and one, two, three, four, five, six, seven, eight or nine residue contacts within residues 155-163 of the human IL-23p19 subunit as described in SEQ ID NO:145. In another embodiment is provided wherein the covered patch formed when the antigen binding protein binds to human IL-23 comprises a residue contact within residues 121-125 of the human IL-23p40 subunit as described in SEQ ID NO:147. In a related embodiment is wherein the covered patch formed when the antigen binding protein is bound to human IL-23 comprises one, two, three, four or five residue contacts within residues 121-125 of the human IL-23p40 subunit as described in SEQ ID NO:147. Within another embodiment is provided wherein the covered patch formed when the antigen binding protein is bound to human IL-23 comprises residue contacts 46, 47, 49, 50, 53, 112-116, 118, 120, 155, 156, 159, 160, and 163 of SEQ ID NO:145. In another embodiment is provided wherein the covered patch formed when the antigen binding protein is bound to human IL-23 comprises residue contacts 46, 47, 49, 50, 53, 112-118, 120, 155, 156, 159, 160, and 163 of SEQ ID NO:145. Within another embodiment is provided wherein the covered patch formed when the antigen binding protein is bound to human IL-23 comprises residues 46, 47, 49, 50, 53-55, 57, 58, 112-116, 118-120, 155, 156, 159, 160, 162 and 163 of SEQ ID NO:145. In a related embodiment is provided wherein the covered patch formed when the antigen binding protein is bound to human IL-23 comprises residue contact 122 of the human IL-23p40 subunit as described in SEQ ID NO:147. In another related embodiment is provided wherein the covered patch formed when the antigen binding protein is bound to human IL-23 comprises residue contacts 122 and 124 of the human IL-23p40 subunit as described in SEQ ID NO:147. In yet another related embodiment is provided wherein the covered patch formed when the antigen binding protein is bound to human IL-23 comprises residue contact 121-123 and 125 of the human IL-23p40 subunit as described in SEQ ID NO:147. In a further related embodiment is provided wherein the covered patch formed when the antigen binding protein is bound to human IL-23 comprises residue contact 121-123, 125 and 283 of the human IL-23p40 subunit as described in SEQ ID NO:147.

[0024] Also provided is an isolated antigen binding protein that binds human IL-23, wherein when said antigen binding protein is bound to human IL-23 said antigen binding protein is 5 Å or less from a residue within residues 46-58, from a residue within residues 112-123, and from a residue within residues 155-163 of the human IL-23p19 subunit as described in SEQ ID NO:145, as determined by X-ray crystallography. In one embodiment, when the antigen binding protein is bound to human IL-23, the antigen binding protein is 5 Å or less from one, two, three, four, five, six, seven, eight, nine, ten, eleven, twelve or thirteen residues within residues 46-58, from one, two, three, four, five, six, seven, eight, nine or ten, residues within residues 112-123, and from one, two, three, four, five, six, seven, eight or nine residues within residues 155-163 of the human IL-23p19 subunit as described in SEQ ID NO:145. Within another embodiment when the antigen binding protein is bound to human IL-23 the antigen binding protein is 5 Å or less from residues 46-50, 113-116, 120, 156, 159, 160 and 163 of SEQ ID NO:145. Within another embodiment when the antigen binding protein is bound to human IL-23, the antigen binding protein is 5 Å or less from residues 46-50, 112-120, 156, 159, 160 and 163 of SEQ ID NO:145. Within a related embodiment when the antigen binding protein is bound to human IL-23, the antigen binding protein is 5 Å or less from residues 46-50, 53, 112-120, 156, 159, 160 and 163 of SEQ ID NO:145. Within another embodiment when the antigen binding protein is bound to human IL-23, the antigen binding protein is 5 Å or less from residues 46-50, 53-55, 58, 113-116, 120, 121, 156, 159, 160, 162 and 163 of SEQ ID NO:145. Within a related embodiment when the antigen binding protein is bound to human IL-23, the antigen binding protein is 5 Å or less from residues 46-51, 53-55, 57, 58, 112-116, 118-121, 123, 155, 156, 159, 160, 162 and 163 of SEQ ID NO:145. Within a further embodiment when the antigen binding protein is bound to human IL-23 the antigen binding protein is 5 Å or less from a residue within residues 121-125, of the human IL-23p40 subunit as described in SEQ ID NO:147, as determined by X-ray crystallography. With a related embodiment when the antigen binding protein is bound to human IL-23, said antigen binding protein is 5 Å or less from residues 122 and 124 of SEQ ID NO:147. Within another embodiment when the antigen binding protein is bound to human IL-23, the antigen binding protein is 5 Å or less from residues 121-123 and 125 of SEQ ID NO:147.

[0025] Also provided is an isolated antigen binding protein as described above, wherein the antigen binding protein has at least one property selected from the group consisting of: a) reducing human IL-23 activity; b) reducing production of a proinflammatory cytokine; c) binding to human IL-23 with a KD of $\leq 5 \times 10^{-8}$ M; d) having a Koff rate of $\leq 5 \times 10^{-6}$ 1/s; and d) having an IC50 of ≤ 400 pM.

[0026] A pharmaceutical composition comprising at least one antigen binding protein as described above and pharmaceutically acceptable excipient is provided. In one embodiment is provided a pharmaceutical composition further comprises a labeling group or an effector group. In yet another embodiment is provided a pharmaceutical composition wherein the labeling group is selected from the group consisting of isotopic labels, magnetic labels, redox active moieties, optical dyes, biotinylated groups and predetermined polypeptide epitopes recognized by a secondary reporter. In yet another embodiment is provided a pharmaceutical composition wherein the effector group is selected from the group consisting of a radioisotope, radionuclide, a toxin, a therapeutic group and a chemotherapeutic group.

[0027] Also provided is a method for treating or preventing a condition associated with IL-23 in a patient, comprising administering to a patient in need thereof an effective amount of at least one isolated antigen binding protein as described above. In one embodiment is provided a method of wherein the condition is selected from the group consisting of an inflammatory disorder, a rheumatic disorder, an autoimmune disorder, an oncological disorder and a gastrointestinal disorder. In yet another embodiment is provided a method wherein the condition is selected from the group consisting of multiple sclerosis, rheumatoid arthritis, cancer, psoriasis, inflammatory bowel disease, Crohn's disease, ulcerative colitis, systemic lupus erythematosus, psoriatic arthritis, autoimmune myocarditis; type 1 diabetes and ankylosing spondylitis. In still another embodiment is provided a method wherein the isolated antigen-binding protein is administered alone or as a combination therapy.

[0028] Also provided is a method of reducing IL-23 activity in a patient comprising administering an effective amount of at least one antigen binding protein as described above. In one embodiment is provided a method of reducing IL-23 activity, wherein said IL-23 activity is inducing production of a proinflammatory cytokine.

Description

BRIEF DESCRIPTION OF THE DRAWINGS

[0029] FIG. 1A: Results of STAT-luciferase reporter assay using recombinant human IL-23. All antibodies completely inhibited recombinant human IL-23

[0030] FIG. 1B: Results from STAT-luciferase reporter assay using native human IL-23. Only half of those antibodies that completely inhibited recombinant human IL-23 were able to completely inhibit native human IL-23

DETAILED DESCRIPTION

[0031] The present invention provides compositions, kits, and methods relating to IL-23 antigen binding proteins, including molecules that antagonize IL-23, such as anti-IL-23 antibodies, antibody fragments, and antibody derivatives, e.g., antagonistic anti-IL-23 antibodies, antibody fragments, or antibody derivatives. Also provided are polynucleotides, and derivatives and fragments thereof, comprising a sequence of nucleic acids that encodes all or a portion of a polypeptide that binds to IL-23, e.g., a polynucleotide encoding all or part of an anti-IL-23 antibody, antibody fragment, or antibody derivative, plasmids and vectors comprising such nucleic acids, and cells or cell lines comprising such polynucleotides and/or vectors and plasmids. The provided methods include, for example, methods of making, identifying, or isolating IL-23 antigen binding proteins, such as anti-IL-23 antibodies, methods of determining whether a molecule binds to IL-23, methods of determining whether a molecule antagonizes IL-23, methods of making compositions, such as pharmaceutical compositions, comprising an IL-23 antigen binding protein, and methods for administering an IL-23 antigen binding

protein to a subject, for example, methods for treating a condition mediated by IL-23, and for antagonizing a biological activity of IL-23, in vivo or in vitro.

[0032] Unless otherwise defined herein, scientific and technical terms used in connection with the present invention shall have the meanings that are commonly understood by those of ordinary skill in the art. Further, unless otherwise required by context, singular terms shall include pluralities and plural terms shall include the singular. Generally, nomenclatures used in connection with, and techniques of, cell and tissue culture, molecular biology, immunology, microbiology, genetics and protein and nucleic acid chemistry and hybridization described herein are those well known and commonly used in the art. The methods and techniques of the present invention are generally performed according to conventional methods well known in the art and as described in various general and more specific references that are cited and discussed throughout the present specification unless otherwise indicated. See, e.g., Sambrook et al., *Molecular Cloning: A Laboratory Manual*, 3rd ed., Cold Spring Harbor Laboratory Press, Cold Spring Harbor, N.Y. (2001) and Ausubel et al., *Current Protocols in Molecular Biology*, Greene Publishing Associates (1992), and Harlow and Lane *Antibodies: A Laboratory Manual* Cold Spring Harbor Laboratory Press, Cold Spring Harbor, N.Y. (1990). Enzymatic reactions and purification techniques are performed according to manufacturer's specifications, as commonly accomplished in the art or as described herein. The terminology used in connection with, and the laboratory procedures and techniques of, analytical chemistry, synthetic organic chemistry, and medicinal and pharmaceutical chemistry described herein are those well known and commonly used in the art. Standard techniques can be used for chemical syntheses, chemical analyses, pharmaceutical preparation, formulation, and delivery, and treatment of patients.

[0033] All patents and other publications identified are expressly incorporated herein by reference in their entirety for the purpose of describing and disclosing, for example, the methodologies described in such publications that might be used in connection with information described herein.

[0034] The polynucleotide and protein sequences of the p19 subunit of human IL-23 (SEQ ID NOs: 144 and 145), the shared p40 subunit (SEQ ID NOs: 146 and 147), the human IL-23 receptor heterodimeric subunits IL-12R β 1 (SEQ ID NOs: 150 and 151) and IL-23R (SEQ ID NOs: 148 and 149), are known in the art, see for example, GenBank Accession Nos. AB030000; M65272, NM_005535, NM_144701, as are those from other mammalian species. Recombinant IL-23 and IL-23 receptor proteins including single chain and Fc proteins as well as cells expressing the IL-23 receptor have been described or are available from commercial sources. (see for example, Oppmann et al., *Immunity*, 2000, 13: 713-715; R&D Systems, Minneapolis, Minnesota; United States Biological, Swampscott, Mass.; WIPO Publication No. WO 2007/076524). Native human IL-23 can be obtained from human cells such as dendritic cells using methods known in the art including those described herein.

[0035] IL-23 is a heterodimeric cytokine comprised of a unique p19 subunit that is covalently bound to a shared p40 subunit. The p19 subunit comprises four α -helices, "A", "B", "C" and "D" in an up-up-down-down motif joined by three intra-helix loops between the A and B helices, between the B and C helices and between the C and D helices, see Oppmann et al., *Immunity*, 2000, 13: 713-715 and Beyer, et al., *J Mol Biol*, 2008, 382(4): 942-55. The A and D helices of 4 helical bundle cytokines are believed to be involved with receptor binding. The p40 subunit comprises three beta-sheet sandwich domains, D1, D2 and D3 (Lupardus and Garcia, *J. Mol. Biol.*, 2008, 382:931-941).

[0036] The term "polynucleotide" includes both single-stranded and double-stranded nucleic acids and includes genomic DNA, RNA, mRNA, cDNA, or synthetic origin or some combination thereof which is not associated with sequences normally found in nature. Isolated polynucleotides comprising specified sequences may include, in addition to the specified sequences, coding sequences for up to ten or even up to twenty other proteins or portions thereof, or may include operably linked regulatory sequences that control expression of the coding region of the recited nucleic acid sequences, and/or may include vector sequences. The nucleotides comprising the polynucleotide can be ribonucleotides or deoxyribonucleotides or a modified form of either type of nucleotide. The modifications include base modifications such as bromouridine and inosine derivatives, ribose modifications such as 2',3'-dideoxyribose, and internucleotide linkage modifications such as phosphorothioate,

phosphorodithioate, phosphoroselenoate, phosphorodiselenoate, phosphoroanilothioate, phosphoraniladate and phosphoroamidate.

[0037] The term “oligonucleotide” means a polynucleotide comprising 100 or fewer nucleotides. In some embodiments, oligonucleotides are 10 to 60 bases in length. In other embodiments, oligonucleotides are 12, 13, 14, 15, 16, 17, 18, 19, or 20 to 40 nucleotides in length. Oligonucleotides may be single stranded or double stranded, e.g., for use in the construction of a mutant gene. Oligonucleotides may be sense or antisense oligonucleotides. An oligonucleotide can include a detectable label, such as a radiolabel, a fluorescent label, a hapten or an antigenic label, for detection assays. Oligonucleotides may be used, for example, as PCR primers, cloning primers or hybridization probes.

[0038] The terms “polypeptide” or “protein” means a macromolecule having the amino acid sequence of a native protein, that is, a protein produced by a naturally-occurring and non-recombinant cell; or it is produced by a genetically-engineered or recombinant cell, and comprise molecules having the amino acid sequence of the native protein, or molecules having one or more deletions from, insertions to, and/or substitutions of the amino acid residues of the native sequence. The term also includes amino acid polymers in which one or more amino acids are chemical analogs of a corresponding naturally-occurring amino acid and polymers. The terms “polypeptide” and “protein” encompass IL-23 antigen binding proteins (such as antibodies) and sequences that have one or more deletions from, additions to, and/or substitutions of the amino acid residues of the antigen binding protein sequence. The term “polypeptide fragment” refers to a polypeptide that has an amino-terminal deletion, a carboxyl-terminal deletion, and/or an internal deletion as compared with the full-length native protein. Such fragments may also contain modified amino acids as compared with the native protein. In certain embodiments, fragments are about five to 500 amino acids long. For example, fragments may be at least 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 50, 70, 100, 110, 150, 200, 250, 300, 350, 400, or 450 amino acids long. Useful polypeptide fragments include immunologically functional fragments of antibodies, including binding domains. In the case of an IL-23 antigen binding protein, such as an antibody, useful fragments include but are not limited to one or more CDR regions, a variable domain of a heavy or light chain, a portion of an antibody chain, a portion of a variable region including less than three CDRs, and the like.

[0039] “Amino acid” includes its normal meaning in the art. The twenty naturally-occurring amino acids and their abbreviations follow conventional usage. See, Immunology-A Synthesis, 2nd Edition, (E. S. Golub and D. R. Gren, eds.), Sinauer Associates: Sunderland, Mass. (1991). Stereoisomers (e.g., D-amino acids) of the twenty conventional amino acids, unnatural amino acids such as [alpha]-, [alpha]-disubstituted amino acids, N-alkyl amino acids, and other unconventional amino acids may also be suitable components for polypeptides. Examples of unconventional amino acids include: 4-hydroxyproline, [gamma]-carboxyglutamate, [epsilon]-N,N,N-trimethyllysine, [epsilon]-N-acetyllysine, O-phosphoserine, N-acetylserine, N-formylmethionine, 3-methylhistidine, 5-hydroxylysine, [sigma]-N-methylarginine, and other similar amino acids and imino acids (e.g., 4-hydroxyproline). In the polypeptide notation used herein, the left-hand direction is the amino terminal direction and the right-hand direction is the carboxyl-terminal direction, in accordance with standard usage and convention.

[0040] The term “isolated protein” refers to a protein, such as an antigen binding protein (an example of which could be an antibody), that is purified from proteins or polypeptides or other contaminants that would interfere with its therapeutic, diagnostic, prophylactic, research or other use. As used herein, “substantially pure” means that the described species of molecule is the predominant species present, that is, on a molar basis it is more abundant than any other individual species in the same mixture. In certain embodiments, a substantially pure molecule is a composition wherein the object species comprises at least 50% (on a molar basis) of all macromolecular species present. In other embodiments, a substantially pure composition will comprise at least 80%, 85%, 90%, 95%, or 99% of all macromolecular species present in the composition. In certain embodiments, an essentially homogeneous substance has been purified to such a degree that contaminating species cannot be detected in the composition by conventional detection methods and thus the composition consists of a single detectable macromolecular species.

[0041] A “variant” of a polypeptide (e.g., an antigen binding protein such as an antibody) comprises an amino acid sequence wherein one or more amino acid residues are inserted into, deleted from and/or substituted into the amino acid sequence relative to another polypeptide sequence. Variants include fusion proteins. A “derivative” of a polypeptide is a polypeptide that has been chemically modified in some manner distinct from insertion, deletion, or substitution variants, e.g., via conjugation to another chemical moiety.

[0042] The terms “naturally occurring” or “native” as used throughout the specification in connection with biological materials such as polypeptides, nucleic acids, host cells, and the like, refers to materials which are found in nature, such as native human IL-23. In certain aspects, recombinant antigen binding proteins that bind native IL-23 are provided. In this context, a “recombinant protein” is a protein made using recombinant techniques, i.e., through the expression of a recombinant nucleic acid as described herein. Methods and techniques for the production of recombinant proteins are well known in the art.

[0043] The term “antibody” refers to an intact immunoglobulin of any isotype, or a fragment thereof that can compete with the intact antibody for specific binding to the target antigen, and includes, for instance, chimeric, humanized, fully human, and bispecific antibodies. An antibody as such is a species of an antigen binding protein. Unless otherwise indicated, the term “antibody” includes, in addition to antibodies comprising two full-length heavy chains and two full-length light chains, derivatives, variants, fragments, and muteins thereof, examples of which are described below. An intact antibody generally will comprise at least two full-length heavy chains and two full-length light chains, but in some instances may include fewer chains such as antibodies naturally occurring in camelids which may comprise only heavy chains. Antibodies may be derived solely from a single source, or may be “chimeric,” that is, different portions of the antibody may be derived from two different antibodies as described further below. The antigen binding proteins, antibodies, or binding fragments may be produced in hybridomas, by recombinant DNA techniques, or by enzymatic or chemical cleavage of intact antibodies.

[0044] The term “functional fragment” (or simply “fragment”) of an antibody or immunoglobulin chain (heavy or light chain), as used herein, is an antigen binding protein comprising a portion (regardless of how that portion is obtained or synthesized) of an antibody that lacks at least some of the amino acids present in a full-length chain but which is capable of specifically binding to an antigen. Such fragments are biologically active in that they bind specifically to the target antigen and can compete with other antigen binding proteins, including intact antibodies, for specific binding to a given epitope. In one aspect, such a fragment will retain at least one CDR present in the full-length light or heavy chain, and in some embodiments will comprise a single heavy chain and/or light chain or portion thereof. These biologically active fragments may be produced by recombinant DNA techniques, or may be produced by enzymatic or chemical cleavage of antigen binding proteins, including intact antibodies. Fragments include, but are not limited to, immunologically functional fragments such as Fab, Fab', F(ab')₂, Fv, domain antibodies and single-chain antibodies, and may be derived from any mammalian source, including but not limited to human, mouse, rat, camelid or rabbit. It is contemplated further that a functional portion of the antigen binding proteins disclosed herein, for example, one or more CDRs, could be covalently bound to a second protein or to a small molecule to create a therapeutic agent directed to a particular target in the body, possessing bifunctional therapeutic properties, or having a prolonged serum half-life.

[0045] The term “compete” when used in the context of antigen binding proteins (e.g., neutralizing antigen binding proteins or neutralizing antibodies) means competition between antigen binding proteins as determined by an assay in which the antigen binding protein (e.g., antibody or immunologically functional fragment thereof) under test prevents or inhibits specific binding of a reference antigen binding protein (e.g., a ligand, or a reference antibody) to a common antigen (e.g., an IL-23 protein or a fragment thereof). Numerous types of competitive binding assays can be used, for example: solid phase direct or indirect radioimmunoassay (RIA), solid phase direct or indirect enzyme immunoassay (EIA), sandwich competition assay (see, e.g., Stahli et al., 1983, *Methods in Enzymology* 92:242-253); solid phase direct biotin-avidin EIA (see, e.g., Kirkland et al., 1986, *J. Immunol.* 137:3614-3619) solid phase direct labeled assay, solid phase direct labeled sandwich assay

(see, e.g., Harlow and Lane, 1988, *Antibodies, A Laboratory Manual*, Cold Spring Harbor Press); solid phase direct label RIA using 1-125 label (see, e.g., Morel et al., 1988, *Molec. Immunol.* 25:7-15); solid phase direct biotin-avidin EIA (see, e.g., Cheung, et al., 1990, *Virology* 176:546-552); and direct labeled RIA (Moldenhauer et al., 1990, *Scand. J. Immunol.* 32:77-82). Typically, such an assay involves the use of purified antigen bound to a solid surface or cells bearing either of these, an unlabelled test antigen binding protein and a labeled reference antigen binding protein.

[0046] Competitive inhibition is measured by determining the amount of label bound to the solid surface or cells in the presence of the test antigen binding protein. Usually the test antigen binding protein is present in excess. Antigen binding proteins identified by competition assay (competing antigen binding proteins) include antigen binding proteins binding to the same epitope as the reference antigen binding proteins and antigen binding proteins binding to an adjacent epitope sufficiently proximal to the epitope bound by the reference antigen binding protein for steric hindrance to occur. Usually, when a competing antigen binding protein is present in excess, it will inhibit specific binding of a reference antigen binding protein to a common antigen by at least 40%, 45%, 50%, 55%, 60%, 65%, 70% or 75%. In some instance, binding is inhibited by at least 80%, 85%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97% 98%, 99% or more.

[0047] The term “epitope” or “antigenic determinant” refers to a site on an antigen to which an antigen binding protein binds. Epitopes can be formed both from contiguous amino acids or noncontiguous amino acids juxtaposed by tertiary folding of a protein. Epitopes formed from contiguous amino acids are typically retained on exposure to denaturing solvents, whereas epitopes formed by tertiary folding are typically lost on treatment with denaturing solvents. Epitope determinants may include chemically active surface groupings of molecules such as amino acids, sugar side chains, phosphoryl or sulfonyl groups, and may have specific three dimensional structural characteristics, and/or specific charge characteristics. An epitope typically includes at least 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 25, 30, 35 amino acids in a unique spatial conformation. Epitopes can be determined using methods known in the art.

IL-23 Antigen Binding Proteins

[0048] An “antigen binding protein” as used herein means a protein that specifically binds a specified target antigen; the antigen as provided herein is IL-23, particularly human IL-23, including native human IL-23. Antigen binding proteins as provided herein interact with at least a portion of the unique p19 subunit of IL-23, detectably binding IL-23; but do not bind with any significance to IL-12 (e.g., the p40 and/or the p35 subunits of IL-12), thus “sparing IL-12”. As a consequence, the antigen binding proteins provided herein are capable of impacting IL-23 activity without the potential risks that inhibition of IL-12 or the shared p40 subunit might incur. The antigen binding proteins may impact the ability of IL-23 to interact with its receptor, for example by impacting binding to the receptor, such as by interfering with receptor association. In particular, such antigen binding proteins totally or partially reduce, inhibit, interfere with or modulate one or more biological activities of IL-23. Such inhibition or neutralization disrupts a biological response in the presence of the antigen binding protein compared to the response in the absence of the antigen binding protein and can be determined using assays known in the art and described herein. Antigen binding proteins provided herein inhibit IL-23-induced proinflammatory cytokine production, for example IL-23-induced IL-22 production in whole blood cells and IL-23-induced IFN γ expression in NK and whole blood cells. Reduction of biological activity can be about 20%, 30%, 40%, 50%, 60%, 70%, 80%, 85%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97% 98%, 99% or more.

[0049] An antigen binding protein may comprise a portion that binds to an antigen and, optionally, a scaffold or framework portion that allows the antigen binding portion to adopt a conformation that promotes binding of the antigen binding protein to the antigen. Examples of antigen binding proteins include antibodies, antibody fragments (e.g., an antigen binding portion of an antibody), antibody derivatives, and antibody analogs. The antigen binding protein can comprise an alternative protein scaffold or artificial scaffold with grafted CDRs or CDR derivatives. Such scaffolds include, but are not limited to, antibody-derived scaffolds comprising mutations introduced to, for example, stabilize the three-dimensional structure of the antigen binding protein as well as wholly synthetic scaffolds

comprising, for example, a biocompatible polymer. See, for example, Korndorfer et al., *Proteins: Structure, Function, and Bioinformatics*, (2003) Volume 53, Issue 1:121-129; Roque et al., *Biotechnol. Prog.*, 2004, 20:639-654. In addition, peptide antibody mimetics (“PAMs”) can be used, as well as scaffolds based on antibody mimetics utilizing fibronectin components as a scaffold.

[0050] Certain antigen binding proteins described herein are antibodies or are derived from antibodies. Such antigen binding proteins include, but are not limited to, monoclonal antibodies, bispecific antibodies, minibodies, domain antibodies, synthetic antibodies, antibody mimetics, chimeric antibodies, humanized antibodies, human antibodies, antibody fusions, antibody conjugates, single chain antibodies, and fragments thereof, respectively. In some instances, the antigen binding protein is an immunological fragment of an antibody (e.g., a Fab, a Fab', a F(ab')₂, or a scFv). The various structures are further described and defined herein.

[0051] Certain antigen binding proteins that are provided may comprise one or more CDRs as described herein (e.g., 1, 2, 3, 4, 5, 6 or more CDRs). In some instances, the antigen binding protein comprises (a) a polypeptide structure and (b) one or more CDRs that are inserted into and/or joined to the polypeptide structure. The polypeptide structure can take a variety of different forms. For example, it can be, or comprise, the framework of a naturally occurring antibody, or fragment or variant thereof, or may be completely synthetic in nature. Examples of various polypeptide structures are further described below.

[0052] An antigen binding protein of the invention is said to “specifically bind” its target antigen when the dissociation equilibrium constant (KD) is $\leq 10^{-8}$ M. The antigen binding protein specifically binds antigen with “high affinity” when the KD is $\leq 5 \times 10^{-9}$ M, and with “very high affinity” when the the KD is $\leq 5 \times 10^{-10}$ M. In one embodiment the antigen binding protein will bind to human IL-23 with a KD of $\leq 5 \times 10^{-12}$ M, and in yet another embodiment it will bind with a $KD \leq 5 \times 10^{-13}$ M. In another embodiment of the invention, the antigen binding protein has a KD of $\leq 5 \times 10^{-12}$ M and an Koff of about $\leq 5 \times 10^{-6}$ 1/s. In another embodiment, the Koff is $\leq 5 \times 10^{-7}$ 1/s.

[0053] Another aspect provides an antigen binding protein having a half-life of at least one day in vitro or in vivo (e.g., when administered to a human subject). In one embodiment, the antigen binding protein has a half-life of at least three days. In another embodiment, the antibody or portion thereof has a half-life of four days or longer. In another embodiment, the antibody or portion thereof has a half-life of eight days or longer. In another embodiment, the antibody or antigen binding portion thereof is derivatized or modified such that it has a longer half-life as compared to the underivatized or unmodified antibody. In another embodiment, the antigen binding protein contains point mutations to increase serum half life, such as described in WIPO Publication No. WO 00/09560.

[0054] In embodiments where the antigen binding protein is used for therapeutic applications, an antigen binding protein can reduce, inhibit, interfere with or modulate one or more biological activities of IL-23, such inducing production of proinflammatory cytokines. IL-23 has many distinct biological effects, which can be measured in many different assays in different cell types; examples of such assays and known and are provided herein.

[0055] Some of the antigen binding proteins that are provided have the structure typically associated with naturally occurring antibodies. The structural units of these antibodies typically comprise one or more tetramers, each composed of two identical couplets of polypeptide chains, though some species of mammals also produce antibodies having only a single heavy chain. In a typical antibody, each pair or couplet includes one full-length “light” chain (in certain embodiments, about 25 kDa) and one full-length “heavy” chain (in certain embodiments, about 50-70 kDa). Each individual immunoglobulin chain is composed of several “immunoglobulin domains”, each consisting of roughly 90 to 110 amino acids and expressing a characteristic folding pattern. These domains are the basic units of which antibody polypeptides are composed. The amino-terminal portion of each chain typically includes a variable region that is responsible for antigen recognition. The carboxy-terminal portion is more conserved evolutionarily than the other end of the chain and is referred to as the “constant region” or “C region”. Human light chains generally are classified as kappa and lambda light chains, and each of these contains one variable region and one constant domain (CL1).z Heavy chains are typically

classified as mu, delta, gamma, alpha, or epsilon chains, and these define the antibody's isotype as IgM, IgD, IgG, IgA, and IgE, respectively. IgG has several subtypes, including, but not limited to, IgG1, IgG2, IgG3, and IgG4. IgM subtypes include IgM, and IgM2. IgA subtypes include IgA1 and IgA2. In humans, the IgA and IgD isotypes contain four heavy chains and four light chains; the IgG and IgE isotypes contain two heavy chains and two light chains; and the IgM isotype contains five heavy chains and five light chains. The heavy chain constant region (CH) typically comprises one or more domains that may be responsible for effector function. The number of heavy chain constant region domains will depend on the isotype. IgG heavy chains, for example, each contains three CH region domains known as CH1, CH2 and CH3. The antibodies that are provided can have any of these isotypes and subtypes, for example, the IL-23 antigen binding protein is of the IgG1, IgG2, or IgG4 subtype. If an IgG4 is desired, it may also be desired to introduce a point mutation (CPSCP->CPPCP) in the hinge region as described in Bloom et al., 1997, Protein Science 6:407) to alleviate a tendency to form intra-H chain disulfide bonds that can lead to heterogeneity in the IgG4 antibodies. Antibodies provided herein that are of one type can be changed to a different type using subclass switching methods. See, e.g., Lantto et al., 2002, Methods Mol. Biol. 178:303-316.

[0056] In full-length light and heavy chains, the variable and constant regions are joined by a "J" region of about twelve or more amino acids, with the heavy chain also including a "D" region of about ten more amino acids. See, e.g., Fundamental Immunology, 2nd ed., Ch. 7 (Paul, W., ed.) 1989, New York: Raven Press. The variable regions of each light/heavy chain pair typically form the antigen binding site.

[0057] Variable Regions

[0058] Various heavy chain and light chain variable regions (or domains) provided herein are depicted in TABLES 1 and 2. Each of these variable regions may be attached, for example, to heavy and light chain constant regions described above. Further, each of the so generated heavy and light chain sequences may be combined to form a complete antigen binding protein structure.

[0059] Provided are antigen binding proteins that contain at least one heavy chain variable region (VH) selected from the group consisting of VH1, VH2, VH3, VH4, VH5, VH6, VH7, VH8, VH9, VH10, VH11, VH12, VH13, VH14, VH15 and VH16 and/or at least one light chain variable region (VL) selected from the group consisting of VL1, VL2, VL3, VL4, VL5, VL6, VL7, VL8, VL9, VL10, VL11, VL12, VL13, VL14, VL15, and VL16 as shown in TABLES 1 and 2 below.

[0060] Each of the heavy chain variable regions listed in TABLE 2 may be combined with any of the light chain variable regions shown in TABLE 1 to form an antigen binding protein. In some instances, the antigen binding protein includes at least one heavy chain variable region and/or one light chain variable region from those listed in TABLES 1 and 2. In some instances, the antigen binding protein includes at least two different heavy chain variable regions and/or light chain variable regions from those listed in TABLES 1 and 2. The various combinations of heavy chain variable regions may be combined with any of the various combinations of light chain variable regions.

[0061] In other instances, the antigen binding protein contains two identical light chain variable regions and/or two identical heavy chain variable regions. As an example, the antigen binding protein may be an antibody or immunologically functional fragment that comprises two light chain variable regions and two heavy chain variable regions in combinations of pairs of light chain variable regions and pairs of heavy chain variable regions as listed in TABLES 1 and 2. Examples of such antigen binding proteins comprising two identical heavy chain and light chain variable regions include: Antibody A VH14/VL14; Antibody B VH9/VL9; Antibody C VH10/VL10; Antibody D VH15/VL15; Antibody E VH1/VL1, Antibody F VH11/VL11; Antibody G VH12/VL12; Antibody H VH13/VL13; Antibody I VH8/VL8; Antibody J VH3/VL3; Antibody K VH7/VL7; Antibody L VH4/VL4; Antibody M VH5/VL5 and Antibody N VH6/VL6.

[0062] Some antigen binding proteins that are provided comprise a heavy chain variable region and/or a light chain variable region comprising a sequence of amino acids that differs from the sequence of a heavy chain variable region and/or a light chain variable region selected from TABLES 1 and 2 at only 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14 or 15 amino acid residues, wherein each such sequence

difference is independently either a deletion, insertion or substitution of one amino acid. The light and heavy chain variable regions, in some antigen binding proteins, comprise sequences of amino acids that have at least 70%, 75%, 80%, 85%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98% or 99% sequence identity to the amino acid sequences provided in TABLES 1 and 2. Still other antigen binding proteins, e.g., antibodies or immunologically functional fragments, also include variant heavy chain region forms and/or variant light chain region forms as described herein.

[0063] The term "identity" refers to a relationship between the sequences of two or more polypeptide molecules or two or more polynucleotides, as determined by aligning and comparing the sequences. "Percent identity" means the percent of identical residues between the amino acids or nucleotides in the compared molecules and is calculated based on the size of the smallest of the molecules being compared.

TABLE-US-00001 TABLE 1 Exemplary Variant Light Chain Region Sequences

FR1	CDRL2	CDRL3	CDRL1	FR4 V.sub.L1	SEQ ID NO:	Region
FR2	CDRL2	CDRL3	CDRL1	FR4 V.sub.L1		
FR3						
QSVLTQPPSVSGAPGQRVTISC	WYQQVPGTAPKLLIY				1	V.sub.L2
GVPDRFSGSKSGTSASLAITGLQAEDEADYYC	FGGGTRLTVL					
QSVLTQPPSVSGAPGQRVTISC	WYQQLPGTAPKLLIY				3	V.sub.L3
GVPDRFSGSKSGTSASLAITGLQAEDEADYYC	FGGGTKLTVL					
QAVLTQPSSLSASPGASASLTC	WYQQKPGSPPQYLLR				4	V.sub.LA
GVPSRFSGSKDASANAGILLISGLQSEDEADYYC	FGGGTKLTVL					
QAVLTQPSSLSASPGASASLTC	WYQQKPGSPPQYLLR				4	V.sub.L5
GVPSRFSGSKDASANAGILLISGLQSEDEADYYC	FGGGTKLTVL					
QPVLTQPPSASASLGASVTLTC	WYQQRPGKGRFVMR				7	V.sub.L6
GIPDRFSVLGSGLNRYLTIKNIQEEDESDYHC	FGTGTKVTVL					
QPVLTQPPSASASLGASVTLTC	WYQQRPGKGRFVMR				9	V.sub.L7
GIPDRFSVLGSGLNRYLTIKNIQEEDESDYHC	FGTGTKVTVL					
QPELTQPPSASASLGASVTLTC	WYQLRPGKGRFVMR				11	V.sub.L8
GIPDRFSVLGSGLNRSLLTIKNIQEEDESDYHC	FGTGTKVTVL					
DIQLTPSPSSVSASVGDRVTITC	WYQQKPGKAPKLLIY				13	V.sub.L9
GVPSRFSGSGSGTDFLTISLQPEDFATYYC	FGGGTKVEIK					
DIQMTQSPSSVSASVGDRVTITC	WYQQKPGKAPSLLIY				15	V.sub.L10
GVPSRFSGSVSGTDFLTISLQPEDFATYYC	FGPGTKVDFK					
DIQMTQSPSSVSASVGDRVTITC	WYQQKPGKAPKLLIY				17	V.sub.L11
GVPSRFSGSGSGTDFLTISLQPEDFATYYC	FGPGTKVDIK					
DSQMTQSPSSVSASVGDRVTITC	WYQQKPGQAPNLLIY				19	V.sub.L12
GVPSRFSGSGSGTEFTLTISLQPEDFATYYC	FGPGTKVDIK					
DIQMTQSPSSVSASVGDRVTITC	WYQQKPGKAPKLLIY				21	V.sub.L13
GVPSRFSGSGSGTDFLTISLQPDFFATYYC	FGGGTKVEIK					
DIQMTQSPSSVSASVGDRVTITC	WYQQKPGKAPKLLIY				23	V.sub.L14
GVPSRFSGSGSGTDFLTISLQPEDFATYYC	FGPGTKVDIK					
DIQLTQSPSSVSASVGDRVTITC	WYQQKPGKAPNLLIY				25	V.sub.L15
GVPSRFSGSGSGTDFLTISLQPADFATYFC	FGPGTKVDVK					
DIQMTQSPSSVSASVGDRVTITC	WYQQKPGKAPKLLIY				27	V.sub.L16
GVPSRFSGSGSGTDFLTISLQPEDFATYYC	FGPGTKVDIK					
DIQMTQSPSSLSASVGDRVTITC	WYQQKPGKAPKRLIY				29	
GVPSRFSGSGSGTEFTLTISLQPEDFATYYC	FGQGTKVEIE					

TABLE-US-00002 TABLE 2 Exemplary Variant Heavy Chain Region Sequences

FR1	CDRH2	CDRH3	CDRH1	FR4 V.sub.H1	SEQ ID NO:	Region
FR2	CDRH2	CDRH3	CDRH1	FR4 V.sub.H1		
FR3						
QVQLVESGGGVVQPGRSLRLSCAASGFTFS	WVRQAPGKGLEWVA				31	V.sub.H2
RFTISRDN SKNTLYLQMN SLRAEDTAVYYCAR	WGQGTMTVSS					
QVQLVESGGGVVQPGRSLRLSCAASGFTFS	WVRQAPGKGLEWVA					

RFTISRDN SKNTLYLQMNSLRAEDTAVYYCAR VWGQGTMTVTVSS SEQ ID NO: 33 V.sub.H3
 QVQLVESGGGVVQPGRSLRLSCAASGFTFS WVRQAPGKGLEWVA
 FTISRDN SKNTLYLQMNSLRAEDTAVYYCAR WGQGT LVTVSS SEQ ID NO: 34 V.sub.H4
 QVQLVESGGGVVQPGRSLRLSCAASGFTFS WVRQAPGKGLEWLS
 RFTISRDN SKNTLYLQMNSLRAEDTAVYYCAR WGQGT LVTVSS SEQ ID NO: 36 V.sub.H5
 EVQLVESGGGLVQP GGS LRLS CAASGFTFS WVRQAPGKGLEWVS
 RFTISRDN AKNSLYLQMNSLRDEDTAVYYCAR WGQGT TVTVSS SEQ ID NO: 38 V.sub.H6
 EVQLVESGGGLVQP GGS LRLS CAASGFTFS WVRQAPGKGLEWVS
 RFTISRDN AKNSLYLQMNSLRDEDTAVYYCAR WGQGT TVTVSS SEQ ID NO: 40 V.sub.H7
 EVQLVESGGGLVQP GGS LRLS CVVSGFTFS WVRQAPGKGLEWVS
 RFTISRDN AKNSLYLQMNSLRDEDTAVYYCAR WGQGT TVTVSS SEQ ID NO: 42 V.sub.H8
 QVQLQESGPGLVKPSETLSLTCTVSGGSIS WIRQPAGKGLEWIG
 RVTMSLDTSKNQFSLRLTSVTAADTAVYYCAR WGQGT TVTVSS SEQ ID NO: 44 V.sub.H9
 QVQLQESGPGLVKPSQTL SLTCTVSGGSIS WIRQHPGKGLEWIG
 RVTISVDTSKNQFSLKLSSVTAADTAVYYCAK WGQGT TVTVSS SEQ ID NO: 46 V.sub.H10
 QVQLQESGPGLVKPSQTL SLTCTVSGGSIN WIRQHP0KGLEWIG
 RVTISVDTSQNQFSLKLSSVTAADTAVYYCAR WGQGT TVTVSS SEQ ID NO: 48 V.sub.H11
 QVQLQESGPGLVKPSQTL SLTCTVSGGSIS WIRQHPGKGLEWIG
 RVTISVDTSKNQFSLKLSSVTAADTAVYYCAR WGQGT TVTVSS SEQ ID NO: 50 V.sub.H12
 QVQLQESGPRLVKPSETLSLTCTVSGDSIS WIRQPPGKGLEWLG
 RVTISIDTSKNQFSLKLSSVTAADTAVYYCTR WGQGT LVTVSS SEQ ID NO: 52 V.sub.H13
 QVQLQESGPGLVKPSQTL SLICTVSGGSIS WIRQHPGKGLEWIG
 RITISVDTSKNQFSLSLSSVTAADTAVYYCAR WGQGT TVTVSS SEQ ID NO: 54 V.sub.H14
 QVQLQESGPGLVKPSQTL SLTCTVSGGSIS WIRQHPGKGLEWIG
 RVIMSVDT SKNQFSLKLSSVTAADTAVYYCAK WGQGT TVTVSS SEQ ID NO: 56 V.sub.H15
 QVQLQESGPGLVKPSQTL SLTCTVSGGSIN WIRQHPGKGLEWIG
 RVTISVDTSKNQFSLKLSSVTAADTAVYYCAR WGQGT TVTVSS SEQ ID NO: 58 V.sub.H16
 QVQLVESGGGVVQPGRSLRLSCAASGFTFS WVRQAPGKGLEWVA
 RFTISRDN SKNTLYLQMNSLRAEDTAVYYCAR WGQGT TVTVSS SEQ ID NO: 60

For these calculations, gaps in alignments (if any) must be addressed by a particular mathematical model or computer program (i.e., an “algorithm”). Methods that can be used to calculate the identity of the aligned nucleic acids or polypeptides include those described in *Computational Molecular Biology*, (Lesk, A. M., ed.), 1988, New York: Oxford University Press; *Biocomputing Informatics and Genome Projects*, (Smith, D. W., ed.), 1993, New York: Academic Press; *Computer Analysis of Sequence Data, Part I*, (Griffin, A. M., and Griffin, H. G., eds.), 1994, New Jersey: Humana Press; von Heinje, G., 1987, *Sequence Analysis in Molecular Biology*, New York: Academic Press; *Sequence Analysis Primer*, (Gribskov, M. and Devereux, J., eds.), 1991, New York: M. Stockton Press; and Carillo et al., 1988, *SIAM J. Applied Math.* 48:1073.

[0064] In calculating percent identity, the sequences being compared are aligned in a way that gives the largest match between the sequences. The computer program used to determine percent identity is the GCG program package, which includes GAP (Devereux et al., 1984, *Nucl. Acid Res.* 12:387; Genetics Computer Group, University of Wisconsin, Madison, Wis.). The computer algorithm GAP is used to align the two polypeptides or polynucleotides for which the percent sequence identity is to be determined. The sequences are aligned for optimal matching of their respective amino acid or nucleotide (the “matched span”, as determined by the algorithm). A gap opening penalty (which is calculated as 3× the average diagonal, wherein the “average diagonal” is the average of the diagonal of the comparison matrix being used; the “diagonal” is the score or number assigned to each perfect amino acid match by the particular comparison matrix) and a gap extension penalty (which is usually 1/10 times the gap opening penalty), as well as a comparison matrix such as PAM 250 or BLOSUM 62 are used in conjunction with the algorithm. In certain embodiments, a standard comparison matrix (see, Dayhoff et al., 1978, *Atlas of Protein Sequence and Structure* 5:345-352 for the PAM 250 comparison matrix; Henikoff et al., 1992, *Proc. Natl. Acad. Sci. U.S.A.* 89:10915-10919 for the BLOSUM 62 comparison matrix) is also used by the algorithm.

[0065] Recommended parameters for determining percent identity for polypeptides or nucleotide sequences using the GAP program are the following: Algorithm: Needleman et al., 1970, *J. Mol. Biol.* 48:443-453; Comparison matrix: BLOSUM 62 from Henikoff et al., 1992, *supra*; Gap Penalty: 12 (but with no penalty for end gaps), Gap Length Penalty: 4, Threshold of Similarity: 0. Certain alignment schemes for aligning two amino acid sequences may result in matching of only a short region of the two sequences and this small aligned region may have very high sequence identity even though there is no significant relationship between the two full-length sequences. Accordingly, the selected alignment method (GAP program) can be adjusted if so desired to result in an alignment that spans at least 50 contiguous amino acids of the target polypeptide.

[0066] The heavy and light chain variable regions disclosed herein include consensus sequences derived from groups of related antigen binding proteins. The amino acid sequences of the heavy and light chain variable regions were analyzed for similarities. Four groups emerged, one group having kappa light chain variable regions, (V.sub.H9/V.sub.L9, V.sub.H10/V.sub.L10, V.sub.H11/V.sub.L11, V.sub.H13/V.sub.L13, V.sub.H14/V.sub.L14 and V.sub.H15/V.sub.L15) and three groups having lambda light chain variable regions: lambda group 1 (V.sub.H5/V.sub.AS, V.sub.H6/V.sub.L6 and V.sub.H7/V.sub.L7), lambda group 2 (V.sub.H3/V.sub.L3 and V.sub.H4/V.sub.L4), and lambda group 3 (V.sub.H1/V.sub.L1 and V.sub.H2/V.sub.L2). Light chain germlines represented include VK1/A30 and VK1/L19. Light chain lambda germlines represented include VL1/1e, VL3/3p, VL5/5c and VL9/9a. Heavy chain germlines represented include VH3/3-30, VH3/3-30.3, VH3/3-33, VH3/3-48, VH4/4-31 and VH4/4-59. As used herein, a "consensus sequence" refers to amino acid sequences having conserved amino acids common among a number of sequences and variable amino acids that vary within given amino acid sequences. Consensus sequences may be determined using standard phylogenetic analyses of the light and heavy chain variable regions corresponding to the IL-23 antigen binding proteins disclosed herein.

[0067] The light chain variable region consensus sequence for the kappa group is
DX.sub.1QX.sub.2TQSPSSVSASVGDRTITCRASQGX.sub.3X.sub.4SX.sub.5WX.sub.6AWYQQKP
GX.sub.7APX.sub.8LLIYAASSLQSGVPSR
FSGSX.sub.9SGTX.sub.10FTLTISLQPX.sub.11DFATYX.sub.12CQQANSFPFTFGPGTKVDX.sub.13
K (SEQ ID NO:30) where X.sub.1 is selected from I or S; X.sub.2 is selected from M or L; X.sub.3 is
selected from G or V and X.sub.4 is selected from S, F or I; X.sub.5 is selected from S or G; X.sub.6 is
selected from F or L; X.sub.7 is selected from K or Q; X.sub.8 is selected from K, N or S; X.sub.9 is
selected from G or V; X.sub.10 is selected from D or E, X.sub.11 is selected from E or A; X.sub.12 is
selected from Y or F; and X.sub.13 is selected from I, V or F.

[0068] The light chain variable region consensus sequence for lambda group 1 is
QPX.sub.1LTQPPSASASLGASVTLTCTLX.sub.2SGYSDYKVDWYQX.sub.3RPGKGPRFVMRVGTGG
X.sub.4VGSKGX.sub.5GI
PDRFSLVLSGLNRX.sub.6LTIKNIQEEDESDYHCGADHSGSX.sub.7NFVYVFGTGTKVTVL (SEQ ID
NO:61) where X.sub.1 is selected from V or E; X.sub.2 is selected from N or S; X.sub.3 is selected
from Q or L and X.sub.4 is selected from I or T; X.sub.5 is selected from D or E; X.sub.6 is selected
from Y or S; and X.sub.7 is selected from S or N.

[0069] The light chain variable region consensus sequence for lambda group 3 is
QSVLTQPPSVSGAPGQRVTISCTGSSSNX.sub.1GAGYDVHWYQQX.sub.2PGTAPKLLIYGSX.sub.3N
RPSGVPDRF SGSKSGTSASLAITGLQAEDEADYYCQSYDSSLSGWVFGGGTX.sub.4RLTVL (SEQ
ID NO:139) where X.sub.1 is selected from T or I; X.sub.2 is selected from V or L; X.sub.3 is selected
from G or N and X.sub.4 is selected from R or K.

[0070] The heavy chain variable region consensus sequence for the kappa group is
QVQLQESGPGLVKPSQTLSTCTVSGGSIX.sub.1SGGYWX.sub.2WIRQHPGKGLEWIGX.sub.3IX.s
ub.4YSGX.sub.5X.sub.6YYNP
SLKSRX.sub.7TX.sub.8SVDTSX.sub.9NQFSLX.sub.10LSSVTAADTAVYYCAX.sub.11X.sub.12RGX.s
ub.13YYGMDVWGQGTTVTVSS (SEQ ID NO:140) where X.sub.1 is selected from N or S; X.sub.2 is
selected from S or T; X.sub.3 is selected from Y or H and X.sub.4 is selected from Y or H; X.sub.5 is
selected from S or N; X.sub.6 is selected from S or T; X.sub.7 is selected from V or I; X.sub.8 is

selected from I or M; X.sub.9 is selected from K or Q; X.sub.10 is selected from K or S, X.sub.11 is selected from R or K; X.sub.12 is selected from D or N; and X.sub.13 is selected from H, F or Y.

[0071] The heavy chain variable region consensus sequence for lambda group 1 is EVQLVESGGGLVQP G G S L R L S C X.sub.1 X.sub.2 S G F T F S X.sub.3 X.sub.4 S M N W V R Q A P G K G L E W V S Y I S S X.sub.5 S S T X.sub.6 Y X.sub.7 A D S V K G R F T I S R D N A K N S L Y L Q M N S L R D E D T A V Y Y C A R R I A A A G X.sub.8 X.sub.9 X.sub.10 Y Y Y A X.sub.11 D V W G Q G T T V T V S S (SEQ ID NO:141) where X.sub.1 is selected from A or V; X.sub.2 is selected from A or V; X.sub.3 is selected from T or S and X.sub.4 is selected from Y or F; X.sub.5 is selected from S or R; X.sub.6 is selected from R or I; X.sub.7 is selected from H, Y or I; X.sub.8 is selected from P or G; X.sub.9 is selected from W or F; X.sub.10 is selected from G or H and X.sub.11 is selected from M or L.

[0072] The heavy chain variable region consensus sequence for lambda group 2 is QVQLVESGGGVVQPGRSLRLS CAASGFTFSSYX.sub.1 MHWVRQAPGKGLEWX.sub.2 X.sub.3 VISX.sub.4 DGSX.sub.5 KY YAD S V K G R F T I S R D N S K N T L Y L Q M N S L R A E D T A V Y Y C A R E R T T L S G S Y F D Y W G Q G T L V T V S S (SEQ ID NO:142) where X.sub.1 is selected from G or A; X.sub.2 is selected from V or L; X.sub.3 is selected from A or S and X.sub.4 is selected from F or H and X.sub.5 is selected from L or I.

[0073] The heavy chain variable region consensus sequence for lambda group 3 is QVQLVESGGGVVQPGRSLRLS CAASGFTFSSYGMHWVRQAPGKGLEWVAVIWDGSNX.sub.1 YYA D S V K G R F T I S R D N S K N T L Y L Q M N S L R A E D T A V Y Y C A R D R G Y X.sub.2 S S W Y P D A F D I W G Q G T M V T V S S (SEQ ID NO: 143) where X.sub.1 is selected from E or K and X.sub.2 is selected from T or S.

[0074] Complementarity Determining Regions

[0075] Complementarity determining regions or “CDRs” are embedded within a framework in the heavy and light chain variable regions where they constitute the regions responsible for antigen binding and recognition. Variable domains of immunoglobulin chains of the same species, for example, generally exhibit a similar overall structure; comprising relatively conserved framework regions (FR) joined by hypervariable CDR regions. An antigen binding protein can have 1, 2, 3, 4, 5, 6 or more CDRs. The variable regions discussed above, for example, typically comprise three CDRs. The CDRs from heavy chain variable regions and light chain variable regions are typically aligned by the framework regions to form a structure that binds specifically on a target antigen (e.g., IL-23). From N-terminal to C-terminal, naturally-occurring light and heavy chain variable regions both typically conform to the following order of these elements: FR1, CDR1, FR2, CDR2, FR3, CDR3 and FR4. The CDR and FR regions of exemplary light chain variable domains and heavy chain variable domains are highlighted in TABLES 1 and 2. It is recognized that the boundaries of the CDR and FR regions can vary from those highlighted. Numbering systems have been devised for assigning numbers to amino acids that occupy positions in each of these domains. Complementarity determining regions and framework regions of a given antigen binding protein may be identified using these systems. Numbering systems are defined in Kabat et al., *Sequences of Proteins of Immunological Interest*, 5.sup.th Ed., US Dept. of Health and Human Services, PHS, NIH, NIH Publication No. 91-3242, 1991, or Chothia & Lesk, 1987, *J. Mol. Biol.* 196:901-917; Chothia et al., 1989, *Nature* 342:878-883. Other numbering systems for the amino acids in immunoglobulin chains include IMGT® (the international ImMunoGeneTics information system; Lefranc et al, *Dev. Comp. Immunol.* 2005, 29:185-203); and AHo (Honegger and Pluckthun, *J. Mol. Biol.* 2001, 309(3):657-670). The CDRs provided herein may not only be used to define the antigen binding domain of a traditional antibody structure, but may be embedded in a variety of other polypeptide structures, as described herein.

[0076] The antigen binding proteins disclosed herein are polypeptides into which one or more CDRs may be grafted, inserted, embedded and/or joined. An antigen binding protein can have, for example, one heavy chain CDR1 (“CDRH1”), and/or one heavy chain CDR2 (“CDRH2”), and/or one heavy chain CDR3 (“CDRH3”), and/or one light chain CDR1 (“CDRL1”), and/or one light chain CDR2 (“CDRL2”), and/or one light chain CDR3 (“CDRL3”). Some antigen binding proteins include both a CDRH3 and a

CDRL3. Specific embodiments generally utilize combinations of CDRs that are non-repetitive, e.g., antigen binding proteins are generally not made with two CDRH2 regions in one variable heavy chain region, etc. Antigen binding proteins may comprise one or more amino acid sequences that are identical to or that differ from the amino acid sequences of one or more of the CDRs presented in TABLE 3 at only 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14 or 15 amino acid residues, wherein each such sequence difference is independently either a deletion, insertion or substitution of one amino acid. The CDRs in some antigen binding proteins comprise sequences of amino acids that have at least 80%, 85%, 90%, 91%, 92, 93%, 94%, 95%, 96%, 97%, 98%, or 99% sequence identity to CDRs sequence listed in TABLE 3. In some antigen binding proteins, the CDRs are embedded into a "framework" region, which orients the CDR(s) such that the proper antigen binding properties of the CDR(s) is achieved.

TABLE-US-00003 TABLE 3 Exemplary CDRH and CDRL Sequences Exemplary CDRL Sequences
CDRL1 CDRL2 CDRL3 TGSSSNTGAGYDVH GSGNRPS QSYDSSLGWSV SEQ ID NO: 62
SEQ ID NO: 63 SEQ ID NO: 64 TGSSNIGAGYDVH GSNRPS MIWHSSASV SEQ ID NO: 65
SEQ ID NO: 66 SEQ ID NO: 67 TLRSGINVTYRIY YKSDSDKQQGS GADHGSGSNFVYV SEQ ID NO: 68
SEQ ID NO: 69 SEQ ID NO: 70 TLNSGYSDYKV VGTGGIVGSKGD GADHGSGNNFVYV SEQ ID NO: 71
SEQ ID NO: 72 SEQ ID NO: 73 TLSSGYSDYKV VGTGGIVGSKGE QQANSFPPT SEQ ID NO: 74
SEQ ID NO: 75 SEQ ID NO: 76 RASQGFSGWLA VGTGGTVGSKGE QQATSFPLT SEQ ID NO: 77
SEQ ID NO: 78 SEQ ID NO: 79 RASQVISSWLA AASSLQS QQADSFPT SEQ ID NO: 80
SEQ ID NO: 81 SEQ ID NO: 82 RASQVISSWFA LQHNSYPPT SEQ ID NO: 83
SEQ ID NO: 84 RASQGSSSWFA SEQ ID NO: 85 RASQGISSWFA SEQ ID NO: 86 RAGQVISSWLA
SEQ ID NO: 87 RASQGIAGWLA SEQ ID NO: 88 RASQGIRNDLG SEQ ID NO: 89
Exemplary CDRH Sequences CDRH1 CDRH2 CDRH3 SYGMH VIWYDGSNEYADSVKGRG
YTSWYPDAFDI SEQ ID NO: 91 SEQ ID NO: 92 SEQ ID NO: 93 SYAMH VIWYDGSNKY
ADSVKGRGYSSWYPDAFDI SEQ ID NO: 94 SEQ ID NO: 95 SEQ ID NO: 96 TYSMN
VISFDGSLKYADSVKGERTTLSGSYFDY SEQ ID NO: 97 SEQ ID NO: 98 SEQ ID NO: 99
SYSMN VISHDGSIKYADSVKGRIAAAGGFHYALDV SEQ ID NO: 100 SEQ ID NO: 101
SEQ ID NO: 102 SFSMN YISSRSSTIYADSVKGRIAAAGPWGYAMDV SEQ ID NO: 103
SEQ ID NO: 104 SEQ ID NO: 105 SGGYYWT YISSSSTRYHADSVKGRNGYYYGMDV
SEQ ID NO: 106 SEQ ID NO: 107 SEQ ID NO: 108 SGGYYWS YISSRSSTIYADSVKGR
NGFYGMDV SEQ ID NO: 109 SEQ ID NO: 110 SEQ ID NO: 111 SYFWS YIYSGNTY
NPSLKS DRGHYGMVDV SEQ ID NO: 112 SEQ ID NO: 113 SEQ ID NO: 114 TYYWS
HIHSGNTYNNPSLKS DRGSYYGSDY SEQ ID NO: 115 SEQ ID NO: 116 SEQ ID NO: 117
YIYSGSTYNNPSLKS DRGYGVDV SEQ ID NO: 118 SEQ ID NO: 119 YIYSGSSYNNPSLKS
ENTVTIYNYGMDV SEQ ID NO: 120 SEQ ID NO: 6 YIYSGSTNYNNPSLKS SEQ ID NO: 121
LIYTSGSTNYNNPSLKS SEQ ID NO: 122 LIWYDGSNKYADSVKGRG SEQ ID NO: 90

[0077] Provided herein are CDR1 regions comprising amino acid residues 23-34 of SEQ ID NOs: 7 and 11; amino acid residues 24-34 of SEQ ID NOs: 9, 13, 15, 17, 19, 21, 23, 25, 27 and 29; amino acid residues 23-36 of SEQ ID NOs: 1, 3 and 4; amino acid residues 31-35 of SEQ ID NOs: 31, 33, 34, 38, 40, 44, 52 and 60 and amino acid residues 31-37 or SEQ ID NOs: 46, 48, 50, 54, 56 and 58.

[0078] CDR2 regions are provided comprising amino acid residues 50-56 of SEQ ID NOs: 9, 13, 15, 17, 19, 21, 23, 25, 27 and 29; amino acid residues 50-61 of SEQ ID NOs: 7 and 11; amino acid residues 52-62 of SEQ ID NO: 4; amino acid residues 50-65 of SEQ ID NOs: 31, 33, 44 and 52; amino acid residues 50-66 of SEQ ID NOs: 36, 38, 40, 42 and 60; amino acid residues 52-58 of SEQ ID NOs: 1 and 3 and amino acid residues 52-67 of SEQ ID NOs: 46, 48, 50, 54, 56 and 58.

[0079] CDR3 regions comprising amino acid residues 89-97 of SEQ ID NOs: 13, 15, 17, 19, 21, 23, 25, 27 and 29; amino acid residues 91-101 of SEQ ID NOs: 1 and 3; amino acid residues 94-106 of SEQ ID NOs: 7, 9 and 11; amino acid residues 98-107 of SEQ ID NOs: 44 and 52; amino acid residues 97-105 of SEQ ID NO: 4; amino acid residues 99-110 of SEQ ID NOs: 34 and 36; amino acid residues 99-112 of SEQ ID NO: 112; amino acid residues 99-113 of SEQ ID NOs: 31 and 33; amino

acid residues 99-114 of SEQ ID NOs: 38, 40 and 42; amino acid residues 100-109 of SEQ ID NOs: 46, 48, 54, 56 and 58; and amino acid residues 101-019 of SEQ ID NO; 50; are also provided.

[0080] The CDRs disclosed herein include consensus sequences derived from groups of related sequences. As described previously, four groups of variable region sequences were identified, a kappa group and three lambda groups. The CDRL1 consensus sequence from the kappa group consists of RASQX.sub.1X.sub.2SX.sub.3WX.sub.4A (SEQ ID NO:123) where X.sub.1 is selected from G or V; X.sub.2 is selected from I, F or S; X.sub.3 is selected from S or G and X.sub.4 is selected from F or L. The CDRL1 consensus sequence from lambda group 1 consists of TLX.sub.1SGYSDYKVD (SEQ ID NO:124) wherein X.sub.1 is selected from N or S. The CDRL1 consensus sequences from lambda group 3 consists of TGSSSNX.sub.1GAGYDVH (SEQ ID NO:125) wherein X.sub.1 is selected from I or T.

[0081] The CDRL2 consensus sequence from lambda group 1 consists of VGTGGX.sub.1VGSKGX.sub.2 (SEQ ID NO: 126) wherein X.sub.1 is selected from I or T and X.sub.2 is selected from D or E. The CDRL2 consensus sequence from lambda group 3 consists of GSX.sub.1NRPS (SEQ ID NO:127) wherein X.sub.1 is selected from N or G.

[0082] The CDRL3 consensus sequences include GADHGSGX.sub.1NFVYV (SEQ IDN NO:128) wherein X.sub.1 is S or N.

[0083] The CDRH1 consensus sequence from the kappa group consists of SGGYYWX.sub.1 (SEQ ID NO:129) wherein X.sub.1 is selected from S or T. The CDRH1 consensus sequence from lambda group 1 consists of X.sub.1X.sub.2SMN (SEQ ID NO:131) wherein X.sub.1 is selected from S or T and X.sub.2 is selected from Y or F. The CDRH1 consensus sequence from lambda group 2 consists of SYX.sub.1MH (SEQ ID NO:130), wherein X.sub.1 is selected from G or A.

[0084] The CDRH2 consensus sequence from the kappa group consists of X.sub.1IX.sub.2YSGX.sub.3X.sub.4YYNPSLKS (SEQ ID NO:132) wherein X.sub.1 is selected from Y or H; X.sub.2 is selected from Y or H; X.sub.3 is selected from S or N and X.sub.4 is selected from T or S. The consensus sequence from lambda group 1 consists of YISSX.sub.1SSTX.sub.2YX.sub.3ADSVKG (SEQ ID NO:134) wherein X.sub.1 is selected from R or S, X.sub.2 is selected from I or R, X.sub.3 is selected from I, H or Y. The consensus sequence from lambda group 2 consists of VISX.sub.1DGSX.sub.2KYYADSVKG (SEQ ID NO:133) wherein X.sub.1 is F or H and X.sub.2 is L or T. The CDRH2 consensus sequence from lambda group 3 consists of VIWYDGSNX.sub.1YYADSVKG (SEQ ID NO:135) wherein X.sub.1 is selected from K or E.

[0085] The CDRH3 consensus sequence from the kappa group consists of X.sub.1RGX.sub.2YYGMDV (SEQ ID NO:136) wherein X.sub.1 is selected from N or D and X.sub.2 is selected from H, Y or F. The CDRH3 consensus sequence from lambda group 1 consists of RIAAAGX.sub.1X.sub.2X.sub.3YYYAX.sub.4DV (SEQ ID NO:137) wherein X.sub.1 is selected from G or P; X.sub.2 is selected from F or W; X.sub.3 is selected from H or G and X.sub.4 is selected from L and M. The CDRH3 consensus sequence from lambda group 3 consists of DRGYX.sub.1SSWYPDAFDI (SEQ ID NO:138) wherein X.sub.1 is selected from S or T.

[0086] Monoclonal Antibodies

[0087] The antigen binding proteins that are provided include monoclonal antibodies that bind to IL-23. Monoclonal antibodies may be produced using any technique known in the art, e.g., by immortalizing spleen cells harvested from the transgenic animal after completion of the immunization schedule. The spleen cells can be immortalized using any technique known in the art, e.g., by fusing them with myeloma cells to produce hybridomas. Myeloma cells for use in hybridoma-producing fusion procedures preferably are non-antibody-producing, have high fusion efficiency, and enzyme deficiencies that render them incapable of growing in certain selective media which support the growth of only the desired fused cells (hybridomas). Examples of suitable cell lines for use in mouse fusions include Sp-20, P3-X63/Ag8, P3-X63-Ag8.653, NS1/1.Ag 4 1, Sp210-Ag14, FO, NSO/U, MPC-11, MPC11-X45-GTG 1.7 and S194/5XXO Bul; examples of cell lines used in rat fusions include

R210.RCY3, Y3-Ag 1.2.3, IR983F and 4B210. Other cell lines useful for cell fusions are U-266, GM1500-GRG2, LICR-LON-HMy2 and UC729-6.

[0088] In some instances, a hybridoma cell line is produced by immunizing an animal (e.g., a transgenic animal having human immunoglobulin sequences) with an IL-23 immunogen; harvesting spleen cells from the immunized animal; fusing the harvested spleen cells to a myeloma cell line, thereby generating hybridoma cells; establishing hybridoma cell lines from the hybridoma cells, and identifying a hybridoma cell line that produces an antibody that binds an IL-23 polypeptide while sparing IL-12. Such hybridoma cell lines, and anti-IL-23 monoclonal antibodies produced by them, are aspects of the present application.

[0089] Monoclonal antibodies secreted by a hybridoma cell line can be purified using any technique known in the art. Hybridomas or mAbs may be further screened to identify mAbs with particular properties, such as the ability to inhibit IL-23-induced activity.

[0090] Chimeric and Humanized Antibodies

[0091] Chimeric and humanized antibodies based upon the foregoing sequences are also provided. Monoclonal antibodies for use as therapeutic agents may be modified in various ways prior to use. One example is a chimeric antibody, which is an antibody composed of protein segments from different antibodies that are covalently joined to produce functional immunoglobulin light or heavy chains or immunologically functional portions thereof. Generally, a portion of the heavy chain and/or light chain is identical with or homologous to a corresponding sequence in antibodies derived from a particular species or belonging to a particular antibody class or subclass, while the remainder of the chain(s) is/are identical with or homologous to a corresponding sequence in antibodies derived from another species or belonging to another antibody class or subclass. For methods relating to chimeric antibodies, see, for example, U.S. Pat. No. 4,816,567; and Morrison et al., 1985, *Proc. Natl. Acad. Sci. USA* 81:6851-6855. CDR grafting is described, for example, in U.S. Pat. Nos. 6,180,370, 5,693,762, 5,693,761, 5,585,089, and 5,530,101.

[0092] One useful type of chimeric antibody is a "humanized" antibody. Generally, a humanized antibody is produced from a monoclonal antibody raised initially in a non-human animal. Certain amino acid residues in this monoclonal antibody, typically from non-antigen recognizing portions of the antibody, are modified to be homologous to corresponding residues in a human antibody of corresponding isotype. Humanization can be performed, for example, using various methods by substituting at least a portion of a rodent variable region for the corresponding regions of a human antibody (see, e.g., U.S. Pat. Nos. 5,585,089, and 5,693,762; Jones et al., 1986, *Nature* 321:522-525; Riechmann et al., 1988, *Nature* 332:323-27; Verhoeyen et al., 1988, *Science* 239:1534-1536),

[0093] In certain embodiments, constant regions from species other than human can be used along with the human variable region(s) to produce hybrid antibodies.

[0094] Fully Human Antibodies

[0095] Fully human antibodies are also provided. Methods are available for making fully human antibodies specific for a given antigen without exposing human beings to the antigen ("fully human antibodies"). One specific means provided for implementing the production of fully human antibodies is the "humanization" of the mouse humoral immune system. Introduction of human immunoglobulin (Ig) loci into mice in which the endogenous Ig genes have been inactivated is one means of producing fully human monoclonal antibodies (mAbs) in mouse, an animal that can be immunized with any desirable antigen. Using fully human antibodies can minimize the immunogenic and allergic responses that can sometimes be caused by administering mouse or mouse-derivatized mAbs to humans as therapeutic agents.

[0096] Fully human antibodies can be produced by immunizing transgenic animals (usually mice) that are capable of producing a repertoire of human antibodies in the absence of endogenous immunoglobulin production. Antigens for this purpose typically have six or more contiguous amino

acids, and optionally are conjugated to a carrier, such as a hapten. See, e.g., Jakobovits et al., 1993, *Proc. Natl. Acad. Sci. USA* 90:2551-2555; Jakobovits et al., 1993, *Nature* 362:255-258; and Bruggemann et al., 1993, *Year in Immunol.* 7:33. In one example of such a method, transgenic animals are produced by incapacitating the endogenous mouse immunoglobulin loci encoding the mouse heavy and light immunoglobulin chains therein, and inserting into the mouse genome large fragments of human genome DNA containing loci that encode human heavy and light chain proteins. Partially modified animals, which have less than the full complement of human immunoglobulin loci, are then cross-bred to obtain an animal having all of the desired immune system modifications. When administered an immunogen, these transgenic animals produce antibodies that are immunospecific for the immunogen but have human rather than murine amino acid sequences, including the variable regions. For further details of such methods, see, for example, WIPO patent publications WO96/33735 and WO94/02602. Additional methods relating to transgenic mice for making human antibodies are described in U.S. Pat. Nos. 5,545,807; 6,713,610; 6,673,986; 6,162,963; 5,545,807; 6,300,129; 6,255,458; 5,877,397; 5,874,299 and 5,545,806; in WIPO patent publications WO91/10741, WO90/04036, and in EP 546073B1 and EP 546073A1.

[0097] The transgenic mice described above contain a human immunoglobulin gene minilocus that encodes unrearranged human heavy ([mu] and [gamma]) and [kappa] light chain immunoglobulin sequences, together with targeted mutations that inactivate the endogenous [mu] and [kappa] chain loci (Lonberg et al., 1994, *Nature* 368:856-859). Accordingly, the mice exhibit reduced expression of mouse IgM or [kappa] and in response to immunization, and the introduced human heavy and light chain transgenes undergo class switching and somatic mutation to generate high affinity human IgG [kappa] monoclonal antibodies (Lonberg et al., supra.; Lonberg and Huszar, 1995, *Intern. Rev. Immunol.* 13: 65-93; Harding and Lonberg, 1995, *Ann. N.Y Acad. Sci.* 764:536-546). The preparation of such mice is described in detail in Taylor et al., 1992, *Nucleic Acids Research* 20:6287-6295; Chen et al., 1993, *International Immunology* 5:647-656; Tuaille et al., 1994, *J. Immunol.* 152:2912-2920; Lonberg et al., 1994, *Nature* 368:856-859; Lonberg, 1994, *Handbook of Exp. Pharmacology* 113:49-101; Taylor et al., 1994, *International Immunology* 6:579-591; Lonberg and Huszar, 1995, *Intern. Rev. Immunol.* 13:65-93; Harding and Lonberg, 1995, *Ann. N.Y Acad. Sci.* 764:536-546; Fishwild et al., 1996, *Nature Biotechnology* 14:845-85. See, further U.S. Pat. Nos. 5,545,806; 5,569,825; 5,625,126; 5,633,425; 5,789,650; 5,877,397; 5,661,016; 5,814,318; 5,874,299; and 5,770,429; as well as U.S. Pat. No. 5,545,807; WIPO Publication Nos. WO 93/1227; WO 92/22646; and WO 92/03918. Technologies utilized for producing human antibodies in these transgenic mice are disclosed also in WIPO Publication No. WO 98/24893, and Mendez et al., 1997, *Nature Genetics* 15:146-156. For example, the HCo7 and HCo12 transgenic mice strains can be used to generate anti-IL-23 antibodies.

[0098] Using hybridoma technology, antigen-specific human mAbs with the desired specificity can be produced and selected from the transgenic mice such as those described above. Such antibodies may be cloned and expressed using a suitable vector and host cell, or the antibodies can be harvested from cultured hybridoma cells.

[0099] Fully human antibodies can also be derived from phage-display libraries (such as disclosed in Hoogenboom et al., 1991, *J. Mol. Biol.* 227:381; Marks et al., 1991, *J. Mol. Biol.* 222:581; WIPO Publication No. WO 99/10494). Phage display techniques mimic immune selection through the display of antibody repertoires on the surface of filamentous bacteriophage, and subsequent selection of phage by their binding to an antigen of choice.

[0100] Bispecific or Bifunctional Antigen Binding Proteins

[0101] A "bispecific," "dual-specific" or "bifunctional" antigen binding protein or antibody is a hybrid antigen binding protein or antibody, respectively, having two different antigen binding sites, such as one or more CDRs or one or more variable regions as described above. In some instances they are an artificial hybrid antibody having two different heavy/light chain pairs and two different binding sites. Multispecific antigen binding protein or "multispecific antibody" is one that targets more than one antigen or epitope. Bispecific antigen binding proteins and antibodies are a species of multispecific antigen binding protein antibody and may be produced by a variety of methods including, but not

limited to, fusion of hybridomas or linking of Fab' fragments. See, e.g., Songsivilai and Lachmann, 1990, *Clin. Exp. Immunol.* 79:315-321; Kostelny et al., 1992, *J. Immunol.* 148:1547-1553.

[0102] Immunological Fragments

[0103] Antigen binding proteins also include immunological fragments of an antibody (e.g., a Fab, a Fab', a F(ab').sub.2, or a scFv). A "Fab fragment" is comprised one light chain (the light chain variable region (V.sub.L) and its corresponding constant domain (C.sub.L)) and one heavy chain (the heavy chain variable region (V.sub.H) and first constant domain (C.sub.H1)). The heavy chain of a Fab molecule cannot form a disulfide bond with another heavy chain molecule. A "Fab' fragment" contains one light chain and a portion of one heavy chain that also contains the region between the C.sub.H1 and C.sub.H2 domains, such that an interchain disulfide bond can be formed between the two heavy chains of two Fab' fragments to form an F(ab').sub.2 molecule. A "F(ab').sub.2 fragment" thus is composed of two Fab' fragments that are held together by a disulfide bond between the two heavy chains. A "Fv fragment" consists of the variable light chain region and variable heavy chain region of a single arm of an antibody. Single-chain antibodies "scFv" are Fv molecules in which the heavy and light chain variable regions have been connected by a flexible linker to form a single polypeptide chain, which forms an antigen binding region. Single chain antibodies are discussed in detail in WIPO Publication No. WO 88/01649, U.S. Pat. Nos. 4,946,778 and 5,260,203; Bird, 1988, *Science* 242:423; Huston et al., 1988, *Proc. Natl. Acad. Sci. U.S.A.* 85:5879; Ward et al., 1989, *Nature* 334:544, de Graaf et al., 2002, *Methods Mol Biol.* 178:379-387; Kortt et al., 1997, *Prot. Eng.* 10:423; Kortt et al., 2001, *Biomol. Eng.* 18:95-108 and Kriangkum et al., 2001, *Biomol. Eng.* 18:31-40. A "Fc" region contains two heavy chain fragments comprising the C.sub.H1 and C.sub.H2 domains of an antibody. The two heavy chain fragments are held together by two or more disulfide bonds and by hydrophobic interactions of the C.sub.H3 domains.

[0104] Also included are domain antibodies, immunologically functional immunoglobulin fragments containing only the variable region of a heavy chain or the variable region of a light chain. In some instances, two or more V.sub.H regions are covalently joined with a peptide linker to create a bivalent domain antibody. The two V.sub.H regions of a bivalent domain antibody may target the same or different antigens. Diabodies are bivalent antibodies comprising two polypeptide chains, wherein each polypeptide chain comprises V.sub.H and V.sub.L domains joined by a linker that is too short to allow for pairing between two domains on the same chain, thus allowing each domain to pair with a complementary domain on another polypeptide chain (see, e.g., Holliger et al., *Proc. Natl. Acad. Sci. USA* 90:6444-48, 1993 and Poljak et al., *Structure* 2:1121-23, 1994). Similarly, tribodies and tetrabodies are antibodies comprising three and four polypeptide chains, respectively, and forming three and four antigen binding sites, respectively, which can be the same or different. Maxibodies comprise bivalent scFvs covalently attached to the Fc region of IgG.sub.1, (see, e.g., Fredericks et al, 2004, *Protein Engineering, Design & Selection*, 17:95-106; Powers et al., 2001, *Journal of Immunological Methods*, 251:123-135; Shu et al., 1993, *Proc. Natl. Acad. Sci. USA* 90:7995-7999; Hayden et al., 1994, *Therapeutic Immunology* 1:3-15).

[0105] Various Other Forms

[0106] Also provided are variant forms of the antigen binding proteins disclosed above, some of the antigen binding proteins having, for example, one or more conservative amino acid substitutions in one or more of the heavy or light chains, variable regions or CDRs listed in TABLES 1 and 2.

[0107] Naturally-occurring amino acids may be divided into classes based on common side chain properties: hydrophobic (norleucine, Met, Ala, Val, Leu, Ile); neutral hydrophilic (Cys, Ser, Thr, Asn, Gln); acidic (Asp, Glu); basic (His, Lys, Arg); residues that influence chain orientation (Gly, Pro); and aromatic (Trp, Tyr, Phe).

[0108] Conservative amino acid substitutions may involve exchange of a member of one of these classes with another member of the same class. Conservative amino acid substitutions may encompass non-naturally occurring amino acid residues, which are typically incorporated by chemical peptide synthesis rather than by synthesis in biological systems. These include peptidomimetics and

may be changed without destroying activity by targeting regions not believed to be important for activity. The skilled artisan also will be able to identify residues and portions of the molecules that are conserved among similar polypeptides. In further embodiments, even areas that may be important for biological activity or for structure may be subject to conservative amino acid substitutions without destroying the biological activity or without adversely affecting the polypeptide structure.

[0116] Additionally, one skilled in the art can review structure-function studies identifying residues in similar polypeptides that are important for activity or structure. In view of such a comparison, one can predict the importance of amino acid residues in a protein that correspond to amino acid residues important for activity or structure in similar proteins. One skilled in the art may opt for chemically similar amino acid substitutions for such predicted important amino acid residues.

[0117] One skilled in the art can also analyze the 3-dimensional structure and amino acid sequence in relation to that structure in similar polypeptides. In view of such information, one skilled in the art may predict the alignment of amino acid residues of an antibody with respect to its three dimensional structure. One skilled in the art may choose not to make radical changes to amino acid residues predicted to be on the surface of the protein, since such residues may be involved in important interactions with other molecules. Moreover, one skilled in the art may generate test variants containing a single amino acid substitution at each desired amino acid residue. These variants can then be screened using assays for IL-23 activity, (see examples below) thus yielding information regarding which amino acids can be changed and which must not be changed. In other words, based on information gathered from such routine experiments, one skilled in the art can readily determine the amino acid positions where further substitutions should be avoided either alone or in combination with other mutations.

[0118] A number of scientific publications have been devoted to the prediction of secondary structure. See, Moulton, 1996, *Curr. Op. in Biotech.* 7:422-427; Chou et al., 1974, *Biochem.* 13:222-245; Chou et al., 1974, *Biochemistry* 113:211-222; Chou et al., 1978, *Adv. Enzymol. Relat. Areas Mol. Biol.* 47:45-148; Chou et al., 1979, *Ann. Rev. Biochem.* 47:251-276; and Chou et al., 1979, *Biophys. J.* 26:367-384. Moreover, computer programs are currently available to assist with predicting secondary structure. One method of predicting secondary structure is based upon homology modeling. For example, two polypeptides or proteins that have a sequence identity of greater than 30%, or similarity greater than 40% often have similar structural topologies. The recent growth of the protein structural database (PDB) has provided enhanced predictability of secondary structure, including the potential number of folds within a polypeptide's or protein's structure. See, Holm et al., 1999, *Nucl. Acid. Res.* 27:244-247. It has been suggested (Brenner et al., 1997, *Curr. Op. Struct. Biol.* 7:369-376) that there are a limited number of folds in a given polypeptide or protein and that once a critical number of structures have been resolved, structural prediction will become dramatically more accurate.

[0119] Additional methods of predicting secondary structure include "threading" (Jones, 1997, *Curr. Opin. Struct. Biol.* 7:377-387; Sippl et al., 1996, *Structure* 4:15-19), "profile analysis" (Bowie et al., 1991, *Science* 253:164-170; Gribskov et al., 1990, *Meth. Enzym.* 183:146-159; Gribskov et al., 1987, *Proc. Nat. Acad. Sci.* 84:4355-4358), and "evolutionary linkage" (See, Holm, 1999, supra; and Brenner, 1997, supra).

[0120] In some embodiments, amino acid substitutions are made that: (1) reduce susceptibility to proteolysis, (2) reduce susceptibility to oxidation, (3) alter binding affinity for forming protein complexes, (4) alter ligand or antigen binding affinities, and/or (4) confer or modify other physicochemical or functional properties on such polypeptides, such as maintaining the structure of the molecular backbone in the area of the substitution, for example, as a sheet or helical conformation; maintaining or altering the charge or hydrophobicity of the molecule at the target site, or maintaining or altering the bulkiness of a side chain.

[0121] For example, single or multiple amino acid substitutions (in certain embodiments, conservative amino acid substitutions) may be made in the naturally-occurring sequence. Substitutions can be made in that portion of the antibody that lies outside the domain(s) forming intermolecular contacts). In such embodiments, conservative amino acid substitutions can be used that do not substantially

change the structural characteristics of the parent sequence (e.g., one or more replacement amino acids that do not disrupt the secondary structure that characterizes the parent or native antigen binding protein). Examples of art-recognized polypeptide secondary and tertiary structures are described in *Proteins, Structures and Molecular Principles* (Creighton, Ed.), 1984, W. H. New York: Freeman and Company; *Introduction to Protein Structure* (Branden and Tooze, eds.), 1991, New York: Garland Publishing; and Thornton et al., 1991, *Nature* 354:105.

[0122] Additional variants include cysteine variants wherein one or more cysteine residues in the parent or native amino acid sequence are deleted from or substituted with another amino acid (e.g., serine). Cysteine variants are useful, inter alia when antibodies (for example) must be refolded into a biologically active conformation. Cysteine variants may have fewer cysteine residues than the native protein, and typically have an even number to minimize interactions resulting from unpaired cysteines.

[0123] The heavy and light chain variable region and CDRs that are disclosed can be used to prepare antigen binding proteins that contain an antigen binding region that can specifically bind to an IL-23 polypeptide. "Antigen binding region" means a protein, or a portion of a protein, that specifically binds a specified antigen, such as the region that contains the amino acid residues that interact with an antigen and confer on the antigen binding protein its specificity and affinity for the target antigen. An antigen binding region may include one or more CDRs and certain antigen binding regions also include one or more "framework" regions. For example, one or more of the CDRs listed in TABLE 3 can be incorporated into a molecule (e.g., a polypeptide) covalently or noncovalently to make an immunoadhesion. An immunoadhesion may incorporate the CDR(s) as part of a larger polypeptide chain, may covalently link the CDR(s) to another polypeptide chain, or may incorporate the CDR(s) noncovalently. The CDR(s) enable the immunoadhesion to bind specifically to a particular antigen of interest (e.g., an IL-23 polypeptide).

[0124] Other antigen binding proteins include mimetics (e.g., "peptide mimetics" or "peptidomimetics") based upon the variable regions and CDRs that are described herein. These analogs can be peptides, non-peptides or combinations of peptide and non-peptide regions. Fauchere, 1986, *Adv. Drug Res.* 15:29; Veber and Freidinger, 1985, *TINS* p. 392; and Evans et al., 1987, *J. Med. Chem.* 30:1229. Peptide mimetics that are structurally similar to therapeutically useful peptides may be used to produce a similar therapeutic or prophylactic effect. Such compounds are often developed with the aid of computerized molecular modeling. Generally, peptidomimetics are proteins that are structurally similar to an antigen binding protein displaying a desired biological activity, such as the ability to bind IL-23, but peptidomimetics have one or more peptide linkages optionally replaced by a linkage selected from, for example: —CH.sub.2NH—, —CH.sub.2S—, —CH.sub.2—CH.sub.2—, —CH—CH— (cis and trans), —COCH.sub.2—, —CH(OH)CH.sub.2—, and —CH.sub.2SO—, by methods well known in the art. Systematic substitution of one or more amino acids of a consensus sequence with a D-amino acid of the same type (e.g., D-lysine in place of L-lysine) may be used in certain embodiments to generate more stable proteins. In addition, constrained peptides comprising a consensus sequence or a substantially identical consensus sequence variation may be generated by methods known in the art (Rizo and Gierasch, 1992, *Ann. Rev. Biochem.* 61:387), for example, by adding internal cysteine residues capable of forming intramolecular disulfide bridges which cyclize the peptide.

[0125] Derivatives of the antigen binding proteins that are described herein are also provided. The derivatized antigen binding proteins can comprise any molecule or substance that imparts a desired property to the antigen binding protein or fragment, such as increased half-life in a particular use. The derivatized antigen binding protein can comprise, for example, a detectable (or labeling) moiety (e.g., a radioactive, colorimetric, antigenic or enzymatic molecule, a detectable bead (such as a magnetic or electrodense (e.g., gold) bead), or a molecule that binds to another molecule (e.g., biotin or Streptavidin)), a therapeutic or diagnostic moiety (e.g., a radioactive, cytotoxic, or pharmaceutically active moiety), or a molecule that increases the suitability of the antigen binding protein for a particular use (e.g., administration to a subject, such as a human subject, or other in vivo or in vitro uses). Examples of molecules that can be used to derivatize an antigen binding protein include albumin (e.g., human serum albumin) and polyethylene glycol (PEG). Albumin-linked and PEGylated derivatives of antigen binding proteins can be prepared using techniques well known in the art. In one embodiment,

the antigen binding protein is conjugated or otherwise linked to transthyretin (TTR) or a TTR variant. The TTR or TTR variant can be chemically modified with, for example, a chemical selected from the group consisting of dextran, poly(n-vinyl pyrrolidone), polyethylene glycols, propylene glycol homopolymers, polypropylene oxide/ethylene oxide co-polymers, polyoxyethylated polyols and polyvinyl alcohols.

[0126] Other derivatives include covalent or aggregative conjugates of IL-23 antigen binding proteins with other proteins or polypeptides, such as by expression of recombinant fusion proteins comprising heterologous polypeptides fused to the N-terminus or C-terminus of an IL-23 antigen binding protein. For example, the conjugated peptide may be a heterologous signal (or leader) polypeptide, e.g., the yeast alpha-factor leader, or a peptide such as an epitope tag. IL-23 antigen binding protein-containing fusion proteins can comprise peptides added to facilitate purification or identification of the IL-23 antigen binding protein (e.g., poly-His). An IL-23 antigen binding protein also can be linked to the FLAG peptide as described in Hopp et al., 1988, *Bio/Technology* 6:1204; and U.S. Pat. No. 5,011,912. The FLAG peptide is highly antigenic and provides an epitope reversibly bound by a specific monoclonal antibody (mAb), enabling rapid assay and facile purification of expressed recombinant protein. Reagents useful for preparing fusion proteins in which the FLAG peptide is fused to a given polypeptide are commercially available (Sigma, St. Louis, Mo.).

[0127] Oligomers that contain one or more IL-23 antigen binding proteins may be employed as IL-23 antagonists. Oligomers may be in the form of covalently-linked or non-covalently-linked dimers, trimers, or higher oligomers. Oligomers comprising two or more IL-23 antigen binding proteins are contemplated for use, with one example being a homodimer. Other oligomers include heterodimers, homotrimers, heterotrimers, homotetramers, heterotetramers, etc. Oligomers comprising multiple IL-23-binding proteins joined via covalent or non-covalent interactions between peptide moieties fused to the IL-23 antigen binding proteins, are also included. Such peptides may be peptide linkers (spacers), or peptides that have the property of promoting oligomerization. Among the suitable peptide linkers are those described in U.S. Pat. Nos. 4,751,180 and 4,935,233. Leucine zippers and certain polypeptides derived from antibodies are among the peptides that can promote oligomerization of IL-23 antigen binding proteins attached thereto. Examples of leucine zipper domains suitable for producing soluble oligomeric proteins are described in WIPO Publication No. WO 94/10308; Hoppe et al., 1994, *FEBS Letters* 344:191; and Fanslow et al., 1994, *Semin. Immunol.* 6:267-278. In one approach, recombinant fusion proteins comprising an IL-23 antigen binding protein fragment or derivative fused to a leucine zipper peptide are expressed in suitable host cells, and the soluble oligomeric IL-23 antigen binding protein fragments or derivatives that form are recovered from the culture supernatant.

[0128] Such oligomers may comprise from two to four IL-23 antigen binding proteins. The IL-23 antigen binding protein moieties of the oligomer may be in any of the forms described above, e.g., variants or fragments. Preferably, the oligomers comprise IL-23 antigen binding proteins that have IL-23 binding activity. Oligomers may be prepared using polypeptides derived from immunoglobulins. Preparation of fusion proteins comprising certain heterologous polypeptides fused to various portions of antibody-derived polypeptides (including the Fc domain) has been described, e.g., by Ashkenazi et al., 1991, *Proc. Natl. Acad. Sci. USA* 88:10535; Byrn et al., 1990, *Nature* 344:677; and Hollenbaugh et al., 1992 "Construction of Immunoglobulin Fusion Proteins", in *Current Protocols in Immunology*, Suppl. 4, pages 10.19.1-10.19.11.

[0129] Also included are dimers comprising two fusion proteins created by fusing an IL-23 antigen binding protein to the Fc region of an antibody. The dimer can be made by, for example, inserting a gene fusion encoding the fusion protein into an appropriate expression vector, expressing the gene fusion in host cells transformed with the recombinant expression vector, and allowing the expressed fusion protein to assemble much like antibody molecules, whereupon interchain disulfide bonds form between the Fc moieties to yield the dimer. Such Fc polypeptides include native and mutein forms of polypeptides derived from the Fc region of an antibody. Truncated forms of such polypeptides containing the hinge region that promotes dimerization also are included. Fusion proteins comprising Fc moieties (and oligomers formed therefrom) offer the advantage of facile purification by affinity chromatography over Protein A or Protein G columns. One suitable Fc polypeptide, described in WIPO Publication No. WO 93/10151 and U.S. Pat. Nos. 5,426,048 and 5,262,522, is a single chain

polypeptide extending from the N-terminal hinge region to the native C-terminus of the Fc region of a human IgG1 antibody. Another useful Fc polypeptide is the Fc mutein described in U.S. Pat. No. 5,457,035, and in Baum et al., 1994, *EMBO J.* 13:3992-4001. The amino acid sequence of this mutein is identical to that of the native Fc sequence presented in WIPO Publication No. WO 93/10151, except that amino acid 19 has been changed from Leu to Ala, amino acid 20 has been changed from Leu to Glu, and amino acid 22 has been changed from Gly to Ala. The mutein exhibits reduced affinity for Fc receptors.

[0130] Glycosylation

[0131] The antigen binding protein may have a glycosylation pattern that is different or altered from that found in the native species. As is known in the art, glycosylation patterns can depend on both the sequence of the protein (e.g., the presence or absence of particular glycosylation amino acid residues, discussed below), or the host cell or organism in which the protein is produced. Particular expression systems are discussed below.

[0132] Glycosylation of polypeptides is typically either N-linked or O-linked. N-linked refers to the attachment of the carbohydrate moiety to the side chain of an asparagine residue. The tri-peptide sequences asparagine-X-serine and asparagine-X-threonine, where X is any amino acid except proline, are the recognition sequences for enzymatic attachment of the carbohydrate moiety to the asparagine side chain. Thus, the presence of either of these tri-peptide sequences in a polypeptide creates a potential glycosylation site. O-linked glycosylation refers to the attachment of one of the sugars N-acetylgalactosamine, galactose, or xylose, to a hydroxyamino acid, most commonly serine or threonine, although 5-hydroxyproline or 5-hydroxylysine may also be used.

[0133] Addition of glycosylation sites to the antigen binding protein is conveniently accomplished by altering the amino acid sequence such that it contains one or more of the above-described tri-peptide sequences (for N-linked glycosylation sites). The alteration may also be made by the addition of, or substitution by, one or more serine or threonine residues to the starting sequence (for O-linked glycosylation sites). For ease, the antigen binding protein amino acid sequence may be altered through changes at the DNA level, particularly by mutating the DNA encoding the target polypeptide at preselected bases such that codons are generated that will translate into the desired amino acids.

[0134] Another means of increasing the number of carbohydrate moieties on the antigen binding protein is by chemical or enzymatic coupling of glycosides to the protein. These procedures are advantageous in that they do not require production of the protein in a host cell that has glycosylation capabilities for N- and O-linked glycosylation. Depending on the coupling mode used, the sugar(s) may be attached to (a) arginine and histidine, (b) free carboxyl groups, (c) free sulfhydryl groups such as those of cysteine, (d) free hydroxyl groups such as those of serine, threonine, or hydroxyproline, (e) aromatic residues such as those of phenylalanine, tyrosine, or tryptophan, or (f) the amide group of glutamine. These methods are described in PCT Publication No. WO 87/05330, and in Aplin and Wriston, 1981, *CRC Crit. Rev, Biochem.*, pp. 259-306.

[0135] Removal of carbohydrate moieties present on the starting antigen binding protein may be accomplished chemically or enzymatically. Chemical deglycosylation requires exposure of the protein to the compound trifluoromethanesulfonic acid, or an equivalent compound. This treatment results in the cleavage of most or all sugars except the linking sugar (N-acetylglucosamine or N-acetylgalactosamine), while leaving the polypeptide intact. Chemical deglycosylation is described by Hakimuddin et al., 1987, *Arch. Biochem. Biophys.* 259:52 and by Edge et al., 1981, *Anal. Biochem.* 118:131. Enzymatic cleavage of carbohydrate moieties on polypeptides can be achieved by the use of a variety of endo- and exo-glycosidases as described by Thotakura et al., 1987, *Meth. Enzymol.* 138:350. Glycosylation at potential glycosylation sites may be prevented by the use of the compound tunicamycin as described by Duskin et al., 1982, *J. Biol. Chem.* 257:3105. Tunicamycin blocks the formation of protein-N-glycoside linkages.

[0136] Hence, aspects include glycosylation variants of the antigen binding proteins wherein the number and/or type of glycosylation site(s) has been altered compared to the amino acid sequences of

the parent polypeptide. In certain embodiments, antigen binding protein variants comprise a greater or a lesser number of N-linked glycosylation sites than the parent polypeptide. Substitutions that eliminate or alter this sequence will prevent addition of an N-linked carbohydrate chain present in the parent polypeptide. For example, the glycosylation can be reduced by the deletion of an Asn or by substituting the Asn with a different amino acid. Antibodies typically have a N-linked glycosylation site in the Fc region.

[0137] Labels And Effector Groups

[0138] Antigen binding proteins may comprise one or more labels. The term "label" or "labeling group" refers to any detectable label. In general, labels fall into a variety of classes, depending on the assay in which they are to be detected: a) isotopic labels, which may be radioactive or heavy isotopes; b) magnetic labels (e.g., magnetic particles); c) redox active moieties; d) optical dyes; enzymatic groups (e.g. horseradish peroxidase, β -galactosidase, luciferase, alkaline phosphatase); e) biotinylated groups; and f) predetermined polypeptide epitopes recognized by a secondary reporter (e.g., leucine zipper pair sequences, binding sites for secondary antibodies, metal binding domains, epitope tags, etc.). In some embodiments, the labeling group is coupled to the antigen binding protein via spacer arms of various lengths to reduce potential steric hindrance. Various methods for labeling proteins are known in the art. Examples of suitable labeling groups include, but are not limited to, the following: radioisotopes or radionuclides (e.g., ³H, ¹⁴C, ¹⁵N, ³⁵S, ⁹⁰Y, ⁹⁹Tc, ¹¹¹In, ¹²⁵I, ¹³¹I), fluorescent groups (e.g., FITC, rhodamine, lanthanide phosphors), enzymatic groups (e.g., horseradish peroxidase, β -galactosidase, luciferase, alkaline phosphatase), chemiluminescent groups, biotinyl groups, or predetermined polypeptide epitopes recognized by a secondary reporter (e.g., leucine zipper pair sequences, binding sites for secondary antibodies, metal binding domains, epitope tags). In some embodiments, the labeling group is coupled to the antigen binding protein via spacer arms of various lengths to reduce potential steric hindrance. Various methods for labeling proteins are known in the art and may be used as is seen fit.

[0139] The term "effector group" means any group coupled to an antigen binding protein that acts as a cytotoxic agent. Examples for suitable effector groups are radioisotopes or radionuclides (e.g., ³H, ¹⁴C, ¹⁵N, ³⁵S, ⁹⁰Y, ⁹⁹Tc, ¹¹¹In, ¹²⁵I, ¹³¹I). Other suitable groups include toxins, therapeutic groups, or chemotherapeutic groups. Examples of suitable groups include calicheamicin, auristatins, geldanamycin and maytansine. In some embodiments, the effector group is coupled to the antigen binding protein via spacer arms of various lengths to reduce potential steric hindrance.

[0140] Polynucleotides Encoding IL-23 Antigen Binding Proteins

[0141] Polynucleotides that encode the antigen binding proteins described herein, or portions thereof, are also provided, including polynucleotides encoding one or both chains of an antibody, or a fragment, derivative, mutein, or variant thereof, polynucleotides encoding heavy chain variable regions or only CDRs, polynucleotides sufficient for use as hybridization probes, PCR primers or sequencing primers for identifying, analyzing, mutating or amplifying a polynucleotide encoding a polypeptide, anti-sense nucleic acids for inhibiting expression of a polynucleotide, and complementary sequences of the foregoing. The polynucleotides can be any length. They can be, for example, 5, 10, 15, 20, 25, 30, 35, 40, 45, 50, 75, 85, 95, 100, 125, 150, 175, 200, 250, 300, 350, 400, 450, 500, 750, 1,000, 1,500, 3,000, 5,000 or more nucleic acids in length, including all values in between, and/or can comprise one or more additional sequences, for example, regulatory sequences, and/or be part of a larger polynucleotide, for example, a vector. The polynucleotides can be single-stranded or double-stranded and can comprise RNA and/or DNA nucleic acids and artificial variants thereof (e.g., peptide nucleic acids).

[0142] Polynucleotides encoding certain antigen binding proteins, or portions thereof (e.g., full length antibody, heavy or light chain, variable domain, or a CDRH1, CDRH2, CDRH3, CDRL1, CDRL2, or CDRL3) may be isolated from B-cells of mice that have been immunized with IL-23 or an immunogenic fragment thereof. The polynucleotide may be isolated by conventional procedures such as polymerase chain reaction (PCR). Phage display is another example of a known technique whereby derivatives of

antibodies and other antigen binding proteins may be prepared. In one approach, polypeptides that are components of an antigen binding protein of interest are expressed in any suitable recombinant expression system, and the expressed polypeptides are allowed to assemble to form antigen binding protein molecules. Phage display is also used to derive antigen binding proteins having different properties (i.e., varying affinities for the antigen to which they bind) via chain shuffling, see Marks et al., 1992, *BioTechnology* 10:779.

[0143] Due to the degeneracy of the genetic code, each of the polypeptide sequences depicted herein are also encoded by a large number of other polynucleotide sequences besides those provided. For example, heavy chain variable domains provided herein may be encoded by polynucleotide sequences SEQ ID NOs: 32, 35, 37, 39, 41, 43, 45, 47, 49, 51, 53, 55, 57, or 59. Light chain variable domains may be encoded by polynucleotide sequences SEQ ID NOs: 2, 5, 6, 8, 10, 12, 14, 16, 18, 20, 22, 24, 26, or 28. One of ordinary skill in the art will appreciate that the present application thus provides adequate written description and enablement for each degenerate nucleotide sequence encoding each antigen binding protein.

[0144] An aspect further provides polynucleotides that hybridize to other polynucleotide molecules under particular hybridization conditions. Methods for hybridizing nucleic acids, basic parameters affecting the choice of hybridization conditions and guidance for devising suitable conditions are well-known in the art. See, e.g., Sambrook, Fritsch, and Maniatis (2001, *Molecular Cloning: A Laboratory Manual*, Cold Spring Harbor Laboratory Press, Cold Spring Harbor, N.Y. and *Current Protocols in Molecular Biology*, 1995, Ausubel et al., eds., John Wiley & Sons, Inc. As defined herein, a moderately stringent hybridization condition uses a prewashing solution containing 5× sodium chloride/sodium citrate (SSC), 0.5% SDS, 1.0 mM EDTA (pH 8.0), hybridization buffer of about 50% formamide, 6×SSC, and a hybridization temperature of 55° C. (or other similar hybridization solutions, such as one containing about 50% formamide, with a hybridization temperature of 42° C.), and washing conditions of 60° C., in 0.5×SSC, 0.1% SDS. A stringent hybridization condition hybridizes in 6×SSC at 45° C., followed by one or more washes in 0.1×SSC, 0.2% SDS at 68° C. Furthermore, one of skill in the art can manipulate the hybridization and/or washing conditions to increase or decrease the stringency of hybridization such that polynucleotides comprising nucleic acid sequences that are at least 65%, 70%, 75%, 80%, 85%, 90%, 91%, 92, 93%, 94%, 95%, 96%, 97%, 98% or 99% identical to each other, including all values in between, typically remain hybridized to each other.

[0145] Changes can be introduced by mutation into a polynucleotide, thereby leading to changes in the amino acid sequence of a polypeptide (e.g., an antigen binding protein or antigen binding protein derivative) that it encodes. Mutations can be introduced using any technique known in the art, such as site-directed mutagenesis and random mutagenesis. Mutant polypeptides can be expressed and selected for a desired property. Mutations can be introduced into a polynucleotide without significantly altering the biological activity of a polypeptide that it encodes. For example, substitutions at non-essential amino acid residues. Alternatively, one or more mutations can be introduced into a polynucleotide that selectively change the biological activity of a polypeptide that it encodes. For example, the mutation can quantitatively or qualitatively change the biological activity, such as increasing, reducing or eliminating the activity and changing the antigen specificity of an antigen binding protein.

[0146] Another aspect provides polynucleotides that are suitable for use as primers or hybridization probes for the detection of nucleic acid sequences. A polynucleotide can comprise only a portion of a nucleic acid sequence encoding a full-length polypeptide, for example, a fragment that can be used as a probe or primer or a fragment encoding an active portion (e.g., an IL-23 binding portion) of a polypeptide. Probes based on the sequence of a nucleic acid can be used to detect the nucleic acid or similar nucleic acids, for example, transcripts encoding a polypeptide. The probe can comprise a label group, e.g., a radioisotope, a fluorescent compound, an enzyme, or an enzyme co-factor. Such probes can be used to identify a cell that expresses the polypeptide.

[0147] Methods of Expressing Antigen Binding Proteins

[0148] The antigen binding proteins provided herein may be prepared by any of a number of conventional techniques. For example, IL-23 antigen binding proteins may be produced by recombinant expression systems, using any technique known in the art. See, e.g., *Monoclonal Antibodies, Hybridomas: A New Dimension in Biological Analyses*, Kennet et al. (eds.) Plenum Press, New York (1980); and *Antibodies: A Laboratory Manual*, Harlow and Lane (eds.), Cold Spring Harbor Laboratory Press, Cold Spring Harbor, N.Y. (1988).

[0149] Expression systems and constructs in the form of plasmids, expression vectors, transcription or expression cassettes that comprise at least one polynucleotide as described above are also provided herein, as well host cells comprising such expression systems or constructs. As used herein, "vector" means any molecule or entity (e.g., nucleic acid, plasmid, bacteriophage or virus) suitable for use to transfer protein coding information into a host cell. Examples of vectors include, but are not limited to, plasmids, viral vectors, non-episomal mammalian vectors and expression vectors, for example, recombinant expression vectors. Expression vectors, such as recombinant expression vectors, are useful for transformation of a host cell and contain nucleic acid sequences that direct and/or control (in conjunction with the host cell) expression of one or more heterologous coding regions operatively linked thereto. An expression construct may include, but is not limited to, sequences that affect or control transcription, translation, and, if introns are present, affect RNA splicing of a coding region operably linked thereto. "Operably linked" means that the components to which the term is applied are in a relationship that allows them to carry out their inherent functions. For example, a control sequence, e.g., a promoter, in a vector that is "operably linked" to a protein coding sequence are arranged such that normal activity of the control sequence leads to transcription of the protein coding sequence resulting in recombinant expression of the encoded protein.

[0150] Another aspect provides host cells into which an expression vector, such as a recombinant expression vector, has been introduced. A host cell can be any prokaryotic cell (for example, *E. coli*) or eukaryotic cell (for example, yeast, insect, or mammalian cells (e.g., CHO cells)). Vector DNA can be introduced into prokaryotic or eukaryotic cells via conventional transformation or transfection techniques. For stable transfection of mammalian cells, it is known that, depending upon the expression vector and transfection technique used, only a small fraction of cells may integrate the foreign DNA into their genome. In order to identify and select these integrants, a gene that encodes a selectable marker (e.g., for resistance to antibiotics) is generally introduced into the host cells along with the gene of interest. Preferred selectable markers include those which confer resistance to drugs, such as G418, hygromycin and methotrexate. Cells stably transfected with the introduced polynucleotide can be identified by drug selection (e.g., cells that have incorporated the selectable marker gene will survive, while the other cells die), among other methods.

[0151] Antigen binding proteins can be expressed in hybridoma cell lines (e.g., in particular antibodies may be expressed in hybridomas) or in cell lines other than hybridomas. Expression constructs encoding the antigen binding proteins can be used to transform a mammalian, insect or microbial host cell. Transformation can be performed using any known method for introducing polynucleotides into a host cell, including, for example packaging the polynucleotide in a virus or bacteriophage and transducing a host cell with the construct by transfection procedures known in the art, as exemplified by U.S. Pat. Nos. 4,399,216; 4,912,040; 4,740,461; 4,959,455. The optimal transformation procedure used will depend upon which type of host cell is being transformed. Methods for introduction of heterologous polynucleotides into mammalian cells are well known in the art and include, but are not limited to, dextran-mediated transfection, calcium phosphate precipitation, polybrene mediated transfection, protoplast fusion, electroporation, encapsulation of the polynucleotide(s) in liposomes, mixing nucleic acid with positively-charged lipids, and direct microinjection of the DNA into nuclei.

[0152] Recombinant expression constructs typically comprise a polynucleotide encoding a polypeptide. The polypeptide may comprise one or more of the following: one or more CDRs such as provided herein; a light chain variable region; a heavy chain variable region; a light chain constant region; a heavy chain constant region (e.g., C.sub.H1, C.sub.H2 and/or C.sub.H3); and/or another scaffold portion of an IL-23 antigen binding protein. These nucleic acid sequences are inserted into an appropriate expression vector using standard ligation techniques. In one embodiment, the heavy or light chain constant region is appended to the C-terminus of a heavy or light chain variable region

provided herein and is ligated into an expression vector. The vector is typically selected to be functional in the particular host cell employed (i.e., the vector is compatible with the host cell machinery, permitting amplification and/or expression of the gene can occur). In some embodiments, vectors are used that employ protein-fragment complementation assays using protein reporters, such as dihydrofolate reductase (see, for example, U.S. Pat. No. 6,270,964). Suitable expression vectors can be purchased, for example, from Invitrogen Life Technologies (Carlsbad, Calif.) or BD Biosciences (San Jose, Calif.). Other useful vectors for cloning and expressing the antibodies and fragments include those described in Bianchi and McGrew, 2003, *Biotech. Biotechnol. Bioeng.* 84:439-44. Additional suitable expression vectors are discussed, for example, in *Methods Enzymol.*, vol. 185 (D. V. Goeddel, ed.), 1990, New York: Academic Press.

[0153] Typically, expression vectors used in any of the host cells will contain sequences for plasmid maintenance and for cloning and expression of exogenous nucleotide sequences. Such sequences, collectively referred to as “flanking sequences” in certain embodiments will typically include one or more of the following nucleotide sequences: a promoter, one or more enhancer sequences, an origin of replication, a transcriptional termination sequence, a complete intron sequence containing a donor and acceptor splice site, a sequence encoding a leader sequence for polypeptide secretion, a ribosome binding site, a polyadenylation sequence, a polylinker region for inserting the polynucleotide encoding the polypeptide to be expressed, and a selectable marker element. The expression vectors that are provided may be constructed from a starting vector such as a commercially available vector. Such vectors may or may not contain all of the desired flanking sequences. Where one or more of the flanking sequences described herein are not already present in the vector, they may be individually obtained and ligated into the vector. Methods used for obtaining each of the flanking sequences are well known to one skilled in the art.

[0154] Optionally, the vector may contain a “tag”-encoding sequence, i.e., an oligonucleotide molecule located at the 5' or 3' end of the IL-23 antigen binding protein coding sequence; the oligonucleotide sequence encodes polyHis (such as hexaHis), or another “tag” such as FLAG®, HA (hemagglutinin influenza virus), or myc, for which commercially available antibodies exist. This tag is typically fused to the polypeptide upon expression of the polypeptide, and can serve as a means for affinity purification or detection of the IL-23 antigen binding protein from the host cell. Affinity purification can be accomplished, for example, by column chromatography using antibodies against the tag as an affinity matrix. Optionally, the tag can subsequently be removed from the purified IL-23 antigen binding protein by various means such as using certain peptidases for cleavage.

[0155] Flanking sequences may be homologous (i.e., from the same species and/or strain as the host cell), heterologous (i.e., from a species other than the host cell species or strain), hybrid (i.e., a combination of flanking sequences from more than one source), synthetic or native. As such, the source of a flanking sequence may be any prokaryotic or eukaryotic organism, any vertebrate or invertebrate organism, or any plant, provided that the flanking sequence is functional in, and can be activated by, the host cell machinery.

[0156] Flanking sequences useful in the vectors may be obtained by any of several methods well known in the art. Typically, flanking sequences useful herein will have been previously identified by mapping and/or by restriction endonuclease digestion and can thus be isolated from the proper tissue source using the appropriate restriction endonucleases. In some cases, the full nucleotide sequence of a flanking sequence may be known. Here, the flanking sequence may be synthesized using the methods described herein for nucleic acid synthesis or cloning.

[0157] Whether all or only a portion of the flanking sequence is known, it may be obtained using polymerase chain reaction (PCR) and/or by screening a genomic library with a suitable probe such as an oligonucleotide and/or flanking sequence fragment from the same or another species. Where the flanking sequence is not known, a fragment of DNA containing a flanking sequence may be isolated from a larger piece of DNA that may contain, for example, a coding sequence or even another gene or genes. Isolation may be accomplished by restriction endonuclease digestion to produce the proper DNA fragment followed by isolation using agarose gel purification, Qiagen® column chromatography

(Qiagen, Chatsworth, Calif.), or other methods known to the skilled artisan. The selection of suitable enzymes to accomplish this purpose will be readily apparent to one of ordinary skill in the art.

[0158] An origin of replication is typically a part of those prokaryotic expression vectors purchased commercially, and the origin aids in the amplification of the vector in a host cell. If the vector of choice does not contain an origin of replication site, one may be chemically synthesized based on a known sequence, and ligated into the vector. For example, the origin of replication from the plasmid pBR322 (New England Biolabs, Beverly, Mass.) is suitable for most gram-negative bacteria, and various viral origins (e.g., SV40, polyoma, adenovirus, vesicular stomatitis virus (VSV), or papillomaviruses such as HPV or BPV) are useful for cloning vectors in mammalian cells. Generally, the origin of replication component is not needed for mammalian expression vectors (for example, the SV40 origin is often used only because it also contains the virus early promoter).

[0159] A transcription termination sequence is typically located 3' to the end of a polypeptide coding region and serves to terminate transcription. Usually, a transcription termination sequence in prokaryotic cells is a G-C rich fragment followed by a poly-T sequence. While the sequence is easily cloned from a library or even purchased commercially as part of a vector, it can also be readily synthesized using methods for nucleic acid synthesis such as those described herein.

[0160] A selectable marker gene encodes a protein necessary for the survival and growth of a host cell grown in a selective culture medium. Typical selection marker genes encode proteins that (a) confer resistance to antibiotics or other toxins, e.g., ampicillin, tetracycline, or kanamycin for prokaryotic host cells; (b) complement auxotrophic deficiencies of the cell; or (c) supply critical nutrients not available from complex or defined media. Specific selectable markers are the kanamycin resistance gene, the ampicillin resistance gene, and the tetracycline resistance gene. Advantageously, a neomycin resistance gene may also be used for selection in both prokaryotic and eukaryotic host cells.

[0161] Other selectable genes may be used to amplify the gene that will be expressed. Amplification is the process wherein genes that are required for production of a protein critical for growth or cell survival are reiterated in tandem within the chromosomes of successive generations of recombinant cells. Examples of suitable selectable markers for mammalian cells include dihydrofolate reductase (DHFR) and promoterless thymidine kinase genes. Mammalian cell transformants are placed under selection pressure wherein only the transformants are uniquely adapted to survive by virtue of the selectable gene present in the vector. Selection pressure is imposed by culturing the transformed cells under conditions in which the concentration of selection agent in the medium is successively increased, thereby leading to the amplification of both the selectable gene and the DNA that encodes another gene, such as an antigen binding protein that binds to IL-23. As a result, increased quantities of a polypeptide such as an antigen binding protein are synthesized from the amplified DNA.

[0162] A ribosome-binding site is usually necessary for translation initiation of mRNA and is characterized by a Shine-Dalgarno sequence (prokaryotes) or a Kozak sequence (eukaryotes). The element is typically located 3' to the promoter and 5' to the coding sequence of the polypeptide to be expressed.

[0163] In some cases, such as where glycosylation is desired in a eukaryotic host cell expression system, one may manipulate the various pre- or pro-sequences to improve glycosylation or yield. For example, one may alter the peptidase cleavage site of a particular signal peptide, or add prosequences, which also may affect glycosylation. The final protein product may have, in the -1 position (relative to the first amino acid of the mature protein), one or more additional amino acids incident to expression, which may not have been totally removed. For example, the final protein product may have one or two amino acid residues found in the peptidase cleavage site, attached to the amino-terminus. Alternatively, use of some enzyme cleavage sites may result in a slightly truncated form of the desired polypeptide, if the enzyme cuts at such area within the mature polypeptide.

[0164] Expression and cloning will typically contain a promoter that is recognized by the host organism and operably linked to the molecule encoding an IL-23 antigen binding protein. Promoters are

untranscribed sequences located upstream (i.e., 5') to the start codon of a structural gene (generally within about 100 to 1000 bp) that control transcription of the structural gene. Promoters are conventionally grouped into one of two classes: inducible promoters and constitutive promoters. Inducible promoters initiate increased levels of transcription from DNA under their control in response to some change in culture conditions, such as the presence or absence of a nutrient or a change in temperature. Constitutive promoters, on the other hand, uniformly transcribe a gene to which they are operably linked, that is, with little or no control over gene expression. A large number of promoters, recognized by a variety of potential host cells, are well known. A suitable promoter is operably linked to the DNA encoding a heavy chain variable region or a light chain variable region of an IL-23 antigen binding protein by removing the promoter from the source DNA by restriction enzyme digestion and inserting the desired promoter sequence into the vector.

[0165] Suitable promoters for use with yeast hosts are also well known in the art. Yeast enhancers are advantageously used with yeast promoters. Suitable promoters for use with mammalian host cells are well known and include, but are not limited to, those obtained from the genomes of viruses such as polyoma virus, fowlpox virus, adenovirus (such as Adenovirus 2), bovine papilloma virus, avian sarcoma virus, cytomegalovirus, retroviruses, hepatitis-B virus, and Simian Virus 40 (SV40). Other suitable mammalian promoters include heterologous mammalian promoters, for example, heat-shock promoters and the actin promoter.

[0166] Additional promoters which may be of interest include, but are not limited to: SV40 early promoter (Benoist and Chambon, 1981, *Nature* 290:304-310); CMV promoter (Thornsen et al., 1984, *Proc. Natl. Acad. U.S.A.* 81:659-663); the promoter contained in the 3' long terminal repeat of Rous sarcoma virus (Yamamoto et al., 1980, *Cell* 22:787-797); herpes thymidine kinase promoter (Wagner et al., 1981, *Proc. Natl. Acad. Sci. U.S.A.* 78:1444-1445); promoter and regulatory sequences from the metallothionein gene (Prinster et al., 1982, *Nature* 296:39-42); and prokaryotic promoters such as the beta-lactamase promoter (Villa-Kamaroff et al., 1978, *Proc. Natl. Acad. Sci. U.S.A.* 75:3727-3731); or the tac promoter (DeBoer et al., 1983, *Proc. Natl. Acad. Sci. U.S.A.* 80:21-25). Also of interest are the following animal transcriptional control regions, which exhibit tissue specificity and have been utilized in transgenic animals: the elastase I gene control region that is active in pancreatic acinar cells (Swift et al., 1984, *Cell* 38:639-646; Ornitz et al., 1986, *Cold Spring Harbor Symp. Quant. Biol.* 50:399-409; MacDonald, 1987, *Hepatology* 7:425-515); the insulin gene control region that is active in pancreatic beta cells (Hanahan, 1985, *Nature* 315:115-122); the immunoglobulin gene control region that is active in lymphoid cells (Grosschedl et al., 1984, *Cell* 38:647-658; Adames et al., 1985, *Nature* 318:533-538; Alexander et al., 1987, *Mol. Cell. Biol.* 7:1436-1444); the mouse mammary tumor virus control region that is active in testicular, breast, lymphoid and mast cells (Leder et al., 1986, *Cell* 45:485-495); the albumin gene control region that is active in liver (Pinkert et al., 1987, *Genes and Devel.* 1:268-276); the alpha-feto-protein gene control region that is active in liver (Krumlauf et al., 1985, *Mol. Cell. Biol.* 5:1639-1648; Hammer et al., 1987, *Science* 253:53-58); the alpha 1-antitrypsin gene control region that is active in liver (Kelsey et al., 1987, *Genes and Devel.* 1:161-171); the beta-globin gene control region that is active in myeloid cells (Mogram et al., 1985, *Nature* 315:338-340; Kollias et al., 1986, *Cell* 46:89-94); the myelin basic protein gene control region that is active in oligodendrocyte cells in the brain (Readhead et al., 1987, *Cell* 48:703-712); the myosin light chain-2 gene control region that is active in skeletal muscle (Sani, 1985, *Nature* 314:283-286); and the gonadotropic releasing hormone gene control region that is active in the hypothalamus (Mason et al., 1986, *Science* 234:1372-1378).

[0167] An enhancer sequence may be inserted into the vector to increase transcription by higher eukaryotes. Enhancers are cis-acting elements of DNA, usually about 10-300 bp in length, that act on the promoter to increase transcription. Enhancers are relatively orientation and position independent, having been found at positions both 5' and 3' to the transcription unit. Several enhancer sequences available from mammalian genes are known (e.g., globin, elastase, albumin, alpha-feto-protein and insulin). Typically, however, an enhancer from a virus is used. The SV40 enhancer, the cytomegalovirus early promoter enhancer, the polyoma enhancer, and adenovirus enhancers known in the art are exemplary enhancing elements for the activation of eukaryotic promoters. While an enhancer may be positioned in the vector either 5' or 3' to a coding sequence, it is typically located at a site 5' from the promoter. A sequence encoding an appropriate native or heterologous signal sequence (leader sequence or signal peptide) can be incorporated into an expression vector, to

promote extracellular secretion of the antibody. The choice of signal peptide or leader depends on the type of host cells in which the antibody is to be produced, and a heterologous signal sequence can replace the native signal sequence. Examples of signal peptides that are functional in mammalian host cells include the following: the signal sequence for interleukin-7 described in U.S. Pat. No. 4,965,195; the signal sequence for interleukin-2 receptor described in Cosman et al., 1984, *Nature* 312:768; the interleukin-4 receptor signal peptide described in EP Patent No. 0367 566; the type I interleukin-1 receptor signal peptide described in U.S. Pat. No. 4,968,607; the type II interleukin-1 receptor signal peptide described in EP Patent No. 0 460 846.

[0168] After the vector has been constructed, the completed vector may be inserted into a suitable host cell for amplification and/or polypeptide expression. The transformation of an expression vector for an antigen binding protein into a selected host cell may be accomplished by well known methods including transfection, infection, calcium phosphate co-precipitation, electroporation, microinjection, lipofection, DEAE-dextran mediated transfection, or other known techniques. The method selected will in part be a function of the type of host cell to be used. These methods and other suitable methods are well known to the skilled artisan, and are set forth, for example, in Sambrook et al., *Molecular Cloning: A Laboratory Manual*, 3rd ed., Cold Spring Harbor Laboratory Press, Cold Spring Harbor, N.Y. (2001).

[0169] A host cell, when cultured under appropriate conditions, synthesizes protein that can be subsequently collected from the culture medium (if the host cell secretes it into the medium) or directly from the host cell producing it (if it is not secreted). The selection of an appropriate host cell will depend upon various factors, such as desired expression levels, polypeptide modifications that are desirable or necessary for activity (such as glycosylation or phosphorylation) and ease of folding into a biologically active molecule.

[0170] Mammalian cell lines available as hosts for expression are well known in the art and include, but are not limited to, immortalized cell lines available from the American Type Culture Collection (ATCC), including but not limited to Chinese hamster ovary (CHO) cells, HeLa cells, baby hamster kidney (BHK) cells, monkey kidney cells (COS), human hepatocellular carcinoma cells (e.g., Hep G2), and a number of other cell lines. In certain embodiments, cell lines may be selected through determining which cell lines have high expression levels and constitutively produce antigen binding proteins with IL-23 binding properties. In another embodiment, a cell line from the B cell lineage that does not make its own antibody but has a capacity to make and secrete a heterologous antibody can be also selected.

[0171] Use of Human IL-23 Antigen Binding Proteins for Diagnostic and Therapeutic Purposes

[0172] Antigen binding proteins are useful for detecting IL-23 in biological samples and identification of cells or tissues that produce IL-23. Antigen binding proteins that specifically bind to IL-23 may be used in diagnosis and/or treatment of diseases related to IL-23 in a patient in need thereof. For one, the IL-23 antigen binding proteins can be used in diagnostic assays, e.g., binding assays to detect and/or quantify IL-23 expressed in blood, serum, cells or tissue. In addition, IL-23 antigen binding proteins can be used to reduce, inhibit, interfere with or modulate one or more biological activities of IL-23 in a cell or tissue. Thus antigen binding proteins that bind to IL-23 may have therapeutic use in ameliorating diseases related to IL-23.

[0173] Indications

[0174] The present invention also relates to the use of IL-23 antigen binding proteins for use in the prevention or therapeutic treatment of medical disorders, such as those disclosed herein. The IL-23 antigen binding proteins are useful to treat a variety of conditions in which IL-23 is associated with or plays a role in contributing to the underlying disease or disorder or otherwise contributes to a negative symptom.

[0175] Conditions effectively treated by IL-23 antigen binding proteins play a role in the inflammatory response. Such inflammatory disorders include periodontal disease; lung disorders such as asthma; skin disorders such as psoriasis, atopic dermatitis, contact dermatitis; rheumatic disorders such as

rheumatoid arthritis, progressive systemic sclerosis (scleroderma); systemic lupus erythematosus; spondyloarthritis including ankylosing spondylitis, psoriatic arthritis, enteropathic arthritis and reactive arthritis. Also contemplated is uveitis including Vogt-Koyanagi-Harada disease, idiopathic anterior and posterior uveitis, and uveitis associated with spondyloarthritis. Use of IL-23 antigen binding proteins is also contemplated for the treatment of autoimmune disorders including multiple sclerosis; autoimmune myocarditis; type 1 diabetes and autoimmune thyroiditis.

[0176] Degenerative conditions of the gastrointestinal system are treatable or preventable with IL-23 antigen binding proteins. Such gastrointestinal disorders including inflammatory bowel disease: Crohn's disease, ulcerative colitis and Celiac disease.

[0177] Also included are use of IL-23 antigen binding proteins in treatments for graft-versus-host disease, and complications such as graft rejection, resulting from solid organ transplantation, such as heart, liver, skin, kidney, lung or other transplants, including bone marrow transplants.

[0178] Also provided herein are methods for using IL-23 antigen binding proteins to treat various oncologic disorders including various forms of cancer including colon, stomach, prostate, renal cell, cervical and ovarian cancers, and lung cancer (SCLC and NSCLC). Also included are solid tumors, including sarcoma, osteosarcoma, and carcinoma, such as adenocarcinoma and squamous cell carcinoma, esophageal cancer, gastric cancer, gall bladder carcinoma, leukemia, including acute myelogenous leukemia, chronic myelogenous leukemia, myeloid leukemia, chronic or acute lymphoblastic leukemia and hairy cell leukemia, and multiple myeloma.

[0179] Diagnostic Methods

[0180] The antigen binding proteins of the described can be used for diagnostic purposes to detect, diagnose, or monitor diseases and/or conditions associated with IL-23. Examples of methods useful in the detection of the presence of IL-23 include immunoassays, such as the enzyme linked immunosorbent assay (ELISA) and the radioimmunoassay (RIA).

[0181] For diagnostic applications, the antigen binding protein typically will be labeled with a detectable labeling group. Suitable labeling groups include, but are not limited to, the following: radioisotopes or radionuclides (e.g., ³H, ¹⁴C, ¹⁵N, ³⁵S, ⁹⁰Y, ⁹⁹Tc, ¹¹¹In, ¹²⁵I, ¹³¹I), fluorescent groups (e.g., FITC, rhodamine, lanthanide phosphors), enzymatic groups (e.g., horseradish peroxidase, β -galactosidase, luciferase, alkaline phosphatase), chemiluminescent groups, biotinyl groups, or predetermined polypeptide epitopes recognized by a secondary reporter (e.g., leucine zipper pair sequences, binding sites for secondary antibodies, metal binding domains, epitope tags). In some embodiments, the labelling group is coupled to the antigen binding protein via spacer arms of various lengths to reduce potential steric hindrance. Various methods for labelling proteins are known in the art and may be used.

[0182] Other diagnostic methods are provided for identifying a cell or cells that express IL-23. In a specific embodiment, the antigen binding protein is labeled with a labeling group and the binding of the labeled antigen binding protein to IL-23 is detected. In a further specific embodiment, the binding of the antigen binding protein to IL-23 is detected in vivo. In a further specific embodiment, the IL-23 antigen binding protein is isolated and measured using techniques known in the art. See, for example, Harlow and Lane, 1988, *Antibodies: A Laboratory Manual*, New York: Cold Spring Harbor (ed. 1991 and periodic supplements); John E. Coligan, ed., 1993, *Current Protocols In Immunology* New York: John Wiley & Sons.

[0183] Other methods provide for detecting the presence of a test molecule that competes for binding to IL-23 with the antigen binding proteins provided. An example of one such assay would involve detecting the amount of free antigen binding protein in a solution containing an amount of IL-23 in the presence or absence of the test molecule. An increase in the amount of free antigen binding protein (i.e., the antigen binding protein not bound to IL-23) would indicate that the test molecule is capable of competing for IL-23 binding with the antigen binding protein. In one embodiment, the antigen binding

protein is labeled with a labeling group. Alternatively, the test molecule is labeled and the amount of free test molecule is monitored in the presence and absence of an antigen binding protein.

[0184] Methods Of Treatment: Pharmaceutical Formulations, Routes Of Administration

[0185] Pharmaceutical compositions that comprise a therapeutically effective amount of one or a plurality of the antigen binding proteins and a pharmaceutically acceptable excipient, diluent, carrier, solubilizer, emulsifier, preservative, and/or adjuvant are provided. In addition, methods of treating a patient by administering such pharmaceutical composition are included. The term "patient" includes human patients. The terms "treat" and "treatment" encompass alleviation or prevention of at least one symptom or other aspect of a disorder, or reduction of disease severity, and the like. The term "therapeutically effective amount" or "effective amount" refers to the amount of an IL-23 antigen binding protein determined to produce any therapeutic response in a mammal. Such therapeutically effective amounts are readily ascertained by one of ordinary skill in the art.

[0186] An antigen binding protein need not affect a complete cure, or eradicate every symptom or manifestation of a disease, to constitute a viable therapeutic agent. As is recognized in the pertinent field, drugs employed as therapeutic agents may reduce the severity of a given disease state, but need not abolish every manifestation of the disease to be regarded as useful therapeutic agents. Similarly, a prophylactically administered treatment need not be completely effective in preventing the onset of a condition in order to constitute a viable prophylactic agent. Simply reducing the impact of a disease (for example, by reducing the number or severity of its symptoms, or by increasing the effectiveness of another treatment, or by producing another beneficial effect), or reducing the likelihood that the disease will occur or worsen in a subject, is sufficient. Certain methods provided herein comprise administering to a patient an IL-23 antagonist (such as the antigen binding proteins disclosed herein) in an amount and for a time sufficient to induce a sustained improvement over baseline of an indicator that reflects the severity of the particular disorder.

[0187] As is understood in the pertinent field, pharmaceutical compositions comprising the molecules of the invention are administered to a patient in a manner appropriate to the indication. Pharmaceutical compositions may be administered by any suitable technique, including but not limited to, parenterally, topically, or by inhalation. If injected, the pharmaceutical composition can be administered, for example, via intra-articular, intravenous, intramuscular, intralesional, intraperitoneal or subcutaneous routes, by bolus injection, or continuous infusion. Localized administration, e.g. at a site of disease or injury is contemplated, as are transdermal delivery and sustained release from implants. Delivery by inhalation includes, for example, nasal or oral inhalation, use of a nebulizer, inhalation of the antagonist in aerosol form, and the like. Other alternatives include eyedrops; oral preparations including pills, syrups, lozenges or chewing gum; and topical preparations such as lotions, gels, sprays, and ointments.

[0188] Use of antigen binding proteins in ex vivo procedures also is contemplated. For example, a patient's blood or other bodily fluid may be contacted with an antigen binding protein that binds IL-23 ex vivo. The antigen binding protein may be bound to a suitable insoluble matrix or solid support material.

[0189] Advantageously, antigen binding proteins are administered in the form of a composition comprising one or more additional components such as a physiologically acceptable carrier, excipient or diluent. Optionally, the composition additionally comprises one or more physiologically active agents for combination therapy. A pharmaceutical composition may comprise an IL-23 antigen binding protein together with one or more substances selected from the group consisting of a buffer, an antioxidant such as ascorbic acid, a low molecular weight polypeptide (such as those having fewer than 10 amino acids), a protein, an amino acid, a carbohydrate such as glucose, sucrose or dextrans, a chelating agent such as EDTA, glutathione, a stabilizer, and an excipient. Neutral buffered saline or saline mixed with conspecific serum albumin are examples of appropriate diluents. In accordance with appropriate industry standards, preservatives such as benzyl alcohol may also be added. The composition may be formulated as a lyophilizate using appropriate excipient solutions (e.g., sucrose) as diluents. Suitable components are nontoxic to recipients at the dosages and concentrations employed. Further examples

of components that may be employed in pharmaceutical formulations are presented in any Remington's Pharmaceutical Sciences including the 21st Ed. (2005), Mack Publishing Company, Easton, Pa.

[0190] Kits for use by medical practitioners include an IL-23 antigen binding protein and a label or other instructions for use in treating any of the conditions discussed herein. In one embodiment, the kit includes a sterile preparation of one or more IL-23 binding antigen binding proteins, which may be in the form of a composition as disclosed above, and may be in one or more vials.

[0191] Dosages and the frequency of administration may vary according to such factors as the route of administration, the particular antigen binding proteins employed, the nature and severity of the disease to be treated, whether the condition is acute or chronic, and the size and general condition of the subject. Appropriate dosages can be determined by procedures known in the pertinent art, e.g. in clinical trials that may involve dose escalation studies.

[0192] A typical dosage may range from about 0.1 $\mu\text{g}/\text{kg}$ to up to about 30 mg/kg or more, depending on the factors mentioned above. In specific embodiments, the dosage may range from 0.1 $\mu\text{g}/\text{kg}$ up to about 30 mg/kg , optionally from 1 $\mu\text{g}/\text{kg}$ up to about 30 mg/kg , optionally from 10 $\mu\text{g}/\text{kg}$ up to about 10 mg/kg , optionally from about 0.1 mg/kg to 5 mg/kg , or optionally from about 0.3 mg/kg to 3 mg/kg .

[0193] Dosing frequency will depend upon the pharmacokinetic parameters of the particular human IL-23 antigen binding protein in the formulation used. Typically, a clinician administers the composition until a dosage is reached that achieves the desired effect. The composition may therefore be administered as a single dose, or as two or more doses (which may or may not contain the same amount of the desired molecule) over time, or as a continuous infusion via an implantation device or catheter. Appropriate dosages may be ascertained through use of appropriate dose-response data. An IL-23 antigen binding protein of the invention may be administered, for example, once or more than once, e.g., at regular intervals over a period of time. In particular embodiments, an IL-23 antigen binding protein is administered over a period of at least a month or more, e.g., for one, two, or three months or even indefinitely. For treating chronic conditions, long-term treatment is generally most effective. However, for treating acute conditions, administration for shorter periods, e.g. from one to six weeks, may be sufficient. In general, the antigen binding protein is administered until the patient manifests a medically relevant degree of improvement over baseline for the chosen indicator or indicators.

[0194] It is contemplated that an IL-23 antigen binding protein be administered to the patient in an amount and for a time sufficient to induce an improvement, preferably a sustained improvement, in at least one indicator that reflects the severity of the disorder that is being treated. Various indicators that reflect the extent of the patient's illness, disease or condition may be assessed for determining whether the amount and time of the treatment is sufficient. Such indicators include, for example, clinically recognized indicators of disease severity, symptoms, or manifestations of the disorder in question. In one embodiment, an improvement is considered to be sustained if the subject exhibits the improvement on at least two occasions separated by two to four weeks. The degree of improvement generally is determined by a physician, who may make this determination based on signs, symptoms, biopsies, or other test results, and who may also employ questionnaires that are administered to the subject, such as quality-of-life questionnaires developed for a given disease.

[0195] Particular embodiments of methods and compositions of the invention involve the use of an IL-23 antigen binding protein and one or more additional IL-23 antagonists, for example, two or more antigen binding proteins of the invention, or an antigen binding protein of the invention and one or more other IL-23 antagonists. Also provided are IL-23 antigen binding proteins administered alone or in combination with other agents useful for treating the condition with which the patient is afflicted. Examples of such agents include both proteinaceous and non-proteinaceous drugs. Such agents include therapeutic moieties having anti-inflammatory properties (for example, non-steroidal anti-inflammatory agents, steroids, immunomodulators and/or other cytokine inhibitors such as those that antagonize, for example, IFN- γ , GM-CSF, IL-6, IL-8, IL-17, IL-22 and TNFs), or of an IL-23 antigen binding protein and one or more other treatments (e.g., surgery, ultrasound, or treatment effective to

reduce inflammation). When multiple therapeutics are co-administered, dosages may be adjusted accordingly, as is recognized or known in the pertinent art. Useful agents that may be combined with IL-23 antigen binding proteins include those used to treat, for example, Crohn's disease or ulcerative colitis, such as aminosalicylate (for example, mesalamine), corticosteroids (including predisone), antibiotics such as metronidazole or ciprofloxacin (or other antibiotics useful for treating, for example, patients afflicted with fistulas), and immunosuppressives such as azathioprine, 6-mercaptopurine, methotrexate, tacrolimus and cyclosporine. Such agent(s) may be administered orally or by another route, for example via suppository or enema. Agents which may be combined with IL-23 binding proteins in treatment of psoriasis include corticosteroids, calcipotriene and other vitamin D derivatives, acetretin and other retinoic acid derivatives, methotrexate, tacrolimus, and cyclosporine used topically or systemically. Such agents can be administered simultaneously, consecutively, alternately, or according to any other regimen that allows the total course of therapy to be effective.

[0196] In addition to human patients, IL-23 antigen binding proteins are useful in the treatment of non-human animals, such as domestic pets (dogs, cats, birds, primates, etc.), domestic farm animals (horses, cattle, sheep, pigs, birds, etc.). In such instances, an appropriate dose may be determined according to the animal's body weight. For example, a dose of 0.2-1 mg/kg may be used. Alternatively, the dose is determined according to the animal's surface area, an exemplary dose ranging from 0.1-20 mg/m², or more preferably, from 5-12 mg/m². For small animals, such as dogs or cats, a suitable dose is 0.4 mg/kg. IL-23 antigen binding protein (preferably constructed from genes derived from the recipient species) is administered by injection or other suitable route one or more times per week until the animal's condition is improved, or it may be administered indefinitely.

[0197] The following examples, including the experiments conducted and the results achieved, are provided for illustrative purposes only and are not to be construed as limiting the scope of the appended claims.

EXAMPLES

Example 1

Generation of Human IL-23 Antibodies

[0198] XenoMouse™ technology (Amgen, Thousand Oaks, Calif.) was used to develop human monoclonal antibodies that recognize and inhibit native human IL-23 activity while sparing human IL-12. The antibodies also recognize and inhibit recombinant cynomolgous IL-23 but do not recognize murine or rat IL-23.

[0199] Antibodies were selected for recognition and complete inhibition of native human IL-23 obtained from human monocyte-derived dendritic cells (MoDCs), using the STAT-luciferase reporter assay described below. Human monocytes were isolated from peripheral blood mononuclear cells from healthy donors using negative selection (Monocyte Isolation Kit II, Miltenyi Biotec, Auburn, Calif.). MoDCs were generated by culturing monocytes with human GM-CSF (50 ng/ml) and human IL-4 (100 ng/ml) for 7 days in RPMI 1640 with 10% fetal bovine serum complete medium. MoDCs were then washed twice with PBS followed by stimulation with human CD40L (1 µg/ml) for an additional 48 hours. CD40L-stimulated MoDC supernatant contains IL-23, IL-12 and IL-12/23p40. ELISAs are used to determine the amount of IL-12p70 (R&D System, Minneapolis, Minn.), IL-23 (eBiosciences, San Diego, Calif.) and IL-12/23p40 (R&D Systems). The STAT-luciferase assay responds to IL-23 and not to IL-12 or to free IL-12/23p40, therefore the assay could be used with crude supernatants to assess IL-23 activity. For use in the NK cell assay, described below, the native human IL-23 crude supernatant was purified using an IL-23 affinity column followed by size exclusion chromatography. Concentration was determined using an IL-23 specific ELISA (eBiosciences).

[0200] The purified antibody supernatants were also tested against recombinant human (rhu) IL-23 and recombinant cynomolgous (cyno) IL-23 in the STAT-luciferase assay. Of the antibodies tested that completely inhibited recombinant human IL-23, only half of those antibodies recognized and completely inhibited native human IL-23. Recognition and complete inhibition of recombinant human

IL-23 was not predictive of, nor correlated to, recognition and complete inhibition of native human IL-23. As shown in FIGS. 1A and 1B, of the antibody supernatants that completely inhibited recombinant human IL-23, only half of those antibodies completely inhibited native human IL-23. Those antibodies that recognized and completely inhibited native human IL-23 were selected for further characterization.

Example 2

Functional Assays

a) STAT-Luciferase Assay

[0201] It is known that IL-23 binds its heterodimeric receptor and signals through JAK2 and Tyk2 to activate STAT 1, 3, 4 and 5. In this assay, cells transfected with a STAT/luciferase reporter gene are used to assess the ability of the IL-23 antibodies to inhibit IL-23-induced bioactivity.

[0202] Chinese hamster ovary cells expressing human IL-23 receptor are transiently transfected with STAT-luciferase reporter overnight. IL-23 antibodies are serially diluted (12 points of 1:4 serial dilutions starting at 37.5 µg/ml) into 96 well plates. Native human IL-23 (preparation method is described in Example 1) is added to each well at a concentration of 2 ng/ml and incubated at room temperature for 15-20 minutes. The transiently transfected cells are added (8×10^3 cells) to a final volume of 100 µl/well and incubated for 5 hours at 37° C., 10% CO₂. Following incubation, cells are lysed using 100 µL/well Glo Lysis buffer (1×) (Promega, Madison, Wis.) at room temperature for 5 minutes. Fifty microliters of cell lysate is added to a 96 well plate along with 50 µL Bright-Glo luciferase substrate (Promega) and read on a luminometer.

[0203] Statistical analysis can be performed using GraphPad PRISM software (GraphPad Software, La Jolla, Calif.). Results can be expressed as the mean±standard deviation (SD).

[0204] As seen in TABLE 5, all IL-23 antibodies potently and completely inhibited native human IL-23-induced STAT/luciferase reporter in a dose dependent manner. The antibodies also potently and completely inhibited recombinant human (rhu) IL-23 and recombinant cyno (cyno) IL-23. The antibodies all had IC₅₀ values in the picomolar range.

TABLE-US-00005 TABLE 5 Table of mean IC₅₀ (pM) values for IL-23 antibodies in the STAT-luciferase assay. Native huIL-23 rhuIL-23 Cyno IL-23 antibody IC₅₀ +/- SD Repeats IC₅₀ +/- SD Repeats IC₅₀ +/- SD Repeats A 114 +/- 70 3 190 +/- 99 3 379 +/- 213 3 B 45 +/- 5 4 100 +/- 59 4 130 +/- 60 3 C 107 +/- 31 3 211 +/- 93 3 376 +/- 89 3 D 65 +/- 5 3 107 +/- 30 3 184 +/- 77 3 E 140 +/- 52 3 142 +/- 52 3 188 +/- 59 3 F 86 +/- 47 4 187 +/- 116 4 366 +/- 219 4 G 156 +/- 74 5 296 +/- 133 5 421 +/- 174 5 H 192 +/- 35 4 253 +/- 184 4 1024 +/- 533 4 I 208 +/- 33 3 338 +/- 140 3 650 +/- 42 3 J 83 +/- 54 2 36 +/- 6 2 56 +/- 2 2 K 71 +/- 38 3 43 +/- 20 3 61 +/- 10 3 L 113 +/- 80 3 23 +/- 7 3 47 +/- 1 3 M 34 +/- 11 2 40 +/- 8 2 56 +/- 6 2 N 361 +/- 164 3 145 1 238 1

[0205] b) NK Cell Assay

[0206] It is known that IL-23 acts on natural killer cells to induce expression of pro-inflammatory cytokines, such as interferon γ (IFNγ). In this assay, human primary natural killer (NK) cells are used to assess the ability of the IL-23 antibodies to inhibit IL-23-induced IFNγ activity in cells expressing the native receptor for human IL-23.

[0207] NK cells are isolated from multiple human donors via negative selection (NK Cell Isolation Kit, Miltenyi Biotec, Auburn, Calif.). Purified NK cells (1×10^6 cells/ml) are added to 6 well plates in RPMI 1640 plus 10% fetal bovine serum complete medium supplemented with recombinant human IL-2 (10 ng/ml, R&D Systems, Minneapolis, Minn.), to a final volume of 10 ml/well. Cells are cultured for 7 days at 37° C., 5% CO₂. The IL-2-activated NK cells are then stimulated with rhuIL-23 or cyno IL-23 (10 ng/ml) and recombinant human IL-18 (20 ng/ml, R&D Systems, Minneapolis, Minn.) in the presence of serial dilutions (11 points of 1:3 serial dilutions starting at 3 µg/ml) of IL-23 antibodies for

24 hours. IFN γ levels are measured in the supernatant by IFN γ ELISA (R&D Systems, Minneapolis, Minn.) according to manufacturer's instructions.

[0208] Statistical analysis can be performed using GraphPad PRISM software. Results can be expressed as the mean \pm standard deviation (SD).

[0209] As seen in TABLE 6, all antibodies potently inhibited rhIL-23 and cyno IL-23-induced IFN γ expression in NK cells in a dose dependent manner. The antibodies all had IC₅₀ values in the picomolar range. The assay was performed on a subset of antibodies using native human IL-23 (30 μ g/ml, preparation method is described in Example 1) and rhIL-18 (40 ng/ml, R&D Systems) and yielded the results shown in TABLE 6. Consistent with the selection for IL-23 specific antibodies, these anti-IL-23 antibodies had no effect on IL-12 stimulated IFN γ production in NK cells using the assay described above, whereas an IL-12p35 specific neutralizing antibody, mAb219 (R&D Systems, Minneapolis, Minn.) potently inhibited recombinant human IL-12.

TABLE-US-00006 TABLE 6 Table of mean IC₅₀ (pM) values for IL-23 antibodies in the NK cell assay. Native huIL-23 rhIL-23 Cyno IL-23 antibody IC₅₀ +/- SD Repeats IC₅₀ +/- SD Repeats IC₅₀ +/- SD Repeats A 42 +/- 12 2 31 +/- 21 2 B 85 +/- 30 2 48 +/- 30 3 19 +/- 8 2 C 32 +/- 19 4 29 +/- 16 2 D 37 +/- 21 2 29 +/- 19 2 E 158 +/- 50 2 57 +/- 14 3 21 +/- 3 2 F 25 +/- 15 2 21 +/- 17 2 G 152 +/- 72 2 45 +/- 30 3 23 +/- 8 2 H 29 +/- 28 2 33 +/- 17 2 I 69 1 52 1 J 4 +/- 3 2 5 +/- 3 2 K 7 +/- 2 2 8 +/- 6 2 L 3 +/- 1 2 4 +/- 1 2 M 8 1 12 1

[0210] c) Human Whole Blood Assay

[0211] Human whole blood is collected from multiple healthy donors using Repludan® (Bayer Pittsburgh, Pa.) as an anti-coagulant. The final concentration of Repludan® in whole blood is 10 μ g/ml. A stimulation mixture of rhIL-23 or cyno IL-23 (final concentration 1 ng/ml)+rhIL-18 (final concentration 20 ng/ml)+rhIL-2 (final concentration 5 ng/ml) in RPMI 1640+10% FBS, is added to a 96 well plate, final volume 20 μ l/well. Serially diluted IL-23 antibodies (11 points of 1:3 serial dilutions starting from 3 μ g/ml) are added at 20 μ l/well and incubated with the stimulation mixture for 30 minutes at room temperature. Whole blood is then added (120 μ l/well) and the final volume adjusted to 200 μ l/well with RPMI 1640+10% FBS. The final concentration of whole blood is 60%. The plates are incubated for 24 hours at 37° C., 5% CO₂. Cell free supernatants are harvested and IFN γ levels are measured from the supernatants by IFN γ ELISA (R&D Systems) according to manufacturer's instructions.

[0212] Statistical analysis can be performed using GraphPad PRISM software. Results can be expressed as the mean \pm standard deviation (SD).

[0213] As seen in TABLE 7, all antibodies potently inhibited rhIL-23-induced and cyno-IL-23-induced IFN γ expression in whole blood cells in a dose dependent manner. The antibodies all had IC₅₀ values in the picomolar range.

TABLE-US-00007 TABLE 7 Table of mean IC₅₀ (pM) values for IL-23 antibodies in the IFN γ human whole blood assay . rhIL-23 Cyno IL-23 antibody IC₅₀ +/- SD Repeats IC₅₀ +/- SD Repeats B 117 +/- 94 7 161 +/- 95 6 E 29 +/- 8 3 54 +/- 33 3 G 53 +/- 13 3 93 +/- 44 3 F 66 +/- 13 3 166 +/- 189 3 D 88 +/- 6 3 110 +/- 14 3 C 97 +/- 31 3 186 +/- 194 3

[0214] d) IL-22 Assay

[0215] It is known that IL-23 is a potent inducer of proinflammatory cytokines. IL-23 acts on activated and memory T cells and promotes the survival and expansion of Th17 cells which produce proinflammatory cytokines including IL-22. In this assay, human whole blood is used to assess the ability of the IL-23 antibodies to inhibit IL-23-induced IL-22 production.

[0216] A whole blood assay is conducted in the same manner as described above with the modification of using rhIL-23 or cynoIL-23 at 1 ng/ml and rhIL-18 at 10 ng/ml to induce IL-22 production. IL-22 concentration is determined by IL-22 ELISA (R&D Systems, Minneapolis, Minn.).

[0217] As seen in TABLE 8, the antibodies potently inhibited rhIL-23-induced and cyno IL-23-induced IL-22 production in whole blood cells in a dose dependent manner. The antibodies all had IC₅₀ values in the picomolar range.

TABLE-US-00008 TABLE 8 Table of mean IC₅₀ (pM) values for IL-23 antibodies in the IL-22 human whole blood assay rhIL-23 Cyno IL-23 antibody IC₅₀ +/- SD Repeats IC₅₀ +/- SD Repeats B 117 +/- 68 4 113 +/- 65 3 E 87 +/- 109 3 56 +/- 60 3 G 83 +/- 59 3 66 +/- 45 3

Example 3

Determining the Equilibrium Dissociation Constant (K_{sub}.D.) for Anti-IL-23 Antibodies Using KinExA Technology

[0218] Binding affinity of rhIL-23 to IL-23 antibodies is evaluated using a kinetic exclusion assay (KinExA assay, Sapidyne Instruments, Inc., Boise, Id.). Normal human serum (NHS)-activated Sepharose 4 fast flow beads (Amersham Biosciences, part of GE Healthcare, Uppsala, Sweden), are pre-coated with rhIL-23 and blocked with 1 M Tris buffer with 10 mg/mL BSA. 50 pM of IL-23 antibody is incubated with rhIL-23 (12 points of 1:2 dilutions starting from 800 pM) at room temperature for 72 hours before it is run through the rhIL-23-coated Sepharose beads. The amount of the bead-bound antibody was quantified by fluorescent (Cy5) labeled goat anti-human-Fc antibody (Jackson Immuno Research, West Grove, Pa.). The binding signal is proportional to the amount of free antibody at equilibrium.

[0219] The dissociation equilibrium constant (K_{sub}.D) and the association rate (K_{sub}.on) are obtained from curve fitting using KinExA Pro software. The dissociation rate (K_{sub}.off) is derived from: $K_{sub}.D = K_{sub}.off / K_{sub}.on$

[0220] As seen in TABLE 9, the antibodies have high affinity for binding to human IL-23. All had K_{sub}.D values in the low to sub pM range.

TABLE-US-00009 TABLE 9 Table of K_{sub}.D (pM), K_{sub}.on (1/MS) and K_{sub}.off (1/s) rates Antibody KD (pM) Kon (1/MS) Koff (1/s) E 0.131 9.12E+05 1.4E-07 D 0.126 1.72E+06 2.2E-07 B 3.99 1.17E+06 4.7E-06 C 2.56 1.36E+06 4.1E-06 F 2.62 5.69E+05 1.5E-06 L 1.08 3.34E+06 3.7E-06 G 2.00 4.00E+05 8.1E-07

Example 4

Structure Determination Using X-Ray Crystallography

[0221] One way to determine the structure of an antibody-antigen complex is by using X-ray crystallography, see for example, Harlow and Lane Antibodies: A Laboratory Manual Cold Spring Harbor Laboratory Press, Cold Spring Harbor, N.Y. (1990), p. 23. The crystal structure of IL-23 has been determined, (see Lupardus and Garcia, J Mol Biol, 2008, 382: 931-941) and the crystal structure of an IL-23/Fab complex has been disclosed, (see Beyer et al. J Mol Biol, 2008. 382(4): 942-55). Structural determination of IL-23 with Fab fragments of antibodies claimed herein was obtained using X-ray crystallography.

Protein for Crystallization

[0222] A recombinantly derived human IL-23 heterodimer was used for the crystallization studies (see Beyer et al., supra). The sequence of the human p19 subunit comprised of residues 20-189 of SEQ ID NO: 145, the signal sequence of SEQ ID NO:154 and a C-terminal 6-His tag SEQ ID NO:155. The sequence of the human p40 subunit was mutated from asparagine to glutamine at position 222 of SEQ ID NO:147 in order to prevent glycosylation at this site (Beyer, et al., supra).

[0223] Fabs derived from Antibody B and Antibody E were expressed on an IgG1 scaffold that incorporated a caspase cleavage site. The Fabs were processed by means of protease cleavage.

Complex Formation and Crystallization

[0224] The IL-23-Antibody B Fab complex was made by mixing a 2× molar excess of the Antibody B Fab with the human heterodimeric IL-23 described above. The complex was purified by size exclusion chromatography to remove excess Antibody B Fab and concentrated to –12 mg/ml for crystallization. The IL-23-Antibody B Fab complex crystallized in 0.1 M Hepes pH 7, 8% PEG 8000.

[0225] The IL-23-Antibody E Fab complex was made by mixing a 2× molar excess of the Antibody E Fab with the human heterodimeric IL-23 described above. The complex was methylated using a JBS Methylation Kit according to manufacturer's instructions (Jena Bioscience, Jena, Germany). The complex was then treated with PNGase to deglycosylate the protein. Following these treatments, the complex was purified by size exclusion chromatography to remove excess Antibody E Fab and concentrated to 13.5 mg/ml for crystallization. The IL-23-Antibody E Fab complex crystallized in 0.1 M Tris pH 8.5, 0.2 M magnesium chloride, 15% PEG 4000.

Data Collection and Structure Determination

[0226] IL-23-Antibody B Fab crystals grew in the P2.sub.1 space group with unit cell dimensions $a=70.93$, $b=71.27$, $c=107.37$ Å, $\beta=104.98^\circ$ and diffract to 2.0 Å resolution. The IL-23-Antibody B Fab structure was solved by molecular replacement with the program MOLREP (CCP4, *The CCP4 suite: programs for protein crystallography*. Acta Crystallogr D Biol Crystallogr, 1994. 50(Pt 5): p. 760-3) using the IL-23 structure (Beyer et al. supra) as the starting search model. Keeping the IL-23 solution fixed, an antibody variable domain was used as a search model. Keeping the IL-23-antibody variable domain solution fixed, an antibody constant domain was used as a search model. The complete structure was improved with multiple rounds of model building with Quanta and refinement with cnx (Brunger, et al., *Acta Crystallogr D Biol Crystallogr*, 1998, 54(Pt 5): p. 905-21).

[0227] Distances between protein atoms were calculated using the program PyMOL (DeLano, W. L. The PyMOL Graphics System. Palo Alto, 2002) (Schrodinger, LLC; New York, N.Y.). Amino acids were chosen if at least one atom was located within the required distance threshold to the partner protein.

[0228] Boundaries of the A, B, C and D helices of the p19 subunit of IL-23 when bound to the Antibody B Fab include A helix residues 28-47, B helix residues 86-105, C helix residues 119-134 and D helix residues 154-187 of SEQ ID NO:145.

[0229] The regions of interaction on the IL-23p19 subunit when bound to the Antibody B Fab include residues within Ser46-Glu58, Glu112-Glu123 and Pro155-Phe163 of SEQ ID NO:145.

[0230] IL-23p19 subunit amino acid residues with atoms 4 Å or less from the Antibody B Fab include Ser46, Ala47, His48, Pro49, Leu50, His53, Met54, Asp55, Glu58, Pro113, Ser114, Leu115, Leu116, Pro120, Val121, Trp156, Leu159, Leu160, Arg162 and Phe163 of SEQ ID NO:145. IL-23p19 amino acid residues with atoms between 4 Å and 5 Å from the Antibody B Fab include Val51, Arg57, Glu112, Asp118, Ser119, Gln123, Pro155 of SEQ ID NO:145.

[0231] IL-23p40 subunit amino acid residues with atoms 4 Å or less from the Antibody B Fab include Glu 122 and Lys 124 of SEQ ID NO:147.

[0232] The Antibody B Fab heavy chain amino acid residues with atoms 4 Å or less from the IL-23 heterodimer include Gly32, Gly33, Tyr34, Tyr35, His54, Asn58, Thr59, Tyr60, Lys66, Arg101, Gly102, Phe103, Tyr104 and Tyr105 of SEQ ID NO:46. The Antibody B Fab heavy chain amino acid residues with atoms 5 Å from the IL-23 heterodimer include Ser31, Gly32, Gly33, Tyr34, Tyr35, His54, Ser56, Asn58, Thr59, Tyr60, Lys66, Arg101, Gly102, Phe103, Tyr104 and Tyr105 of SEQ ID NO:46.

[0233] The Antibody B Fab light chain amino acid residues with atoms 4 Å or less from the IL-23 heterodimer include Ser30, Ser31, Trp32, Tyr49, Ser52, Ser53, Ala91, Asn92, Ser93, Phe94, and Phe96 of SEQ ID NO:15. The Antibody B Fab light chain amino acid residues with atoms 5 Å from the

IL-23 heterodimer include Ser30, Ser31, Trp32, Tyr49, Ala50, Ser52, Ser53, Ser56, Ala91, Asn92, Ser93, Phe94, and Phe96 of SEQ ID NO:15

[0234] The IL-23-Antibody E Fab complex crystals grew in the P222.sub.1 space group with unit cell dimensions $a=61.60$, $b=97.59$, $c=223.95$ Å and diffract to 3.5 Å resolution. The IL-23-Antibody E Fab complex structure was solved by molecular replacement with the program Phaser (CCP4, supra) using the IL-23 structure, an antibody variable domain, and an antibody constant domain as the three starting search models, as described above. The complete structure was improved with multiple rounds of model building with Quanta and refinement with cns (Brunger, et al., supra). The Antibody E Fab constant domain was left out of the final refined structure due to very poor electron density for that portion of the protein.

[0235] The regions of interaction on the IL-23p19 subunit identified when bound to the Antibody E Fab include residues within Ser46-His53, Glu112-Val120 and Trp156-Phe163 of SEQ ID NO:145.

[0236] IL-23p19 amino acid residues with atoms 4 Å or less from the Antibody E Fab include Ser46, Ala47, His48, Pro49, Leu50, Glu112, Pro113, Ser114, Leu115, Leu116, Pro117, Asp118, Ser119, Pro120, Trp156, Leu159, Leu160 and Phe163 of SEQ ID NO: 145. IL-23p19 amino acid residues with atoms between 4 Å and 5 Å from the Antibody E Fab include His53 of SEQ ID NO:145.

[0237] IL-23p40 amino acid residues with atoms 4 Å or less from the Antibody E Fab include Lys121, Glu 122, Pro123 and Asn 125 of SEQ ID NO:147.

[0238] The Antibody E Fab heavy chain amino acid residues with atoms 4 Å or less from the IL-23 heterodimer include Gly26, Phe27, Thr28, Ser31, Tyr53, Tyr59, Tyr102, Ser104, Ser105, Trp106, Tyr107, and Pro108 of SEQ ID NO:31. The Antibody E Fab heavy chain amino acid residues with atoms ≤ 5 Å from the IL-23 heterodimer include Gln1, Gly26, Phe27, Thr28, Ser30, Ser31, Tyr32, Trp52, Tyr53, Tyr59, Arg100, Tyr102, Thr103, Ser104, Ser105, Trp106, Tyr107, and Pro108 of SEQ ID NO:31.

[0239] The Antibody E Fab light chain amino acid residues with atoms 4 Å or less from the IL-23 heterodimer include Ala31, Gly32, Tyr33, Asp34, Tyr51, Gly52, Asn55, Lys68, and Tyr93 of SEQ ID NO:1. The Antibody B Fab light chain amino acid residues with atoms 5 Å from the IL-23 heterodimer include Thr29, Ala31, Gly32, Tyr33, Asp34, Tyr51, Gly52, Asn55, Lys68, Tyr93, and Trp100 of SEQ ID NO:1.

Example 5

Determination of IL-23-Antibody Complex Contact Residues Through Solvent Accessible Surface Area Differences

[0240] The residue contacts in the paratope (the portion of the antibody that recognizes the antigen) and the portion of the antigen that it binds bound by the paratope in a human IL-23-Antibody B Fab complex and in a human IL-23-Antibody E Fab complex were determined using solvent accessible surface area differences. The solvent accessible surface area calculations were performed using Molecular Operating Environment (Chemical Computing Group, Montreal, Quebec).

[0241] The solvent accessible surface area differences of the paratope residues in the IL-23-Antibody B Fab complex were calculated by setting the Antibody B Fab residues as the desired set. The structural information obtained in Example 4 for the IL-23-Antibody B Fab complex was used and the residue solvent accessible surface area of the amino acid residues of the Antibody B Fab in the presence of the IL-23 heterodimer were calculated and represent the “bound areas” for the set.

[0242] The residue solvent accessible surface area of each of the Antibody B Fab residues in the absence of the IL-23 antigen were calculated and represent the “free areas” of the set.

[0243] The “bound areas” were then subtracted from the “free areas” resulting in the “solvent exposed surface area difference” for each residue in the set. The Antibody B Fab residues that had no change

in surface area, or a zero difference, had no contact with the residues of the IL-23 antigen when complexed. The Antibody B Fab residues that had a difference value $0 \text{ \AA} \cdot \text{sup.2}$ were considered to be in significant contact with residues in the IL-23 antigen such that these Antibody B Fab residues were at least partially to completely occluded when the Antibody B Fab was bound to human IL-23. This set of Antibody B Fab residues make up the "covered patch", the residues involved in the structure of the interface when Antibody B Fab is bound to human IL-23, see Tables 10 and 11. The Antibody B Fab residues in this covered patch may not be involved in binding interactions with residues of the IL-23 antigen, but mutation of any single residue within the covered patch could introduce energetic differences that would impact the binding of Antibody B Fab to human IL-23. With the exception of Tyr49, all of the residues are located in the CDR regions of the Antibody B Fab light and heavy chains. These residues were also within 5 Å or less of the IL-23 antigen when bound to the Antibody B Fab, as described in Example 4.

TABLE-US-00010 TABLE 10 Solvent Accessibility Surface Area Differences for Antibody B Fab Light Chain Residue Residue Position Solvent exposed surface area AHO Number SEQ ID NO: 15 difference ($\text{\AA} \cdot \text{sup.2}$) Ser32 Ser30 44.9 Ser33 Ser31 41.1 Trp40 Trp32 79.0 Tyr57 Tyr49 40.7 Ala58 Ala50 20.3 Ser68 Ser52 43.6 Ser69 Ser53 38.9 Ser72 Ser56 19.1 Asn110 Asn92 34.0 Phe135 Phe94 51.4

TABLE-US-00011 TABLE 11 Solvent Accessibility Surface Area Differences for Antibody B Fab Heavy Chain Residue Residue Position Solvent exposed surface area AHO Number SEQ ID NO: 46 difference ($\text{\AA} \cdot \text{sup.2}$) Ser33 Ser31 18.2 Gly34 Gly32 49.5 Gly38 Gly33 33.8 Tyr39 Tyr34 51.4 Tyr40 Tyr35 30.7 His59 His54 29.5 Asn67 Asn58 66.7 Thr68 Thr59 26.0 Tyr69 Tyr60 59.4 Lys75 Lys66 32.6 Arg110 Arg101 47.2 Gly111 Gly102 21.7 Phe112 Phe103 35.5 Tyr133 Tyr104 83.0 Tyr134 Tyr105 91.7

[0244] The solvent accessible surface area differences of the residues in the IL-23-Antibody E Fab complex were calculated as described above. The Antibody E Fab residues that had a difference value $0 \text{ \AA} \cdot \text{sup.2}$ were considered to be in significant contact with residues in the IL-23 antigen and these Antibody E Fab residues were at least partially to completely occluded when the Antibody E Fab was bound to human IL-23. This set of Antibody E Fab residues make up the covered patch, the residues involved in the structure of the interface when the Antibody E Fab is bound to human IL-23, see Tables 12 and 13. The Antibody E Fab residues in this covered patch may not be involved in binding interactions with residues of the IL-23 antigen, but mutation of any single residue within the covered patch could introduce energetic differences that would impact the binding of Antibody E Fab to human IL-23. For the most part, these covered patch residues were located within the CDR regions of the Antibody E Fab heavy and light chains. These residues were also within 5 Å or less of the IL-23 antigen when bound to the Antibody E Fab, as described in Example 4.

TABLE-US-00012 TABLE 12 Solvent Accessibility Surface Area Differences for Antibody E Fab Light Chain Residue Residue Position Solvent exposed surface area AHO Number SEQ ID NO: 1 difference ($\text{\AA} \cdot \text{sup.2}$) Ala33 Ala31 11.6 Gly34 Gly32 51.2 Tyr39 Tyr33 47.2 Asp40 Asp34 36.8 Tyr57 Tyr51 16.1 Gly58 Gly52 11.1 Asn69 Asn55 29.4 Lys82 Lys68 20.1 Tyr109 Tyr93 27.3 Ser135 Ser98 11.3

TABLE-US-00013 TABLE 13 Solvent Accessibility Surface Area Differences for Antibody E Fab Heavy Chain Residue Residue Position Solvent exposed surface area AHO Number SEQ ID NO: 31 difference ($\text{\AA} \cdot \text{sup.2}$) Gln1 Gln1 41.1 Gly27 Gly26 24.6 Thr30 Thr28 82.2 Ser33 Ser31 40.7 Tyr39 Tyr32 30.7 Trp59 Trp52 11.3 Tyr60 Tyr53 44.7 Tyr69 Tyr59 42.4 Lys86 Lys76 17.4 Gly111 Gly101 12.8 Tyr112 Tyr102 103.1 Ser114 Ser104 21.0 Ser115 Ser105 91.4 Trp131 Trp106 145.0 Tyr132 Tyr107 71.6 Pro133 Pro108 20.4

[0245] The solvent accessible surface area differences of the portion of the IL-23 heterodimer bound by the paratope of the Antibody B Fab were calculated by setting the IL-23 heterodimer residues as the desired set. The structural information obtained in Example 4 for the Antibody B Fab-IL-23 complex was used and the residue solvent accessible surface area of the amino acid residues of the IL-23 heterodimer in the presence of the Antibody B Fab were calculated and represent the bound areas for the set.

[0246] The residue solvent accessible surface area of each of the IL-23 heterodimer residues in the absence of the Antibody B Fab were calculated and represent the free areas of the set.

[0247] As described above, the bound areas were subtracted from the free areas resulting in the solvent exposed surface area difference for each IL-23 residue. The IL-23 heterodimer residues that had no change in surface area, or a zero difference, had no contact with the residues of the Antibody B Fab when complexed. The IL-23 heterodimer residues that had a difference value $\geq 10 \text{ \AA}^2$ were considered to be in significant contact with residues of the Antibody B Fab and these 11-23 heterodimer residues were at least partially to completely occluded when the human IL-23 heterodimer was bound to the Antibody B Fab. This set of IL-23 heterodimer residues make up the covered patch, the residues involved in the structure of the interface when the human IL-23 heterodimer is bound to the Antibody E Fab, see Table 14. The 11-23 heterodimer residues in this covered patch may not all be involved in binding interactions with residues on the Antibody B Fab, but mutation of any single residue within the covered patch could introduce energetic differences that would impact the binding of Antibody B Fab to human IL-23. These residues are also within 4 Å or less from the Antibody B Fab, as described Example 4.

TABLE-US-00014 TABLE 14 Solvent Accessibility Surface Area Differences for IL-23 heterodimer residues Solvent exposed surface area difference (\AA^2) p19 residues (SEQ ID NO: 145) Ser46 26.5 Ala47 12.7 Pro49 59.6 Leu50 122.2 His53 47.8 Met54 13.9 Asp55 20.5 Arg57 14.6 Glu58 96.5 Glu112 29.7 Pro113 64.8 Ser114 30.0 Leu115 31.4 Leu116 60.0 Asp118 14.4 Ser119 19.7 Pro120 64.7 Pro155 19.4 Tyr156 61.9 Leu159 72.8 Leu160 27.0 Arg162 14.4 Phe163 67.5 p40 residues (SEQ ID NO: 147) Glu122 29.1 Lys124 60.9

[0248] The solvent accessible surface area differences of the portion of the IL-23 heterodimer bound by the paratope of the Antibody E Fab were calculated as described above. The IL-23 heterodimer residues that had a difference value $\geq 0 \text{ \AA}^2$ were considered to be in significant contact with residues of the Antibody E Fab and these 11-23 heterodimer residues were at least partially to completely occluded when the human IL-23 heterodimer was bound to the Antibody E Fab. This set of IL-23 heterodimer residues make up the covered patch, the residues involved in the structure of the interface when the human IL-23 heterodimer is bound to the Antibody E Fab, see Table 15. The 11-23 heterodimer residues in this covered patch may not all be involved in binding interactions with residues on the Antibody E Fab, but mutation of any single residue within the covered patch could introduce energetic differences that would impact the binding of Antibody E Fab to human IL-23. These residues are also within 5 Å or less from the Antibody E Fab, as described in Example 4.

TABLE-US-00015 TABLE 15 Solvent Accessibility Surface Area Differences for IL-23 heterodimer residues Solvent exposed surface area difference (\AA^2) p19 residues (SEQ ID NO: 145) Ser46 18.7 Ala47 14.9 Pro49 79.8 Leu50 99.5 His53 61.2 Glu112 62.8 Pro113 45.7 Ser114 69.5 Leu115 50.3 Leu116 127.2 Pro117 54.1 Asp118 37.0 Pro120 18.8 Pro155 16.9 Trp156 140.7 Leu159 21.8 Leu160 17.0 Phe163 56.6 p40 residues (SEQ ID NO: 147) Lys121 86.2 Glu122 21.8 Pro123 22.1 Asn125 26.7 Arg283 22.6

Claims

1. An isolated antigen binding protein that binds IL-23, comprising at least one heavy chain variable region comprising a CDRH1, a CDRH2 and a CDRH3 selected from the group consisting of: a) a CDRH1 that differs by no more than one amino acid substitution, insertion or deletion from a CDRH1 as shown in TABLE 3; b) a CDRH2 that differs by no more than three amino acid substitutions, insertions and/or deletions from a CDRH2 as shown in TABLE 3; c) a CDRH3 that differs by no more than three amino acid substitutions, insertions and/or deletions from a CDRH3 as shown in TABLE 3; and comprising at least one light chain variable region comprising a CDRL1, a CDRL2 and a CDRL3 selected from the group consisting of: d) a CDRL1 that differs by no more than three amino acid substitutions, insertions and/or deletions from a CDRL1 as shown in TABLE 3; e) a CDRL2 that differs by no more than one amino acid substitution, insertion or deletion from a CDRL2 as shown in TABLE 3; f) a CDRL3 that differs by no more than one amino acid substitution, insertion or deletion from a CDRL3 as shown in TABLE 3.

2. An isolated antigen binding protein of claim 1, comprising at least one heavy chain variable region comprising a CDRH1, a CDRH2 and a CDRH3 selected from the group consisting of: a) a CDRH1 that differs by no more than one amino acid substitution, insertion or deletion from a CDRH1 as shown in TABLE 3; b) a CDRH2 that differs by no more than two amino acid substitutions, insertions and/or deletions from a CDRH2 as shown in TABLE 3; c) a CDRH3 that differs by no more than two amino acid substitutions, insertions and/or deletions from a CDRH3 as shown in TABLE 3; and also comprising at least one light chain variable region comprising a CDRL1, a CDRL2 and a CDRL3 selected from the group consisting of: d) a CDRL1 that differs by no more than two amino acid substitutions, insertions and/or deletions from a CDRL1 as shown in TABLE 3; e) a CDRL2 that differs by no more than one amino acid substitution, insertions or deletion from a CDRL2 as shown in TABLE 3; f) a CDRL3 that differs by no more than one amino acid substitution, insertions or deletion from a CDRL3 as shown in TABLE 3.

3. An isolated antigen binding protein of claim 1, comprising at least one heavy chain variable region comprising a CDRH1, CDRH2 and a CDRH3 selected from the group consisting of: a) a CDRH1 that differs by no more than one amino acid substitution, insertion or deletion from a CDRH1 as shown in TABLE 3; b) a CDRH2 that differs by no more than one amino acid substitution, insertion or deletion from a CDRH2 as shown in TABLE 3; c) a CDRH3 that differs by no more than one amino acid substitution, insertion or deletion from a CDRH3 as shown in TABLE 3; and also comprising at least one light chain variable region comprising a CDRL1, a CDRL2 and a CDRL3 selected from the group consisting of: d) a CDRL1 that differs by no more than one amino acid substitution, insertion or deletion from a CDRL1 as shown in TABLE 3; e) a CDRL2 that differs by no more than one amino acid substitution, insertion or deletion from a CDRL2 as shown in TABLE 3; f) a CDRL3 that differs by no more than one amino acid substitution, insertion or deletion from a CDRL3 as shown in TABLE 3.

4. An isolated antigen binding protein that binds IL-23 selected from the group consisting of a) an antigen binding protein having CDRH1 of SEQ ID NO:129, CDRH2 of SEQ ID NO:132, CDRH3 of SEQ ID NO:136, and CDRL1 of SEQ ID NO:123, CDRL2 of SEQ ID NO:81, and CDRL3 of SEQ ID NO: 76; b) an antigen binding protein having CDRH1 of SEQ ID NO:131, CDRH2 of SEQ ID NO: 134, CDRH3 of SEQ ID NO:137 and CDRL1 of SEQ ID NO:124, CDRL2 of SEQ ID NO:126 and CDRL3 of SEQ ID NO:128; c) a) an antigen binding protein having CDRH1 of SEQ ID NO:130, CDRH2 of SEQ ID NO:133, CDRH3 of SEQ ID NO:99 and CDRL1 of SEQ ID NO:68, CDRL2 of SEQ ID NO:69, and CDRL3 of SEQ ID NO:67; and d) an antigen binding protein having CDRH1 SEQ ID NO:91, CDRH2 SEQ ID NO: 135, CDRH3 SEQ ID NO:138 and CDRL1 SEQ ID NO:125, CDRL2 SEQ ID NO:127, and CDRL3 SEQ ID NO:64.

5. An isolated antigen binding protein of claim 1 comprising: a CDRH1 selected from the group consisting of SEQ ID NO: 91, 94, 97, 100, and 103; a CDRH2 selected from the group consisting of SEQ ID NO:92, 95, 98, 101, 104, 107, and 110; a CDRH3 selected from the group consisting of SEQ ID NO: 93, 96, 99, 102, and 105; a CDRL1 selected from the group consisting of SEQ ID NO: 62, 65, 68, 71, and 74; a CDRL2 selected from the group consisting of SEQ ID NO:63, 66, 69, 72, 75, and 78; and a CDRL3 selected from the group consisting of SEQ ID NO:64, 67, 70 and 73.

6. An isolated antigen binding protein of claim 1, comprising: a CDRH1 selected from the group consisting of SEQ ID NO: 91, 106, 109, 112, and 115; a CDRH2 selected from the group consisting of SEQ ID NO: 113, 116, 118, 120, 121, and 122; a CDRH3 selected from the group consisting of SEQ ID NO: 108, 111, 114, 117, and 119; a CDRL1 selected from the group consisting of SEQ ID NO: 77, 80, 83, 85, 86, 87, 88, 89 and 90; a CDRL2 is SEQ ID NO: 81; and a CDRL3 selected from the group consisting of SEQ ID NO: 76, 79, 82 and 84.

7. An isolated antigen binding protein that binds IL-23 comprising at least one heavy chain variable region and at least one light chain variable region, selected from the group consisting of: a) a heavy chain variable region comprising amino acid residues 31-35, 50-65 and 99-113 of SEQ ID NO:31; and a light chain variable region comprising amino acid residues 23-36, 52-58 and 91-101 of SEQ ID NO:1; b) a heavy chain variable region comprising amino acid residues 31-35, 50-65 and 99-110 of SEQ ID NO:34 and heavy chain variable region comprising amino acid residues 31-35, 50-66 and 99-110 of SEQ ID NO:36; and a light chain variable region comprising amino acid residues 23-36, 52-62

and 97-105 of SEQ ID NO:4; c) a heavy chain variable region comprising amino acid residues 31-35, 50-66 and 99-114 of SEQ ID NO:38; and a light chain variable region comprising amino acid residues 23-34, 50-61 and 94-106 of SEQ ID NO:7; d) a heavy chain variable region comprising amino acid residues 31-35, 50-66 and 99-114 of SEQ ID NO:40; and a light chain variable region comprising amino acid residues 24-34, 50-56 and 94-106 of SEQ ID NO:9; e) a heavy chain variable region comprising amino acid residues 31-35, 50-66 and 99-114 of SEQ ID NO:42; and a light chain variable region comprising amino acid residues 23-34, 50-61 and 94-106 of SEQ ID NO:11; f) a heavy chain variable region comprising amino acid residues 31-35, 50-65 and 98-107 of SEQ ID NO:44; and a light chain variable region comprising amino acid residues 24-34, 50-56 and 89-97 of SEQ ID NO:13; g) a heavy chain variable region comprising amino acid residues 31-37, 52-67 and 100-109 of SEQ ID NO:46 or 153; and a light chain variable region comprising amino acid residues 24-34, 50-56 and 89-97 of SEQ ID NO:15; h) a heavy chain variable region comprising amino acid residues 31-37, 52-67 and 100-109 of SEQ ID NO:48; and a light chain variable region comprising amino acid residues 24-34, 50-56 and 89-97 of SEQ ID NO:17; i) a heavy chain variable region comprising amino acid residues 31-37, 52-67 and 101-109 of SEQ ID NO:50; and a light chain variable region comprising amino acid residues 24-34, 50-56 and 89-97 of SEQ ID NO:19; j) a heavy chain variable region comprising amino acid residues 31-35, 50-65 and 98-107 of SEQ ID NO: 52; and a light chain variable region comprising amino acid residues 24-34, 50-56 and 98-107 of SEQ ID NO:21; k) a heavy chain variable region comprising amino acid residues 31-37, 52-67 and 100-109 of SEQ ID NO:54; and a light chain variable region comprising amino acid residues 24-34, 50-56 and 89-97 of SEQ ID NO:23; l) a heavy chain variable region comprising amino acid residues 31-37, 52-67 and 100-109 of SEQ ID NO:56; and a light chain variable region comprising amino acid residues 24-34, 50-56 and 89-97 of SEQ ID NO:25; and m) a heavy chain variable region comprising amino acid residues 31-37, 52-57 and 100-109 of SEQ ID NO:58; and a light chain variable region comprising amino acid residues 24-34, 50-56 and 89-97 of SEQ ID NO:27.

8. An isolated antigen-binding protein of claim 1 that comprises at least one heavy chain variable region and at least one light chain variable region.

9. An isolated antigen-binding protein of claim 1 that comprises at least two heavy chain variable regions and at least two light chain variable regions.

10. An isolated antigen binding protein that binds IL-23 comprising a heavy chain variable region and a light chain variable region, wherein the heavy chain variable region sequence differs by no more than 13 amino acid substitutions, additions and/or deletions from a heavy chain variable region sequence as shown in TABLE 2; and wherein the light chain variable region sequence differs by no more than 13 amino acid substitutions, additions and/or deletions from a light chain variable region sequence as shown in TABLE 1.

11. An isolated antigen binding protein of claim 10, wherein the heavy chain variable region sequence differs by no more than 11 amino acid substitutions, insertions and/or deletions from a heavy chain variable region sequence as shown in TABLE 2.

12. An isolated antigen binding protein of claim 10, wherein the heavy chain variable region sequence differs by no more than 5 amino acid substitutions, insertions and/or deletions from a heavy chain variable region sequence as shown in TABLE 2.

13. An isolated antigen binding protein of claim 10, wherein the heavy chain variable region sequence differs by no more than 2 amino acid substitutions, insertions and/or deletions from a heavy chain variable region sequence as shown in TABLE 2.

14. An isolated antigen binding protein of claim 10, wherein the heavy chain variable region sequence differs by no more than 1 amino acid substitution, insertion or deletion from a heavy chain variable region sequence as shown in TABLE 2.

15. An isolated antigen binding protein of claim 10, wherein the light chain variable region sequence differs by no more than 7 amino acid substitutions, insertions and/or deletions from a light chain

variable region sequence as shown in TABLE 1.

16. An isolated antigen binding protein of claim 10, wherein the light chain variable region sequence differs by no more than 4 amino acid substitutions, insertions and/or deletions from a light chain variable region sequence as shown in TABLE 1.

17. An isolated antigen binding protein of claim 10, wherein the light chain variable region sequence differs by no more than 2 amino acid substitutions, insertions and/or deletions from a light chain variable region sequence as shown in TABLE 1.

18. An isolated antigen binding protein of claim 10, wherein the light chain variable region sequence differs by no more than 1 amino acid substitutions, insertions and/or deletions from a light chain variable region sequence as shown in TABLE 1.

19. An isolated antigen binding protein that binds IL-23 selected from the group consisting of a) a heavy chain variable region of SEQ ID NO:140 and a light chain variable region of SEQ ID NO: 30; b) a heavy chain variable region of SEQ ID NO:141 and a light chain variable region of SEQ ID NO:61; c) a heavy chain variable region of SEQ ID NO:142 and a light chain variable region of SEQ ID NO:4; and d) a heavy chain variable region of SEQ ID NO:143 and a light chain variable region of SEQ ID NO:139.

20. An isolated antigen binding protein that binds IL-23 comprising a heavy chain variable region comprising of an amino acid sequence having at least 90% sequence identity to SEQ ID NO:31, 34, 36, 38, 40, 42, 44, 46, 48, 50, 52, 54, 56 and 58; and a light chain variable region comprising an amino acid sequence having at least 90% sequence identity to SEQ ID NO: 1, 4, 7, 9, 11, 13, 15, 17, 19, 21, 23, 25 and 27.

21. An isolated antigen binding protein of claim 20, a heavy chain variable region comprising of an amino acid sequence having at least 95% sequence identity to SEQ ID NO:31, 34, 36, 38, 40, 42, 44, 46, 48, 50, 52, 54, 56 and 58; and a light chain variable region comprising an amino acid sequence having at least 95% sequence identity to SEQ ID NO: 1, 4, 7, 9, 11, 13, 15, 17, 19, 21, 23, 25 and 27.

22. An isolated antigen binding protein of claim 20, comprising a heavy chain variable region comprising of an amino acid sequence having at least 97% sequence identity to SEQ ID NO:31, 34, 36, 38, 40, 42, 44, 46, 48, 50, 52, 54, 56 and 58; and a light chain variable region comprising an amino acid sequence having at least 97% sequence identity to SEQ ID NO: 1, 4, 7, 9, 11, 13, 15, 17, 19, 21, 23, 25 and 27.

23. An isolated antigen binding protein of claim 20, comprising a heavy chain variable region comprising of an amino acid sequence having at least 98% sequence identity to SEQ ID NO:31, 34, 36, 38, 40, 42, 44, 46, 48, 50, 52, 54, 56 and 58; and a light chain variable region comprising an amino acid sequence having at least 98% sequence identity to SEQ ID NO: 1, 4, 7, 9, 11, 13, 15, 17, 19, 21, 23, 25 and 27.

24. An isolated antigen binding protein of claim 20, comprising a heavy chain variable region selected from the group consisting of SEQ ID NO: 44, 46, 48, 50, 52, 54, 56, 58 and 153, and a light chain variable region selected from the group consisting of SEQ ID NO:13, 15, 17, 19, 21, 23, 25, and 27.

25. An isolated antigen binding protein of claim 20, comprising a heavy chain variable region selected from the group consisting of SEQ ID NO: 31, 34, 36, 38, 40 and 42, and a light chain variable region selected from the group consisting of SEQ ID NO: 1, 4, 7, 9 and 11.

26. An isolated antigen binding protein that binds IL-23 comprising a heavy chain variable region and a light chain variable region selected from the group consisting of: a) a heavy chain variable region of SEQ ID NO:31 and a light chain variable region of SEQ ID NO:1; b) a heavy chain variable region of SEQ ID NO:34 or 36 and a light chain variable region of SEQ ID NO:4; c) a heavy chain variable region of SEQ ID NO:38 and a light chain variable region of SEQ ID NO: 7; d) a heavy chain variable region of SEQ ID NO:40 and a light chain variable region of SEQ ID NO:9; e) a heavy chain variable

region of SEQ ID NO:42 and a light chain variable region of SEQ ID NO: 11; f) a heavy chain variable region of SEQ ID NO:44 and a light chain variable region of SEQ ID NO:13; g) a heavy chain variable region of SEQ ID NO:46 or 153 and a light chain variable region of SEQ ID NO:15; h) a heavy chain variable region of SEQ ID NO:48 and a light chain variable region of SEQ ID NO:17; i) a heavy chain variable region of SEQ ID NO:50 and a light chain variable region of SEQ ID NO: 19; j) a heavy chain variable region of SEQ ID NO:52 and a light chain variable region of SEQ ID NO:21; k) a heavy chain variable region of SEQ ID NO:54 and a light chain variable region of SEQ ID NO:23; l) a heavy chain variable region of SEQ ID NO:56 and a light chain variable region of SEQ ID NO:25; and m) a heavy chain variable region of SEQ ID NO:58 and a light chain variable region of SEQ ID NO:27.

27. An isolated antigen binding protein that binds human IL-23, wherein the covered patch formed when said antigen binding protein is bound to human IL-23 comprises residue contacts 30, 31, 32, 49, 50, 52, 53, 56, 92 and 94 of SEQ ID NO:15, wherein said residue contacts have a difference value of greater than or equal to 10 \AA^2 as determined by solvent exposed surface area.

28. An isolated antigen binding protein that binds human IL-23, wherein the covered patch formed when said antigen binding protein is bound to human IL-23 comprises residue contacts 31-35, 54, 58-60, 66, and 101-105 of SEQ ID NO:46, wherein said residue contacts have a difference value of greater than or equal to 10 \AA^2 as determined by solvent exposed surface area.

29. An isolated antigen binding protein that binds human IL-23, wherein the covered patch formed when said antigen binding protein is bound to human IL-23 comprises residue contacts 31-34, 51, 52, 55, 68, 93 and 98 of SEQ ID NO:1, wherein said residue contacts have a difference value of greater than or equal to 10 \AA^2 as determined by solvent exposed surface area.

30. An isolated antigen binding protein that binds human IL-23, wherein the covered patch formed when said antigen binding protein is bound to human IL-23 comprises residue contacts 1, 26, 28, 31, 32, 52, 53, 59, 76, 101, 102 and 104-108 of SEQ ID NO:31, wherein said residue contacts have a difference value of greater than or equal to 10 \AA^2 as determined by solvent exposed surface area.

31. An isolated antigen binding protein that binds human IL-23, wherein when said antigen binding protein is bound to human IL-23, said antigen binding protein is 5 \AA or less from residues 32-35, 54, 58-60, 66 and 101-105 of SEQ ID NO:46, as determined by X-ray crystallography.

32. An isolated antigen binding protein according to claim 31, wherein when said antigen binding protein is bound to human IL-23, said antigen binding protein is 5 \AA or less from residues 31-35, 54, 56, 58-60, 66 and 101-105 of SEQ ID NO:46.

33. An isolated antigen binding protein that binds human IL-23, wherein when said antigen binding protein is bound to human IL-23, said antigen binding protein is 5 \AA or less from residues 30-32, 49, 52, 53, 91-94 and 96 of SEQ ID NO:15, as determined by X-ray crystallography.

34. An isolated antigen binding protein according to claim 33, wherein when said antigen binding protein is bound to human IL-23, said antigen binding protein is 5 \AA or less from residues 30-32, 49, 50, 52, 53, 56, 91-94 and 96 of SEQ ID NO:15.

35. An isolated antigen binding protein that binds human IL-23, wherein when said antigen binding protein is bound to human IL-23, said antigen binding protein is 5 \AA or less from residues 26-28, 31, 53, 59, 102 and 104-108 of SEQ ID NO:31, as determined by X-ray crystallography.

36. An isolated antigen binding protein according to claim 35, wherein when said antigen binding protein is bound to human IL-23, said antigen binding protein is 5 \AA or less from residues 1, 26-28, 30-32, 52, 53, 59, 100, and 102-108 of SEQ ID NO:31.

37. An isolated antigen binding protein that binds human IL-23, wherein when said antigen binding protein is bound to human IL-23, said antigen binding protein is 5 \AA or less from residues 31-34, 51, 52, 55, 68 and 93 of SEQ ID NO:1 as determined by X-ray crystallography.

- 38.** An isolated antigen binding protein according to claim 37, wherein when said antigen binding protein is bound to human IL-23, said antigen binding protein is 5 Å or less from residues 29, 31-34, 51, 52, 55, 68, 93 and 100 of SEQ ID NO:1.
- 39.** An isolated antigen binding protein of claim 1, 4, 7, 10, 19, 20, 26-31, 33, 35 or 37, wherein said antigen binding protein is an antibody.
- 40.** An isolated antigen binding protein of claim 39, wherein said antibody is a monoclonal antibody, a recombinant antibody, a human antibody, a humanized antibody, a chimeric antibody, a multispecific antibody, or an antibody fragment thereof.
- 41.** An isolated antigen binding protein of claim 40, wherein said antibody fragment is a Fab fragment, a Fab' fragment, a F(ab').sub.2 fragment, a Fv fragment, a diabody, or a single chain antibody molecule.
- 42.** An isolated antigen binding protein of claim 40, wherein said antigen binding protein is a human antibody.
- 43.** An isolated antigen binding protein of claim 40, wherein said antigen binding protein is a monoclonal antibody.
- 44.** An isolated antigen binding protein of claim 39, wherein said antigen binding protein is of the IgG1-, IgG2- IgG3- or IgG4-type.
- 45.** An isolated antigen binding protein of claim 44, wherein said antigen binding protein is of the IgG1- or IgG2-type.
- 46.** An isolated nucleic acid molecule encoding an antigen binding protein of claim 1, 4, 7, 10, 19, 20 or 26.
- 47.** An isolated nucleic acid molecule of claim 46, wherein at least one heavy chain variable region is encoded by an isolated nucleic acid molecule selected from the group consisting of SEQ ID NOs:32, 35, 37, 39, 41, 43, 45, 47, 49, 51, 53, 55, 57, 59 and 152 and at least one light chain variable region is encoded by an isolated nucleic acid molecule selected from the group consisting of SEQ ID NOs:2, 5, 6, 8, 10, 12, 14, 16, 18, 20, 22, 24, 26, and 28.
- 48.** A nucleic acid molecule according to claim 47, wherein said nucleic acid molecule is operably linked to a control sequence.
- 49.** A vector comprising a nucleic acid molecule according to claim 46.
- 50.** A host cell comprising the nucleic acid molecule according to claim 46.
- 51.** A host cell comprising the vector according to claim 49.
- 52.** An isolated polynucleotide sufficient for use as a hybridization probe, PCR primer or sequencing primer that is a fragment of the nucleic acid molecule of claim 47 or its complement.
- 53.** A method of making the antigen binding protein of claim 1, 4, 7, 10, 19, 20 or 26 comprising the step of preparing said antigen binding protein from a host cell that secretes said antigen binding protein.
- 54.** An isolated antigen binding protein that binds human IL-23, wherein the covered patch formed when said antigen binding protein is bound to human IL-23 comprises a residue contact within residues 46-58, a residue contact within residues 112-120, and a residue contact within residues 155-163 of the human IL-23p19 subunit as described in SEQ ID NO:145, wherein said residue contact has a difference value greater than or equal to 10 Å.sup.2 as determined by solvent exposed surface area.

- 55.** An isolated antigen binding protein according to claim 54, wherein the covered patch formed when said antigen binding protein binds to human IL-23 further comprises a residue contact within residues 121-125 of the human IL-23p40 subunit as described in SEQ ID NO:147, wherein said residue contact has a difference value greater than or equal to 10 \AA^2 as determined by solvent exposed surface area.
- 56.** An isolated antigen binding protein that binds human IL-23, wherein when said antigen binding protein is bound to human IL-23 said antigen binding protein is 5 \AA or less from a residue within residues 46-58, from a residue within residues 112-123, and from a residue within residues 155-163 of the human IL-23p19 subunit as described in SEQ ID NO:145, as determined by X-ray crystallography.
- 57.** An isolated antigen binding protein according to claim 56, wherein when said antigen binding protein is bound to human IL-23 said antigen binding protein is 5 \AA or less from a residue within residues 121-125, of the human IL-23p40 subunit as described in SEQ ID NO:147, as determined by X-ray crystallography.
- 58.** An isolated antigen binding protein of claim 1, 4, 7, 10, 19, 20, 26-31, 33, 35, 37, 54 or 56 wherein said antigen binding protein has at least one property selected from the group consisting of: a) reducing human IL-23 activity; b) reducing production of a proinflammatory cytokine; c) binding to human IL-23 with a $K_{\text{sub.D}}$ of $\leq 5 \times 10^{-8} \text{ M}$; d) having a $K_{\text{sub.off}}$ rate of $\leq 5 \times 10^{-6} \text{ 1/s}$; and e) having an IC_{50} of $\leq 400 \text{ pM}$.
- 59.** A pharmaceutical composition comprising at least one antigen binding protein of claim 1, 4, 7, 10, 19, 20 or 26 and pharmaceutically acceptable excipient.
- 60.** A pharmaceutical composition of claim 59, further comprises a labeling group or an effector group.
- 61.** A pharmaceutical composition of claim 60, wherein said labeling group is selected from the group consisting of isotopic labels, magnetic labels, redox active moieties, optical dyes, biotinylated groups and predetermined polypeptide epitopes recognized by a secondary reporter.
- 62.** A pharmaceutical composition of claim 60, wherein said effector group is selected from the group consisting of a radioisotope, radionuclide, a toxin, a therapeutic group and a chemotherapeutic group.
- 63.** An isolated antigen binding protein of claim 60, wherein said antigen binding protein is coupled to a labeling group.
- 64.** A method for treating or preventing a condition associated with IL-23 in a patient, comprising administering to a patient in need thereof an effective amount of at least one isolated antigen binding protein of claim 1, 4, 7, 10, 19, 20 or 26.
- 65.** A method of claim 64, wherein the condition is selected from the group consisting of an inflammatory disorder, a rheumatic disorder, an autoimmune disorder, an oncological disorder and a gastrointestinal disorder.
- 66.** A method of claim 65, wherein the condition is selected from the group consisting of multiple sclerosis, rheumatoid arthritis, cancer, psoriasis, inflammatory bowel disease, Crohn's disease, ulcerative colitis, systemic lupus erythematosus, psoriatic arthritis, autoimmune myocarditis; type 1 diabetes and ankylosing spondylitis.
- 67.** A method of claim 64, wherein the isolated antigen-binding protein is administered alone or as a combination therapy.
- 68.** A method of reducing IL-23 activity in a patient comprising administering an effective amount of at least one antigen binding protein of claim 1, 4, 7, 10, 19, 20 or 26.
- 69.** A method of reducing IL-23 activity of claim 68, wherein said IL-23 activity is inducing production of a proinflammatory cytokine.

Sequence Listing

CWU

1

SEQUENCE LISTING NEW RULES

1551111PRTHomo sapiens 1Gln Ser Val Leu Thr Gln Pro Pro Ser Val Ser Gly Ala Pro Gly Gln1 5 10
15Arg Val Thr Ile Ser Cys Thr Gly Ser Ser Ser Asn Thr Gly Ala Gly 20 25 30Tyr Asp Val His Trp Tyr
Gln Gln Val Pro Gly Thr Ala Pro Lys Leu 35 40 45Leu Ile Tyr Gly Ser Gly Asn Arg Pro Ser Gly Val Pro
Asp Arg Phe 50 55 60Ser Gly Ser Lys Ser Gly Thr Ser Ala Ser Leu Ala Ile Thr Gly Leu65 70 75 80Gln
Ala Glu Asp Glu Ala Asp Tyr Tyr Cys Gln Ser Tyr Asp Ser Ser 85 90 95Leu Ser Gly Trp Val Phe Gly
Gly Gly Thr Arg Leu Thr Val Leu 100 105 1102333DNAHomo sapiens 2cagtctgtgc tgacgcagcc
gccctcagtg tctggggccc cagggcagag ggtcaccatc 60tctctgactg ggagcagctc caacaccggg gcagggtatg
atgtacactg gtaccagcaa 120gttccaggaa cagcccccaa actcctcatt tatggtagcg gcaatcggcc ctcaggggtc
180cctgaccgat tctctggctc caagtctggc acctcagcct cctggccat cactggactc 240caggctgagg atgaggctga
ttactactgc cagtcctatg acagcagcct gactggttg 300gtgttcggcg gagggaccag gctgaccgtc ctg
3333111PRTHomo sapiens 3Gln Ser Val Leu Thr Gln Pro Pro Ser Val Ser Gly Ala Pro Gly Gln1 5 10
15Arg Val Thr Ile Ser Cys Thr Gly Ser Ser Ser Asn Ile Gly Ala Gly 20 25 30Tyr Asp Val His Trp Tyr Gln
Gln Leu Pro Gly Thr Ala Pro Lys Leu 35 40 45Leu Ile Tyr Gly Ser Asn Asn Arg Pro Ser Gly Val Pro
Asp Arg Phe 50 55 60Ser Gly Ser Lys Ser Gly Thr Ser Ala Ser Leu Ala Ile Thr Gly Leu65 70 75 80Gln
Ala Glu Asp Glu Ala Asp Tyr Tyr Cys Gln Ser Tyr Asp Ser Ser 85 90 95Leu Ser Gly Trp Val Phe Gly
Gly Gly Thr Lys Leu Thr Val Leu 100 105 1104115PRTHomo sapiens 4Gln Ala Val Leu Thr Gln Pro
Ser Ser Leu Ser Ala Ser Pro Gly Ala1 5 10 15Ser Ala Ser Leu Thr Cys Thr Leu Arg Ser Gly Ile Asn Val
Gly Thr 20 25 30Tyr Arg Ile Tyr Trp Tyr Gln Gln Lys Pro Gly Ser Pro Pro Gln Tyr 35 40 45Leu Leu Arg
Tyr Lys Ser Asp Ser Asp Lys Gln Gln Gly Ser Gly Val 50 55 60Pro Ser Arg Phe Ser Gly Ser Lys Asp
Ala Ser Ala Asn Ala Gly Ile65 70 75 80Leu Leu Ile Ser Gly Leu Gln Ser Glu Asp Glu Ala Asp Tyr Tyr
Cys 85 90 95Met Ile Trp His Ser Ser Ala Ser Val Phe Gly Gly Gly Thr Lys Leu 100 105 110Thr Val Leu
1155345DNAHomo sapiens 5caggctgtgc tgactcagcc gtcttccctc tctgcatctc ctggagcatc agccagctc
60acctgcacct tacgcaagt ggtacctaca ggatatactg gtaccagcag 120aagccaggga gtctccccca
gtatctctg aggtacaaat cagactcaga taagcagcag 180ggctctggag tccccagccg ctctctgga tcaaagatg
cttcggccaa tgcaggatt 240ttactcatct ctgggtcca gctgaggat gaggctgact attactgtat gatttgac
300agcagcgctt cggtattcgg cggaggacc aagctgaccg tcta 345614PRTHomo sapiens 6Glu Asn Thr Val
Thr Ile Tyr Tyr Asn Tyr Gly Met Asp Val1 5 107116PRTHomo sapiens 7Gln Pro Val Leu Thr Gln Pro
Pro Ser Ala Ser Ala Ser Leu Gly Ala1 5 10 15Ser Val Thr Leu Thr Cys Thr Leu Asn Ser Gly Tyr Ser
Asp Tyr Lys 20 25 30Val Asp Trp Tyr Gln Gln Arg Pro Gly Lys Gly Pro Arg Phe Val Met 35 40 45Arg
Val Gly Thr Gly Gly Ile Val Gly Ser Lys Gly Asp Gly Ile Pro 50 55 60Asp Arg Phe Ser Val Leu Gly Ser
Gly Leu Asn Arg Tyr Leu Thr Ile65 70 75 80Lys Asn Ile Gln Glu Glu Asp Glu Ser Asp Tyr His Cys Gly
Ala Asp 85 90 95His Gly Ser Gly Ser Asn Phe Val Tyr Val Phe Gly Thr Gly Thr Lys 100 105 110Val Thr
Val Leu 1158348DNAHomo sapiens 8cagcctgtgc tgactcagcc accttctgca tcagcctccc tgggagcctc
ggtcacactc 60acctgcaccc tgaacagcgg ctacagtgat tataaagtgg actggtacca gcagagacca 120gggaagggcc
cccgtttgt gatgagtg ggcactggg ggattgtgg atccaagggg 180gatggcatcc ctgatcgctt ctactcttg
ggctcaggcc tgaatcggt cctgaccatc 240aagaatatcc aggaagagga tgagagtgc taccactgtg gggcagacca
tggcagtggg 300agcaacttcg tgtatgtctt cggaactggg accaaggta cctccta 3489116PRTHomo sapiens
9Gln Pro Val Leu Thr Gln Pro Pro Ser Ala Ser Ala Ser Leu Gly Ala1 5 10 15Ser Val Thr Leu Thr Cys
Thr Leu Ser Ser Gly Tyr Ser Asp Tyr Lys 20 25 30Val Asp Trp Tyr Gln Gln Arg Pro Gly Lys Gly Pro Arg
Phe Val Met 35 40 45Arg Val Gly Thr Gly Gly Ile Val Gly Ser Lys Gly Glu Gly Ile Pro 50 55 60Asp Arg
Phe Ser Val Leu Gly Ser Gly Leu Asn Arg Tyr Leu Thr Ile65 70 75 80Lys Asn Ile Gln Glu Glu Asp Glu
Ser Asp Tyr His Cys Gly Ala Asp 85 90 95His Gly Ser Gly Asn Asn Phe Val Tyr Val Phe Gly Thr Gly
Thr Lys 100 105 110Val Thr Val Leu 11510348DNAHomo sapiens 10cagcctgtgc tgactcagcc accttctgca
tcagcctccc tgggagcctc ggtcacactc 60acctgcaccc tgagcagcgg ctacagtgat tataaagtgg actggtacca
gcagagacca 120gggaagggcc cccgtttgt gatgagtg ggcactggg ggattgtgg atccaagggg 180gaaggcatcc
ctgatcgctt ctactcttg ggctcaggcc tgaatcggt cctgaccatc 240aagaacatcc aggaagagga tgagagtgc
taccactgtg gggcagacca tggcagtggg 300aacaacttcg tgtatgtctt cggaactggg accaaggta cctccta
34811116PRTHomo sapiens 11Gln Pro Glu Leu Thr Gln Pro Pro Ser Ala Ser Ala Ser Leu Gly Ala1 5
10 15Ser Val Thr Leu Thr Cys Thr Leu Ser Ser Gly Tyr Ser Asp Tyr Lys 20 25 30Val Asp Trp Tyr Gln

Leu Arg Pro Gly Lys Gly Pro Arg Phe Val Met 35 40 45Arg Val Gly Thr Gly Gly Thr Val Gly Ser Lys Gly
Glu Gly Ile Pro 50 55 60Asp Arg Phe Ser Val Leu Gly Ser Gly Leu Asn Arg Ser Leu Thr Ile65 70 75
80Lys Asn Ile Gln Glu Glu Asp Glu Ser Asp Tyr His Cys Gly Ala Asp 85 90 95His Gly Ser Gly Ser Asn
Phe Val Tyr Val Phe Gly Thr Gly Thr Lys 100 105 110Val Thr Val Leu 11512348DNAHomo sapiens
12cagcctgagt tgactcagcc acctctgca tcagcctccc tgggagcctc ggtcacactc 60acctgcaccc tgagcagcgg
ctacagtgat tataaagtgg actggtacca gctgagacca 120gggaagggcc cccggtttgt gatgcgagtg ggcactggtg
ggactgttg atccaagggg 180gaagcatcc ctgatcgctt ctactgtt ggctcaggcc tgaatcggct cctgaccatc
240aagaacatcc aggaagagga tgagagtgc taccactgtg gggcagacca tggcagtggg 300agcaacttcg tgatgtctt
cggaactggg accaaggtca ccgtccta 34813107PRTTHomo sapiens 13Asp Ile Gln Leu Thr Pro Ser Pro Ser
Ser Val Ser Ala Ser Val Gly1 5 10 15Asp Arg Val Thr Ile Thr Cys Arg Ala Ser Gln Gly Ile Ala Gly Trp 20
25 30Leu Ala Trp Tyr Gln Gln Lys Pro Gly Lys Ala Pro Lys Leu Leu Ile 35 40 45Tyr Ala Ala Ser Ser Leu
Gln Ser Gly Val Pro Ser Arg Phe Ser Gly 50 55 60Ser Gly Ser Gly Thr Asp Phe Thr Leu Thr Ile Ser
Ser Leu Gln Pro65 70 75 80Glu Asp Phe Ala Thr Tyr Tyr Cys Gln Gln Ala Asp Ser Phe Pro Pro 85 90
95Thr Phe Gly Gly Gly Thr Lys Val Glu Ile Lys 100 10514321DNAHomo sapiens 14gacatccagt
tgaccccgct tccatcttcc gtgtctgcat ctgtaggaga cagagtcacc 60atcactgtc gggcgagtca gggattgcc ggctggttag
cctggtatca gcagaaacca 120gggaaagccc ctaagctct gatctatgct gcatccagtt tgcaaagtgg ggtcccatca
180aggttcagcg gcagtggatc tgggacagat ttactctca ccatcagcag cctgcagcct 240gaagatttg caacttacta
ttgtcaacag gctgacagtt tccctccac ttcggcgga 300gggaccaagg tggagatcaa a 32115107PRTTHomo
sapiens 15Asp Ile Gln Met Thr Gln Ser Pro Ser Ser Val Ser Ala Ser Val Gly1 5 10 15Asp Arg Val Thr
Ile Thr Cys Arg Ala Ser Gln Val Ile Ser Ser Trp 20 25 30Leu Ala Trp Tyr Gln Gln Lys Pro Gly Lys Ala
Pro Ser Leu Leu Ile 35 40 45Tyr Ala Ala Ser Ser Leu Gln Ser Gly Val Pro Ser Arg Phe Ser Gly 50 55
60Ser Val Ser Gly Thr Asp Phe Thr Leu Thr Ile Ser Ser Leu Gln Pro65 70 75 80Glu Asp Phe Ala Thr
Tyr Tyr Cys Gln Gln Ala Asn Ser Phe Pro Phe 85 90 95Thr Phe Gly Pro Gly Thr Lys Val Asp Phe Lys
100 10516321DNAHomo sapiens 16gacatccaga tgaccagtc tccatcttcc gtgtctgcat ctgtaggaga
cagagtcacc 60atcactgtc gggcgagtca ggttattagc agctggttag cctggtatca gcagaaacca 120gggaaagccc
ctagcctct gatctatgct gcatccagtt tgcaaagtgg ggtcccatca 180aggttcagcg gcagtgtatc tgggacagat
ttactctca ccatcagcag cctgcagcct 240gaagatttg caacttacta ttgtcaacag gtaacagtt tccattcac ttcggccct
300gggaccaaag tggattcaa a 32117107PRTTHomo sapiens 17Asp Ile Gln Met Thr Gln Ser Pro Ser Ser
Val Ser Ala Ser Val Gly1 5 10 15Asp Arg Val Thr Ile Thr Cys Arg Ala Ser Gln Gly Ser Ser Ser Trp 20
25 30Phe Ala Trp Tyr Gln Gln Lys Pro Gly Lys Ala Pro Lys Leu Leu Ile 35 40 45Tyr Ala Ala Ser Ser
Leu Gln Ser Gly Val Pro Ser Arg Phe Ser Gly 50 55 60Ser Gly Ser Gly Thr Asp Phe Thr Leu Thr Ile
Ser Ser Leu Gln Pro65 70 75 80Glu Asp Phe Ala Thr Tyr Tyr Cys Gln Gln Ala Asn Ser Phe Pro Phe 85
90 95Thr Phe Gly Pro Gly Thr Lys Val Asp Ile Lys 100 10518321DNAHomo sapiens 18gacatccaga
tgaccagtc tccatcttcc gtgtctgcat ctgtaggaga cagagtcacc 60atcactgtc gggcgagtca gggaagtagc
agctggttg cctggtatca gcagaaacca 120gggaaagccc caagctct gatctatgct gcatccagtt tgcaaagtgg
ggtcccatca 180aggttcagcg gcagtggatc tgggacagac ttactctca ccatcagcag cctgcagcct 240gaagatttg
caacttacta ttgtcaacag gtaacagtt tccattcac ttcggccct 300gggaccaaag tggatatcaa a
32119107PRTTHomo sapiens 19Asp Ser Gln Met Thr Gln Ser Pro Ser Ser Val Ser Ala Ser Val Gly1 5
10 15Asp Arg Val Thr Ile Thr Cys Arg Ala Ser Gln Gly Ile Ser Ser Trp 20 25 30Phe Ala Trp Tyr Gln Gln
Lys Pro Gly Gln Ala Pro Asn Leu Leu Ile 35 40 45Tyr Ala Ala Ser Ser Leu Gln Ser Gly Val Pro Ser Arg
Phe Ser Gly 50 55 60Ser Gly Ser Gly Thr Glu Phe Thr Leu Thr Ile Ser Ser Leu Gln Pro65 70 75 80Glu
Asp Phe Ala Thr Tyr Tyr Cys Gln Gln Ala Asn Ser Phe Pro Phe 85 90 95Thr Phe Gly Pro Gly Thr Lys
Val Asp Ile Lys 100 10520321DNAHomo sapiens 20gacagccaga tgaccagtc tccatcttcc gtgtctgct
ctgtaggaga cagagtcacc 60atcactgtc gggcgagtca gggattagc agctggttg cctggtatca gcagaaacca
120gggcaagccc ctaacctct gatctatgct gcatccagtt tgcaaagtgg ggtcccatca 180aggttcagcg gcagtggatc
tgggacagaa ttactctca ccatcagcag cctgcagcct 240gaagatttg caacttacta ttgtcaacag gtaacagtt
tccattcac ttcggccct 300gggaccaaag tggatatcaa a 32121107PRTTHomo sapiens 21Asp Ile Gln Met Thr
Gln Ser Pro Ser Ser Val Ser Ala Ser Val Gly1 5 10 15Asp Arg Val Thr Ile Thr Cys Arg Ala Gly Gln Val
Ile Ser Ser Trp 20 25 30Leu Ala Trp Tyr Gln Gln Lys Pro Gly Lys Ala Pro Lys Leu Leu Ile 35 40 45Tyr
Ala Ala Ser Ser Leu Gln Ser Gly Val Pro Ser Arg Phe Ser Gly 50 55 60Ser Gly Ser Gly Thr Asp Phe
Thr Leu Thr Ile Ser Ser Leu Gln Pro65 70 75 80Asp Asp Phe Ala Thr Tyr Tyr Cys Gln Gln Ala Thr Ser
Phe Pro Leu 85 90 95Thr Phe Gly Gly Gly Thr Lys Val Glu Ile Lys 100 10522321DNAHomo sapiens
22gacatccaga tgaccagtc tccatcttcc gtgtctgcat ctgtaggaga cagagtcacc 60atcactgtc gggcggtca
ggttattagc agctggttag cctggtatca gcagaaacca 120gggaaagccc ctaagctct gatctatgct gcatccagtt
tgcaaagtgg ggtcccatcg 180aggttcagcg gcagtggatc tgggacagat ttactctca ccatcagcag cctgcagcct
240gacgatttg caacttacta ttgtcaacag gctaccagtt tccctcac ttcggcgga 300gggaccaagg tggagatcaa a

32123107PRT Homo sapiens 23 Asp Ile Gln Met Thr Gln Ser Pro Ser Ser Val Ser Ala Ser Val Gly1 5 10
15 Asp Arg Val Thr Ile Thr Cys Arg Ala Ser Gln Gly Phe Ser Gly Trp 20 25 30 Leu Ala Trp Tyr Gln Gln
Lys Pro Gly Lys Ala Pro Lys Leu Leu Ile 35 40 45 Tyr Ala Ala Ser Ser Leu Gln Ser Gly Val Pro Ser Arg
Phe Ser Gly 50 55 60 Ser Gly Ser Gly Thr Asp Phe Thr Leu Thr Ile Ser Ser Leu Gln Pro 65 70 75 80 Glu
Asp Phe Ala Thr Tyr Tyr Cys Gln Gln Ala Asn Ser Phe Pro Phe 85 90 95 Thr Phe Gly Pro Gly Thr Lys
Val Asp Ile Lys 100 105 24321 DNA Homo sapiens 24 gacatccaga tgaccagtc tccatctcc gtgtctgcat
ctgtaggaga cagagtcacc 60 atcactgtc gggcgagtc ggggttagc ggtggtag cctggtatca gcagaaacca
120 gggaaagccc ctaagctct gatctatgct gcatccagtt tgcaaagtg ggtccatca 180 aggttcagtg gcagtgatc
tgggacagat ttactctca ccatcagcag cctgcagcct 240 gaagatttg caactacta ctgtcaacag gtaacagtt
tccattcac ttcggcct 300 gggaccaag tggatcaaa a 32125107PRT Homo sapiens 25 Asp Ile Gln Leu Thr
Gln Ser Pro Ser Ser Val Ser Ala Ser Val Gly1 5 10 15 Asp Arg Val Thr Ile Thr Cys Arg Ala Ser Gln Val
Ile Ser Ser Trp 20 25 30 Phe Ala Trp Tyr Gln Gln Lys Pro Gly Lys Ala Pro Asn Leu Leu Ile 35 40 45 Tyr
Ala Ala Ser Ser Leu Gln Ser Gly Val Pro Ser Arg Phe Ser Gly 50 55 60 Ser Gly Ser Gly Thr Asp Phe
Thr Leu Thr Ile Ser Ser Leu Gln Pro 65 70 75 80 Ala Asp Phe Ala Thr Tyr Phe Cys Gln Gln Ala Asn Ser
Phe Pro Phe 85 90 95 Thr Phe Gly Pro Gly Thr Lys Val Asp Val Lys 100 105 26321 DNA Homo sapiens
26 gacatccagt tgaccagtc tccatctcc gtgtctgcat ctgtaggaga cagagtcacc 60 atcactgtc gggcgagtc
ggttattagc agctggttg cctggtatca gcagaaacca 120 gggaaagccc ctaactct gatctatgct gcatccagtt
tgcaaagtg ggtccatca 180 aggttcagcg gcagtgatc tgggacagat ttactctca ccatcagcag cctgcagcct
240 gcagatttg caactacta ttgtcaacag gtaacagtt tccattcac ttcggcct 300 gggaccaag tggatcaaa a
32127107PRT Homo sapiens 27 Asp Ile Gln Met Thr Gln Ser Pro Ser Ser Val Ser Ala Ser Val Gly1 5 10
15 Asp Arg Val Thr Ile Thr Cys Arg Ala Ser Gln Gly Ser Ser Ser Trp 20 25 30 Phe Ala Trp Tyr Gln Gln
Lys Pro Gly Lys Ala Pro Lys Leu Leu Ile 35 40 45 Tyr Ala Ala Ser Ser Leu Gln Ser Gly Val Pro Ser Arg
Phe Ser Gly 50 55 60 Ser Gly Ser Gly Thr Asp Phe Thr Leu Thr Ile Ser Ser Leu Gln Pro 65 70 75 80 Glu
Asp Phe Ala Thr Tyr Tyr Cys Gln Gln Ala Asn Ser Phe Pro Phe 85 90 95 Thr Phe Gly Pro Gly Thr Lys
Val Asp Ile Lys 100 105 28321 DNA Homo sapiens 28 gacatccaga tgaccagtc tccatctcc gtgtctgcat
ctgtaggaga cagagtcacc 60 atcactgtc gggcgagtc gggtagtagc agctggttg cctggtatca acagaaacca
120 gggaaagccc caagctct gatctatgct gcatccagtt tgcaaagtg ggtccatca 180 aggttcagcg gcagtgatc
tgggacagat ttactctca ccatcagcag cctgcagcct 240 gaagatttg caactacta ttgtcaacag gtaacagtt
tccattcac ttcggcct 300 gggaccaag tggatcaaa a 32129107PRT Homo sapiens 29 Asp Ile Gln Met Thr
Gln Ser Pro Ser Ser Leu Ser Ala Ser Val Gly1 5 10 15 Asp Arg Val Thr Ile Thr Cys Arg Ala Ser Gln Gly
Ile Arg Asn Asp 20 25 30 Leu Gly Trp Tyr Gln Gln Lys Pro Gly Lys Ala Pro Lys Arg Leu Ile 35 40 45 Tyr
Ala Ala Ser Ser Leu Gln Ser Gly Val Pro Ser Arg Phe Ser Gly 50 55 60 Ser Gly Ser Gly Thr Glu Phe
Thr Leu Thr Ile Ser Ser Leu Gln Pro 65 70 75 80 Glu Asp Phe Ala Thr Tyr Tyr Cys Leu Gln His Asn Ser
Tyr Pro Pro 85 90 95 Thr Phe Gly Gln Gly Thr Lys Val Glu Ile Glu 100
105 30108PRT Artificial Consensus Sequence MISC_FEATURE(2)..(2) Xaa can be Ile or
Ser MISC_FEATURE(4)..(4) Xaa can be Met or Leu MISC_FEATURE(29)..(29) Xaa can be Gly or
Val MISC_FEATURE(30)..(30) Xaa can be Ser, Phe or Ile MISC_FEATURE(32)..(32) Xaa can be Ser or
Gly MISC_FEATURE(34)..(34) Xaa can be Phe or Leu MISC_FEATURE(43)..(43) Xaa can be Lys or
Gln MISC_FEATURE(46)..(46) Xaa can be Lys, Asn or Ser MISC_FEATURE(67)..(67) Xaa can be Gly or
Val MISC_FEATURE(71)..(71) Xaa can be Asp or Glu MISC_FEATURE(82)..(82) Xaa can be Glu or
Ala MISC_FEATURE(88)..(88) Xaa can be Tyr or Phe MISC_FEATURE(107)..(107) Xaa can be Ile, Val
or Phe 30 Asp Xaa Gln Xaa Thr Gln Ser Pro Ser Ser Val Ser Ala Ser Val Gly1 5 10 15 Asp Arg Val Thr
Ile Thr Cys Arg Ala Ser Gln Gly Xaa Xaa Ser Xaa 20 25 30 Trp Xaa Ala Trp Tyr Gln Gln Lys Pro Gly
Xaa Ala Pro Xaa Leu Leu 35 40 45 Ile Tyr Ala Ala Ser Ser Leu Gln Ser Gly Val Pro Ser Arg Phe Ser 50
55 60 Gly Ser Xaa Ser Gly Thr Xaa Phe Thr Leu Thr Ile Ser Ser Leu Gln 65 70 75 80 Pro Xaa Asp Phe
Ala Thr Tyr Xaa Cys Gln Gln Ala Asn Ser Phe Pro 85 90 95 Phe Thr Phe Gly Pro Gly Thr Lys Val Asp
Xaa Lys 100 105 31124PRT Homo sapiens 31 Gln Val Gln Leu Val Glu Ser Gly Gly Gly Val Val Gln Pro
Gly Arg1 5 10 15 Ser Leu Arg Leu Ser Cys Ala Ala Ser Gly Phe Thr Phe Ser Ser Tyr 20 25 30 Gly Met
His Trp Val Arg Gln Ala Pro Gly Lys Gly Leu Glu Trp Val 35 40 45 Ala Val Ile Trp Tyr Asp Gly Ser Asn
Glu Tyr Tyr Ala Asp Ser Val 50 55 60 Lys Gly Arg Phe Thr Ile Ser Arg Asp Asn Ser Lys Asn Thr Leu
Tyr 65 70 75 80 Leu Gln Met Asn Ser Leu Arg Ala Glu Asp Thr Ala Val Tyr Tyr Cys 85 90 95 Ala Arg Asp
Arg Gly Tyr Thr Ser Ser Trp Tyr Pro Asp Ala Phe Asp 100 105 110 Ile Trp Gly Gln Gly Thr Met Val Thr
Val Ser Ser 115 120 32372 DNA Homo sapiens 32 caggtgcagc tggtagagtc tggggaggc gtgtccagc
ctgggaggtc cctgagactc 60 tcctgtgcag cgctggatt cacctcagt agctatggca tgactgggt ccgcccagct
120 ccaggcaagg ggctggagtg ggtggcagtt atatggtatg atggaagtaa tgaatactat 180 gcagactccg tgaagggccg
attcaccatc tccagagaca attcaagaa cacgctgtat 240 ctgcaaatga acagcctgag agccgaggac acggctgtg

attactgtgc gagagatcgg 300gggtatacca gtagctggta cctgatgct tttgatatct gggccaagg gacaatggc
360accgtctctt ca 37233124PRT Homo sapiens 33Gln Val Gln Leu Val Glu Ser Gly Gly Gly Val Val Gln
Pro Gly Arg1 5 10 15Ser Leu Arg Leu Ser Cys Ala Ala Ser Gly Phe Thr Phe Ser Ser Tyr 20 25 30Gly
Met His Trp Val Arg Gln Ala Pro Gly Lys Gly Leu Glu Trp Val 35 40 45Ala Val Ile Trp Tyr Asp Gly Ser
Asn Lys Tyr Tyr Ala Asp Ser Val 50 55 60Lys Gly Arg Phe Thr Ile Ser Arg Asp Asn Ser Lys Asn Thr
Leu Tyr65 70 75 80Leu Gln Met Asn Ser Leu Arg Ala Glu Asp Thr Ala Val Tyr Tyr Cys 85 90 95Ala Arg
Asp Arg Gly Tyr Ser Ser Ser Trp Tyr Pro Asp Ala Phe Asp 100 105 110Ile Trp Gly Gln Gly Thr Met Val
Thr Val Ser Ser 115 12034121PRT Homo sapiens 34Gln Val Gln Leu Val Glu Ser Gly Gly Gly Val Val
Gln Pro Gly Arg1 5 10 15Ser Leu Arg Leu Ser Cys Ala Ala Ser Gly Phe Thr Phe Ser Ser Tyr 20 25
30Gly Met His Trp Val Arg Gln Ala Pro Gly Lys Gly Leu Glu Trp Val 35 40 45Ala Val Ile Ser Phe Asp
Gly Ser Leu Lys Tyr Tyr Ala Asp Ser Val 50 55 60Lys Gly Arg Phe Thr Ile Ser Arg Asp Asn Ser Lys
Asn Thr Leu Tyr65 70 75 80Leu Gln Met Asn Ser Leu Arg Ala Glu Asp Thr Ala Val Tyr Tyr Cys 85 90
95Ala Arg Glu Arg Thr Thr Leu Ser Gly Ser Tyr Phe Asp Tyr Trp Gly 100 105 110Gln Gly Thr Leu Val
Thr Val Ser Ser 115 12035363DNA Homo sapiens 35caggtgcagc tgggtggagtc tgggggaggc gtggtccagc
ctgggaggc cctgagactc 60tcctgtgcag cctctggatt cacctcagt agctatggca tgcactgggt ccgccaggct
120ccaggcaagg ggctggagtg ggtggcagtt atatcattg atggaagtct taaatactat 180gcagactccg tgaagggccg
attcaccatc tccagagaca attccaagaa caccctgtat 240ctgcaaatga acagcctgag agctgaggac acggctgtgt
attactgtgc gagagaacgg 300actactttaa gtgggagcta ctttgactac tggggccagg gaaccctggt caccgtctcc 360tca
36336121PRT Homo sapiens 36Gln Val Gln Leu Val Glu Ser Gly Gly Gly Val Val Gln Pro Gly Arg1 5 10
15Ser Leu Arg Leu Ser Cys Ala Ala Ser Gly Phe Thr Phe Ser Ser Tyr 20 25 30Ala Met His Trp Val Arg
Gln Ala Pro Gly Lys Gly Leu Glu Trp Leu 35 40 45Ser Val Ile Ser His Asp Gly Ser Ile Lys Tyr Tyr Ala
Asp Ser Val 50 55 60Lys Gly Arg Phe Thr Ile Ser Arg Asp Asn Ser Lys Asn Thr Leu Tyr65 70 75 80Leu
Gln Met Asn Ser Leu Arg Ala Glu Asp Thr Ala Val Tyr Tyr Cys 85 90 95Ala Arg Glu Arg Thr Thr Leu
Ser Gly Ser Tyr Phe Asp Tyr Trp Gly 100 105 110Gln Gly Thr Leu Val Thr Val Ser Ser 115
12037363DNA Homo sapiens 37caggtgcagc tgggtggagtc tgggggaggc gtggtccagc ctgggaggc cctgagactc
60tcctgtgcag cctctggatt cacctcagt agctatgcca tgcactgggt ccgccaggct 120ccaggcaagg ggctggagtg
gtgtcagtt atatcacatg atggaagtat taaatactat 180gcagactccg tgaagggccg attcaccatc tccagagaca
attccaagaa cacgctgtat 240ctgcaaatga acagcctgag agctgaggac acggctgtgt attactgtgc gagagaacgg
300actactctaa gtgggagcta ctttgactac tggggccagg gaaccctggt caccgtctcc 360tca 36338125PRT Homo
sapiens 38Glu Val Gln Leu Val Glu Ser Gly Gly Gly Leu Val Gln Pro Gly Gly1 5 10 15Ser Leu Arg Leu
Ser Cys Ala Ala Ser Gly Phe Thr Phe Ser Ser Tyr 20 25 30Ser Met Asn Trp Val Arg Gln Ala Pro Gly
Lys Gly Leu Glu Trp Val 35 40 45Ser Tyr Ile Ser Ser Arg Ser Ser Thr Ile Tyr Ile Ala Asp Ser Val 50 55
60Lys Gly Arg Phe Thr Ile Ser Arg Asp Asn Ala Lys Asn Ser Leu Tyr65 70 75 80Leu Gln Met Asn Ser
Leu Arg Asp Glu Asp Thr Ala Val Tyr Tyr Cys 85 90 95Ala Arg Arg Ile Ala Ala Ala Gly Gly Phe His Tyr
Tyr Tyr Ala Leu 100 105 110Asp Val Trp Gly Gln Gly Thr Thr Val Thr Val Ser Ser 115 120
12539375DNA Homo sapiens 39gaggtgcagc tgggtggagtc tgggggaggc ctggtacagc ctggggggtc cctgagactc
60tcctgtgcag cctctggatt cacctcagt agctatagta tgaactgggt ccgccaggct 120ccagggaagg ggctggagtg
ggtttcgtac attagtagta ggagtagtac catatacatc 180gcagactctg tgaagggccg attcaccatc tccagagaca
atgccaagaa ctactgtat 240ctgcaaatga acagcctgag agacgaagac acggctgtgt attactgtgc gagacggata
300gcagcagctg gtgggtcca ctactactac gctttggagc tctggggcca agggaccagc 360gtcaccgtct cctca
37540125PRT Homo sapiens 40Glu Val Gln Leu Val Glu Ser Gly Gly Gly Leu Val Gln Pro Gly Gly1 5
10 15Ser Leu Arg Leu Ser Cys Ala Ala Ser Gly Phe Thr Phe Ser Thr Tyr 20 25 30Ser Met Asn Trp Val
Arg Gln Ala Pro Gly Lys Gly Leu Glu Trp Val 35 40 45Ser Tyr Ile Ser Ser Ser Ser Ser Thr Arg Tyr His
Ala Asp Ser Val 50 55 60Lys Gly Arg Phe Thr Ile Ser Arg Asp Asn Ala Lys Asn Ser Leu Tyr65 70 75
80Leu Gln Met Asn Ser Leu Arg Asp Glu Asp Thr Ala Val Tyr Tyr Cys 85 90 95Ala Arg Arg Ile Ala Ala
Ala Gly Pro Trp Gly Tyr Tyr Tyr Ala Met 100 105 110Asp Val Trp Gly Gln Gly Thr Thr Val Thr Val Ser
Ser 115 120 12541375DNA Homo sapiens 41gaggtgcagc tgggtggagtc tgggggaggc ttgtacaac
ctggggggtc cctgagactc 60tcctgtgcag cctctggatt cacctcagt acctatagca tgaactgggt ccgccaggct
120ccagggaagg ggctggagtg ggtttcatac attagtagca gtagtagtac cagataccac 180gcagactctg tgaagggccg
attcaccatc tccagagaca atgccaagaa ctactgtat 240ctgcaaatga acagcctgag agacgaggac acggctgtgt
attactgtgc gagacgtata 300gcagcagctg gtccgtgggg ctactactac gctatggagc tctggggcca agggaccagc
360gtcaccgtct cctca 37542125PRT Homo sapiens 42Glu Val Gln Leu Val Glu Ser Gly Gly Gly Leu Val
Gln Pro Gly Gly1 5 10 15Ser Leu Arg Leu Ser Cys Val Val Ser Gly Phe Thr Phe Ser Ser Phe 20 25
30Ser Met Asn Trp Val Arg Gln Ala Pro Gly Lys Gly Leu Glu Trp Val 35 40 45Ser Tyr Ile Ser Ser Arg
Ser Ser Thr Ile Tyr Tyr Ala Asp Ser Val 50 55 60Lys Gly Arg Phe Thr Ile Ser Arg Asp Asn Ala Lys Asn
Ser Leu Tyr65 70 75 80Leu Gln Met Asn Ser Leu Arg Asp Glu Asp Thr Ala Val Tyr Tyr Cys 85 90

95Ala Arg Arg Ile Ala Ala Ala Gly Pro Trp Gly Tyr Tyr Tyr Ala Met 100 105 110Asp Val Trp Gly Gln Gly Thr Thr Val Thr Val Ser Ser 115 120 12543375DNAHomo sapiens 43gaggtgcagc tgggtggagtc tgggggaggc tgggtacagc ctggggggc cctgagactc 60tctctgttag tctctggatt caccttcagt agtttagca tgaactggct cgcaggct 120ccaggaagg ggctggagtg ggttcatac atagtagtc gtagtagtac catatactac 180gcagactctg tgaagggccg attcaccatc tccagagaca atgccaagaa ctactgtat 240ctgcaaatga acagcctgag agacgaggac acggctgtgt attattgtgc gagacgtata 300gcagcagctg gtccgtggg ctactactac gctatggagc tctggggcca agggaccacg 360gtcaccgtct cctca 37544118PRTTHomo sapiens 44Gln Val Gln Leu Gln Glu Ser Gly Pro Gly Leu Val Lys Pro Ser Glu1 5 10 15Thr Leu Ser Leu Thr Cys Thr Val Ser Gly Gly Ser Ile Ser Thr Tyr 20 25 30Tyr Trp Ser Trp Ile Arg Gln Pro Ala Gly Lys Gly Leu Glu Trp Ile 35 40 45Gly Leu Ile Tyr Thr Ser Gly Ser Thr Asn Tyr Asn Pro Ser Leu Lys 50 55 60Ser Arg Val Thr Met Ser Leu Asp Thr Ser Lys Asn Gln Phe Ser Leu65 70 75 80Arg Leu Thr Ser Val Thr Ala Ala Asp Thr Ala Val Tyr Tyr Cys Ala 85 90 95Arg Asp Arg Gly Tyr Tyr Tyr Gly Val Asp Val Trp Gly Gln Gly Thr 100 105 110Thr Val Thr Val Ser Ser 11545354DNAHomo sapiens 45caggtgcagc tgcaggagtc gggcccagga ctggtgaagc ctccggagac cctgtccctc 60acctgcactg tctctgggtg ctccatcagt acttactact ggagctggat cggcgagccc 120gccgggaagg gactggagtg gattgggctt atctatacca gtgggagcac caactacaac 180ccctccctca agagtgcagt caccatgtca ttagacacgt ccaagaacca gtctccctg 240aggctgacct ctgtgaccgc cgcggacacg gccgttatt actgtgagc agatcgtggg 300tactactacg gtgtggacgt ctggggccag gggaccacg tcaccgtctc ctca 35446120PRTTHomo sapiens 46Gln Val Gln Leu Gln Glu Ser Gly Pro Gly Leu Val Lys Pro Ser Gln1 5 10 15Thr Leu Ser Leu Thr Cys Thr Val Ser Gly Gly Ser Ile Ser Ser Gly 20 25 30Gly Tyr Tyr Trp Ser Trp Ile Arg Gln His Pro Gly Lys Gly Leu Glu 35 40 45Trp Ile Gly His Ile His Tyr Ser Gly Asn Thr Tyr Tyr Asn Pro Ser 50 55 60Leu Lys Ser Arg Val Thr Ile Ser Val Asp Thr Ser Lys Asn Gln Phe65 70 75 80Ser Leu Lys Leu Ser Ser Val Thr Ala Ala Asp Thr Ala Val Tyr Tyr 85 90 95Cys Ala Lys Asn Arg Gly Phe Tyr Tyr Gly Met Asp Val Trp Gly Gln 100 105 110Gly Thr Thr Val Thr Val Ser Ser 115 12047360DNAHomo sapiens 47caggtgcagc tgcaggagtc gggcccagga ctggtgaagc cttcacagac cctgtccctc 60acctgcactg tctctgggtg ctccatcagc agtgggtggt actactggag ctggatccgc 120cagcaccag ggaagggcct ggagtggatt gggcacatcc attacagtg gaacacctac 180tacaaccctg cctcaagag tgcagttacc atatcagtag acacgtctaa gaatcagttc 240tccctgaaac tgagctctgt gactgccgcg gacacggccg tgtattactg tgcgaaaaat 300cgcgggttct actacggtat ggacgtctgg ggccaagga ccacggtcac cgtctcctca 36048120PRTTHomo sapiens 48Gln Val Gln Leu Gln Glu Ser Gly Pro Gly Leu Val Lys Pro Ser Gln1 5 10 15Thr Leu Ser Leu Thr Cys Thr Val Ser Gly Gly Ser Ile Asn Ser Gly 20 25 30Gly Tyr Tyr Trp Ser Trp Ile Arg Gln His Pro Gly Lys Gly Leu Glu 35 40 45Trp Ile Gly Tyr Ile Tyr Tyr Ser Gly Ser Ser Tyr Tyr Asn Pro Ser 50 55 60Leu Lys Ser Arg Val Thr Ile Ser Val Asp Thr Ser Gln Asn Gln Phe65 70 75 80Ser Leu Lys Leu Ser Ser Val Thr Ala Ala Asp Thr Ala Val Tyr Tyr 85 90 95Cys Ala Arg Asp Arg Gly His Tyr Tyr Gly Met Asp Val Trp Gly Gln 100 105 110Gly Thr Thr Val Thr Val Ser Ser 115 12049360DNAHomo sapiens 49caggtgcagc tgcaggagtc gggcccagga ctggtgaagc cttcacagac cctgtccctc 60acctgcactg tctctgggtg ctccatcaac agtgggtggt actactggag ctggatccgc 120cagcaccag ggaagggcct ggagtggatt ggtacatct attacagtg gagctcctac 180tacaaccctg cctcaagag tgcagttacc atatcagtag acacgtctaa gaaccagttc 240tccctgaagc tgagctctgt gactgccgcg gacacggccg tgtattactg tgcgagagat 300cgggggact actacggtat ggacgtctgg ggccaagga ccacggtcac cgtctcctca 36050120PRTTHomo sapiens 50Gln Val Gln Leu Gln Glu Ser Gly Pro Gly Leu Val Lys Pro Ser Gln1 5 10 15Thr Leu Ser Leu Thr Cys Thr Val Ser Gly Gly Ser Ile Ser Ser Gly 20 25 30Gly Tyr Tyr Trp Ser Trp Ile Arg Gln His Pro Gly Lys Gly Leu Glu 35 40 45Trp Ile Gly Tyr Ile Tyr Tyr Ser Gly Ser Thr Tyr Tyr Asn Pro Ser 50 55 60Leu Lys Ser Arg Val Thr Ile Ser Val Asp Thr Ser Lys Asn Gln Phe65 70 75 80Ser Leu Lys Leu Ser Ser Val Thr Ala Ala Asp Thr Ala Val Tyr Tyr 85 90 95Cys Ala Arg Asp Arg Gly His Tyr Tyr Gly Met Asp Val Trp Gly Gln 100 105 110Gly Thr Thr Val Thr Val Ser Ser 115 12051360DNAHomo sapiens 51caggtgcagc tgcaggagtc gggcccagga ctggtgaagc cttcacagac cctgtccctc 60acctgcactg tctctgggtg ctccatcagt agtgggtggt actactggag ctggatccgc 120cagcaccag ggaagggcct ggagtggatt ggtacatt attacagtg gagcacctac 180tacaaccctg cctcaagag tgcagttacc atatcagtag acacgtctaa gaaccagttc 240tccctgaagc tgagctctgt gactgccgcg gacacggccg tgtattactg tgcgagagat 300cgggggact actatggaat ggacgtctgg ggccaagga ccacggtcac cgtctcctca 36052118PRTTHomo sapiens 52Gln Val Gln Leu Gln Glu Ser Gly Pro Arg Leu Val Lys Pro Ser Glu1 5 10 15Thr Leu Ser Leu Thr Cys Thr Val Ser Gly Asp Ser Ile Ser Ser Tyr 20 25 30Phe Trp Ser Trp Ile Arg Gln Pro Pro Gly Lys Gly Leu Glu Trp Leu 35 40 45Gly Tyr Ile Tyr Tyr Ser Gly Ser Thr Asn Tyr Asn Pro Ser Leu Lys 50 55 60Ser Arg Val Thr Ile Ser Ile Asp Thr Ser Lys Asn Gln Phe65 70 75 80Lys Leu Ser Ser Val Thr Ala Ala Asp Thr Ala Val Tyr Tyr Cys Thr 85 90 95Arg Asp Arg Gly Ser Tyr Tyr Gly Ser Asp Tyr Trp Gly Gln Gly Thr 100 105 110Leu Val Thr Val Ser Ser 11553354DNAHomo sapiens 53caggtgcagc tgcaggagtc gggccaaga ctggtgaagc ctccggagac cctgtccctc 60acctgcactg tctctgggtg ctccatcagt agttactct ggagtggat cggcgagccc

120ccaggggaagg gactggagtg gcttggtat atctattaca gtgggagcac caactacaac 180ccctcctca agagtcgagt
cacatatca atagacacgt ccaagaacca gttctcctg 240aagctgagct ctgtgaccgc tgcggacacg gccgtgtatt
actgtacgag agatcggggg 300agctactacg gatctgacta ctggggccag ggaaccctgg tcaccgtctc ctca
35454120PRT Homo sapiens 54Gln Val Gln Leu Gln Glu Ser Gly Pro Gly Leu Val Lys Pro Ser Gln1 5
10 15Thr Leu Ser Leu Thr Cys Thr Val Ser Gly Gly Ser Ile Ser Ser Gly 20 25 30Gly Tyr Tyr Trp Thr Trp
Ile Arg Gln His Pro Gly Lys Gly Leu Glu 35 40 45Trp Ile Gly Tyr Ile Tyr Tyr Ser Gly Asn Thr Tyr Tyr Asn
Pro Ser 50 55 60Leu Lys Ser Arg Ile Thr Ile Ser Val Asp Thr Ser Lys Asn Gln Phe65 70 75 80Ser Leu
Ser Leu Ser Ser Val Thr Ala Ala Asp Thr Ala Val Tyr Tyr 85 90 95Cys Ala Arg Asn Arg Gly Tyr Tyr Tyr
Gly Met Asp Val Trp Gly Gln 100 105 110Gly Thr Thr Val Thr Val Ser Ser 115 12055360DNA Homo
sapiens 55caggtgcagc tgcaggagtc gggcccagga ctggtgaagc cttcacagac cctgtccctc 60acctgcactg
tctctggtgg ctccatcagc agtgggtggt actactggac ctggatccgc 120cagcaccag ggaagggcct ggagtggatt
gggtacatct attacagtgg gaacacctac 180tacaaccctg cctcaagag tgaattacc atatcagtgg acacgtctaa
gaaccagttc 240tccctgagcc tgagctctgt gactgccgcg gacacggccg tgtattactg tgcgagaaat 300cgcggttact
actacggtat ggacgtctgg ggccaagga ccacgggtcac cgtctcctca 36056120PRT Homo sapiens 56Gln Val Gln
Leu Gln Glu Ser Gly Pro Gly Leu Val Lys Pro Ser Gln1 5 10 15Thr Leu Ser Leu Thr Cys Thr Val Ser
Gly Gly Ser Ile Ser Ser Gly 20 25 30Gly Tyr Tyr Trp Ser Trp Ile Arg Gln His Pro Gly Lys Gly Leu Glu 35
40 45Trp Ile Gly Tyr Ile Tyr Tyr Ser Gly Ser Thr Tyr Tyr Asn Pro Ser 50 55 60Leu Lys Ser Arg Val Thr
Met Ser Val Asp Thr Ser Lys Asn Gln Phe65 70 75 80Ser Leu Lys Leu Ser Ser Val Thr Ala Ala Asp Thr
Ala Val Tyr Tyr 85 90 95Cys Ala Lys Asn Arg Gly Phe Tyr Tyr Gly Met Asp Val Trp Gly Gln 100 105
110Gly Thr Thr Val Thr Val Ser Ser 115 12057360DNA Homo sapiens 57caggtgcagc tgcaggagtc
gggcccagga ctggtgaagc cttcacagac cctgtccctc 60acctgcactg tctctggtgg ctccatcagc agtgggtggt
actactggag ctggatccgc 120cagcaccag ggaagggcct ggagtggatt gggtacatct attacagtgg gagcacctac
180tacaaccctg cctcaagag tcgagttacc atgtcagtag acacgtctaa gaaccagttc 240tccctgaaac tgagctctgt
gactgccgcg gacacggccg tgtattactg tgcgaaaaat 300cgcggttct actacggtat ggacgtctgg ggccaagga
ccacgggtcac cgtctcctca 36058120PRT Homo sapiens 58Gln Val Gln Leu Gln Glu Ser Gly Pro Gly Leu
Val Lys Pro Ser Gln1 5 10 15Thr Leu Ser Leu Thr Cys Thr Val Ser Gly Gly Ser Ile Asn Ser Gly 20 25
30Gly Tyr Tyr Trp Ser Trp Ile Arg Gln His Pro Gly Lys Gly Leu Glu 35 40 45Trp Ile Gly Tyr Ile Tyr Tyr
Ser Gly Ser Ser Tyr Tyr Asn Pro Ser 50 55 60Leu Lys Ser Arg Val Thr Ile Ser Val Asp Thr Ser Lys Asn
Gln Phe65 70 75 80Ser Leu Lys Leu Ser Ser Val Thr Ala Ala Asp Thr Ala Val Tyr Tyr 85 90 95Cys Ala
Arg Asp Arg Gly His Tyr Tyr Gly Met Asp Val Trp Gly Gln 100 105 110Gly Thr Thr Val Thr Val Ser Ser
115 12059360DNA Homo sapiens 59caggtgcagc tgcaggagtc gggcccagga ctggtgaagc cttcacagac
cctgtccctc 60acctgcactg tctctggtgg ctccatcaat agtgggtggt actactggag ctggatccgc 120cagcaccag
ggaagggcct ggagtggatt gggtacatct attacagtgg gagcagctac 180tacaaccctg cctcaagag tcgagttacc
atatcagttg acacgtctaa gaaccagttc 240tccctgaagc tgagttctgt gactgccgcg gacacggccg tgtattactg
tgcgagagat 300cgggggcact actacggtat ggacgtctgg ggccaagga ccacgggtcac cgtctcctca
36060123PRT Homo sapiens 60Gln Val Gln Leu Val Glu Ser Gly Gly Gly Val Val Gln Pro Gly Arg1 5 10
15Ser Leu Arg Leu Ser Cys Ala Ala Ser Gly Phe Thr Phe Ser Ser Tyr 20 25 30Gly Met His Trp Val Arg
Gln Ala Pro Gly Lys Gly Leu Glu Trp Val 35 40 45Ala Leu Ile Trp Tyr Asp Gly Ser Asn Lys Tyr Tyr Ala
Asp Ser Val 50 55 60Lys Gly Arg Phe Thr Ile Ser Arg Asp Asn Ser Lys Asn Thr Leu Tyr65 70 75 80Leu
Gln Met Asn Ser Leu Arg Ala Glu Asp Thr Ala Val Tyr Tyr Cys 85 90 95Ala Arg Glu Asn Thr Val Thr Ile
Tyr Tyr Asn Tyr Gly Met Asp Val 100 105 110Trp Gly Gln Gly Thr Thr Val Thr Val Ser Ser 115
12061116PRT ArtificialConsensus Sequence MISC_FEATURE(3)..(3)Xaa can be Val or
Glu MISC_FEATURE(25)..(25)Xaa can be Asn or Ser MISC_FEATURE(38)..(38)Xaa can be Gln or
Leu MISC_FEATURE(55)..(55)Xaa can be Ile or Thr MISC_FEATURE(61)..(61)Xaa can be Asp or
Glu MISC_FEATURE(77)..(77)Xaa can be Tyr or Ser MISC_FEATURE(101)..(101)Xaa can be Ser or
Asn 61Gln Pro Xaa Leu Thr Gln Pro Pro Ser Ala Ser Ala Ser Leu Gly Ala1 5 10 15Ser Val Thr Leu Thr
Cys Thr Leu Xaa Ser Gly Tyr Ser Asp Tyr Lys 20 25 30Val Asp Trp Tyr Gln Xaa Arg Pro Gly Lys Gly
Pro Arg Phe Val Met 35 40 45Arg Val Gly Thr Gly Gly Xaa Val Gly Ser Lys Gly Xaa Gly Ile Pro 50 55
60Asp Arg Phe Ser Val Leu Gly Ser Gly Leu Asn Arg Xaa Leu Thr Ile65 70 75 80Lys Asn Ile Gln Glu
Glu Asp Glu Ser Asp Tyr His Cys Gly Ala Asp 85 90 95His Gly Ser Gly Xaa Asn Phe Val Tyr Val Phe
Gly Thr Gly Thr Lys 100 105 110Val Thr Val Leu 1156214PRT Homo sapiens 62Thr Gly Ser Ser Ser
Asn Thr Gly Ala Gly Tyr Asp Val His1 5 10637PRT Homo sapiens 63Gly Ser Gly Asn Arg Pro Ser1
56411PRT Homo sapiens 64Gln Ser Tyr Asp Ser Ser Leu Ser Gly Trp Val1 5 106514PRT Homo
sapiens 65Thr Gly Ser Ser Ser Asn Ile Gly Ala Gly Tyr Asp Val His1 5 10667PRT Homo sapiens 66Gly
Ser Asn Asn Arg Pro Ser1 5679PRT Homo sapiens 67Met Ile Trp His Ser Ser Ala Ser Val1
56814PRT Homo sapiens 68Thr Leu Arg Ser Gly Ile Asn Val Gly Thr Tyr Arg Ile Tyr1 5

106911PRT Homo sapiens 69 Tyr Lys Ser Asp Ser Asp Lys Gln Gln Gly Ser1 5 107013PRT Homo sapiens 70 Gly Ala Asp His Gly Ser Gly Ser Asn Phe Val Tyr Val1 5 107111PRT Homo sapiens 71 Thr Leu Asn Ser Gly Tyr Ser Asp Tyr Lys Val1 5 107212PRT Homo sapiens 72 Val Gly Thr Gly Gly Ile Val Gly Ser Lys Gly Asp1 5 107313PRT Homo sapiens 73 Gly Ala Asp His Gly Ser Gly Asn Asn Phe Val Tyr Val1 5 107411PRT Homo sapiens 74 Thr Leu Ser Ser Gly Tyr Ser Asp Tyr Lys Val1 5 107512PRT Homo sapiens 75 Val Gly Thr Gly Gly Ile Val Gly Ser Lys Gly Glu1 5 10769PRT Homo sapiens 76 Gln Gln Ala Asn Ser Phe Pro Phe Thr1 57711PRT Homo sapiens 77 Arg Ala Ser Gln Gly Phe Ser Gly Trp Leu Ala1 5 107812PRT Homo sapiens 78 Val Gly Thr Gly Gly Thr Val Gly Ser Lys Gly Glu1 5 10799PRT Homo sapiens 79 Gln Gln Ala Thr Ser Phe Pro Leu Thr1 58011PRT Homo sapiens 80 Arg Ala Ser Gln Val Ile Ser Ser Trp Leu Ala1 5 10817PRT Homo sapiens 81 Ala Ala Ser Ser Leu Gln Ser1 5829PRT Homo sapiens 82 Gln Gln Ala Asp Ser Phe Pro Pro Thr1 58311PRT Homo sapiens 83 Arg Ala Ser Gln Val Ile Ser Ser Trp Phe Ala1 5 10849PRT Homo sapiens 84 Leu Gln His Asn Ser Tyr Pro Pro Thr1 58511PRT Homo sapiens 85 Arg Ala Ser Gln Gly Ser Ser Ser Trp Phe Ala1 5 108611PRT Homo sapiens 86 Arg Ala Ser Gln Gly Ile Ser Ser Trp Phe Ala1 5 108711PRT Homo sapiens 87 Arg Ala Gly Gln Val Ile Ser Ser Trp Leu Ala1 5 108811PRT Homo sapiens 88 Arg Ala Ser Gln Gly Ile Ala Gly Trp Leu Ala1 5 108911PRT Homo sapiens 89 Arg Ala Ser Gln Gly Ile Arg Asn Asp Leu Gly1 5 109017PRT Homo sapiens 90 Leu Ile Trp Tyr Asp Gly Ser Asn Lys Tyr Tyr Ala Asp Ser Val Lys1 5 10 15Gly915PRT Homo sapiens 91 Ser Tyr Gly Met His1 59217PRT Homo sapiens 92 Val Ile Trp Tyr Asp Gly Ser Asn Glu Tyr Tyr Ala Asp Ser Val Lys1 5 10 15Gly9315PRT Homo sapiens 93 Asp Arg Gly Tyr Thr Ser Ser Trp Tyr Pro Asp Ala Phe Asp Ile1 5 10 15945PRT Homo sapiens 94 Ser Tyr Ala Met His1 59517PRT Homo sapiens 95 Val Ile Trp Tyr Asp Gly Ser Asn Lys Tyr Tyr Ala Asp Ser Val Lys1 5 10 15Gly9615PRT Homo sapiens 96 Asp Arg Gly Tyr Ser Ser Ser Trp Tyr Pro Asp Ala Phe Asp Ile1 5 10 15975PRT Homo sapiens 97 Thr Tyr Ser Met Asn1 59817PRT Homo sapiens 98 Val Ile Ser Phe Asp Gly Ser Leu Lys Tyr Tyr Ala Asp Ser Val Lys1 5 10 15Gly9912PRT Homo sapiens 99 Glu Arg Thr Thr Leu Ser Gly Ser Tyr Phe Asp Tyr1 5 101005PRT Homo sapiens 100 Ser Tyr Ser Met Asn1 510117PRT Homo sapiens 101 Val Ile Ser His Asp Gly Ser Ile Lys Tyr Tyr Ala Asp Ser Val Lys1 5 10 15Gly10216PRT Homo sapiens 102 Arg Ile Ala Ala Ala Gly Gly Phe His Tyr Tyr Tyr Ala Leu Asp Val1 5 10 151035PRT Homo sapiens 103 Ser Phe Ser Met Asn1 510417PRT Homo sapiens 104 Tyr Ile Ser Ser Arg Ser Ser Thr Ile Tyr Ile Ala Asp Ser Val Lys1 5 10 15Gly10516PRT Homo sapiens 105 Arg Ile Ala Ala Ala Gly Pro Trp Gly Tyr Tyr Tyr Ala Met Asp Val1 5 10 151067PRT Homo sapiens 106 Ser Gly Gly Tyr Tyr Trp Thr1 510717PRT Homo sapiens 107 Tyr Ile Ser Ser Ser Ser Thr Arg Tyr His Ala Asp Ser Val Lys1 5 10 15Gly10810PRT Homo sapiens 108 Asn Arg Gly Tyr Tyr Tyr Gly Met Asp Val1 5 101097PRT Homo sapiens 109 Ser Gly Gly Tyr Tyr Trp Ser1 511017PRT Homo sapiens 110 Tyr Ile Ser Ser Arg Ser Ser Thr Ile Tyr Tyr Ala Asp Ser Val Lys1 5 10 15Gly11110PRT Homo sapiens 111 Asn Arg Gly Phe Tyr Tyr Gly Met Asp Val1 5 101125PRT Homo sapiens 112 Ser Tyr Phe Trp Ser1 511316PRT Homo sapiens 113 Tyr Ile Tyr Tyr Ser Gly Asn Thr Tyr Tyr Asn Pro Ser Leu Lys Ser1 5 10 1511410PRT Homo sapiens 114 Asp Arg Gly His Tyr Tyr Gly Met Asp Val1 5 101155PRT Homo sapiens 115 Thr Tyr Tyr Trp Ser1 511616PRT Homo sapiens 116 His Ile His Tyr Ser Gly Asn Thr Tyr Tyr Asn Pro Ser Leu Lys Ser1 5 10 1511710PRT Homo sapiens 117 Asp Arg Gly Ser Tyr Tyr Gly Ser Asp Tyr1 5 1011816PRT Homo sapiens 118 Tyr Ile Tyr Tyr Ser Gly Ser Thr Tyr Tyr Asn Pro Ser Leu Lys Ser1 5 10 1511910PRT Homo sapiens 119 Asp Arg Gly Tyr Tyr Tyr Gly Val Asp Val1 5 1012016PRT Homo sapiens 120 Tyr Ile Tyr Tyr Ser Gly Ser Ser Tyr Tyr Asn Pro Ser Leu Lys Ser1 5 10 1512116PRT Homo sapiens 121 Tyr Ile Tyr Tyr Ser Gly Ser Thr Asn Tyr Asn Pro Ser Leu Lys Ser1 5 10 1512216PRT Homo sapiens 122 Leu Ile Tyr Thr Ser Gly Ser Thr Asn Tyr Asn Pro Ser Leu Lys Ser1 5 10 1512311PRT ArtificialConsensus sequence MISC_FEATURE(5)..(5) Xaa can be Gly or Val MISC_FEATURE(6)..(6) Xaa can be Ile, Phe or Ser MISC_FEATURE(8)..(8) Xaa can be Ser or Gly MISC_FEATURE(10)..(10) Xaa can be Phe or Leu. 123 Arg Ala Ser Gln Xaa Xaa Ser Xaa Trp Xaa Ala1 5 1012412PRT ArtificialConsensus sequence MISC_FEATURE(3)..(3) Xaa can be Asn or Ser 124 Thr Leu Xaa Ser Gly Tyr Ser Asp Tyr Lys Val Asp1 5 1012514PRT ArtificialConsensus sequence MISC_FEATURE(7)..(7) Xaa can be Ile or Thr 125 Thr Gly Ser Ser Ser Asn Xaa Gly Ala Gly Tyr Asp Val His1 5 1012612PRT ArtificialConsensus sequence MISC_FEATURE(6)..(6) Xaa can be Ile or Thr MISC_FEATURE(12)..(12) Xaa can be Asp or Glu 126 Val Gly Thr Gly Gly Xaa Val Gly Ser Lys Gly Xaa1 5 101277PRT ArtificialConsensus sequence MISC_FEATURE(3)..(3) Xaa can be Asn or Gly 127 Gly Ser Xaa Asn Arg Pro Ser1 512813PRT ArtificialConsensus sequence MISC_FEATURE(8)..(8) Xaa can be Ser or Asn 128 Gly Ala Asp His Gly Ser Gly Xaa Asn Phe Val Tyr Val1 5 101297PRT ArtificialConsensus sequence MISC_FEATURE(7)..(7) Xaa can be Ser or Thr 129 Ser Gly

Gly Tyr Tyr Trp Xaa1 51305PRTArtificialConsensus sequenceMISC_FEATURE(3)..(3)Xaa can be Gly or Ala 130Ser Tyr Xaa Met His1 51315PRTArtificialConsensus sequenceMISC_FEATURE(1)..(1)Xaa can be Ser or ThrMISC_FEATURE(1)..(1)Xaa can be Ser or ThrMISC_FEATURE(2)..(2)Xaa can be Tyr or Phe 131Xaa Xaa Ser Met Asn1 513216PRTArtificialConsensus sequenceMISC_FEATURE(1)..(1)Xaa can be Tyr or HisMISC_FEATURE(3)..(3)Xaa can be Thr or HisMISC_FEATURE(7)..(7)Xaa can be Ser or AsnMISC_FEATURE(8)..(8)Xaa can be Thr or Ser 132Xaa Ile Xaa Tyr Ser Gly Xaa Xaa Tyr Tyr Asn Pro Ser Leu Lys Ser1 5 10 1513317PRTArtificialConsensus sequenceMISC_FEATURE(4)..(4)Xaa can be Phe or HisMISC_FEATURE(8)..(8)Xaa can be Leu or Thr 133Val Ile Ser Xaa Asp Gly Ser Xaa Lys Tyr Tyr Ala Asp Ser Val Lys1 5 10 15Gly13417PRTArtificialConsensus sequenceMISC_FEATURE(5)..(5)Xaa can be Arg or SerMISC_FEATURE(9)..(9)Xaa can be Ile or ArgMISC_FEATURE(11)..(11)Xaa can be Ile, His or Try 134Tyr Ile Ser Ser Xaa Ser Ser Thr Xaa Tyr Xaa Ala Asp Ser Val Lys1 5 10 15Gly13517PRTArtificialConsensus sequenceMISC_FEATURE(9)..(9)Xaa can be Lys or Glu 135Val Ile Trp Tyr Asp Gly Ser Asn Xaa Tyr Tyr Ala Asp Ser Val Lys1 5 10 15Gly13610PRTArtificialConsensus sequenceMISC_FEATURE(1)..(1)Xaa can be Asn or AspMISC_FEATURE(4)..(4)Xaa can be His, Tyr or Phe 136Xaa Arg Gly Xaa Tyr Tyr Gly Met Asp Val1 5 1013716PRTArtificialConsensus sequenceMISC_FEATURE(7)..(7)Xaa can be Gly or PheMISC_FEATURE(8)..(8)Xaa can be Phe or TrpMISC_FEATURE(9)..(9)Xaa can be His or GlyMISC_FEATURE(14)..(14)Xaa can be Leu and Met 137Arg Ile Ala Ala Ala Gly Xaa Xaa Xaa Tyr Tyr Tyr Ala Xaa Asp Val1 5 10 1513815PRTArtificialConsensus sequenceMISC_FEATURE(5)..(5)Xaa can be Ser or Thr 138Asp Arg Gly Tyr Xaa Ser Ser Trp Tyr Pro Asp Ala Phe Asp Ile1 5 10 15139112PRTArtificialConsensus SequenceMISC_FEATURE(29)..(29)Xaa can be Ile or ThrMISC_FEATURE(41)..(41)Xaa can be Val or LeuMISC_FEATURE(54)..(54)Xaa can be Gly or AsnMISC_FEATURE(107)..(107)Xaa can be Arg or Lys 139Gln Ser Val Leu Thr Gln Pro Pro Ser Val Ser Gly Ala Pro Gly Gln1 5 10 15Arg Val Thr Ile Ser Cys Thr Gly Ser Ser Ser Asn Xaa Gly Ala Gly 20 25 30Tyr Asp Val His Trp Tyr Gln Gln Xaa Pro Gly Thr Ala Pro Lys Leu 35 40 45Leu Ile Tyr Gly Ser Xaa Asn Arg Pro Ser Gly Val Pro Asp Arg Phe 50 55 60Ser Gly Ser Lys Ser Gly Thr Ser Ala Ser Leu Ala Ile Thr Gly Leu65 70 75 80Gln Ala Glu Asp Glu Ala Asp Tyr Tyr Cys Gln Ser Tyr Asp Ser Ser 85 90 95Leu Ser Gly Trp Val Phe Gly Gly Gly Thr Xaa Arg Leu Thr Val Leu 100 105 110140120PRTArtificialConsensus SequenceMISC_FEATURE(30)..(30)Xaa can be And or SerMISC_FEATURE(37)..(37)Xaa can be Ser or ThrMISC_FEATURE(52)..(52)Xaa can be Tyr or HisMISC_FEATURE(54)..(54)Xaa can be Tyr or HisMISC_FEATURE(58)..(58)Xaa can be Ser or AsnMISC_FEATURE(59)..(59)Xaa can be Ser or AsnMISC_FEATURE(69)..(69)Xaa can be Ser or ThrMISC_FEATURE(71)..(71)Xaa can be Val or IleMISC_FEATURE(77)..(77)Xaa can be Ile or MetMISC_FEATURE(83)..(83)Xaa can be Lys or GlnMISC_FEATURE(99)..(99)Xaa can be Arg or LysMISC_FEATURE(100)..(100)Xaa can be Asp or AsnMISC_FEATURE(103)..(103)Xaa can be His, Phe or Try 140Gln Val Gln Leu Gln Glu Ser Gly Pro Gly Leu Val Lys Pro Ser Gln1 5 10 15Thr Leu Ser Leu Thr Cys Thr Val Ser Gly Gly Ser Ile Xaa Ser Gly 20 25 30Gly Tyr Tyr Trp Xaa Trp Ile Arg Gln His Pro Gly Lys Gly Leu Glu 35 40 45Trp Ile Gly Xaa Ile Xaa Tyr Ser Gly Xaa Xaa Tyr Tyr Asn Pro Ser 50 55 60Leu Lys Ser Arg Xaa Thr Xaa Ser Val Asp Thr Ser Xaa Asn Gln Phe65 70 75 80Ser Leu Xaa Leu Ser Ser Val Thr Ala Ala Asp Thr Ala Val Tyr Tyr 85 90 95Cys Ala Xaa Xaa Arg Gly Xaa Tyr Tyr Gly Met Asp Val Trp Gly Gln 100 105 110Gly Thr Thr Val Thr Val Ser Ser 115 120141125PRTArtificialConsensus SequenceMISC_FEATURE(23)..(23)Xaa can be Ala or ValMISC_FEATURE(24)..(24)Xaa can be Ala or ValMISC_FEATURE(31)..(31)Xaa can be Thr or SerMISC_FEATURE(32)..(32)Xaa can be Tyr or PheMISC_FEATURE(54)..(54)Xaa can be Ser or ArgMISC_FEATURE(58)..(58)Xaa can be Arg or IleMISC_FEATURE(60)..(60)Xaa can be His, Try or IleMISC_FEATURE(105)..(105)Xaa can be Pro or GlyMISC_FEATURE(106)..(106)Xaa can be Trp or PheMISC_FEATURE(107)..(107)Xaa can be Gly or HisMISC_FEATURE(112)..(112)Xaa can be Met or Leu 141Glu Val Gln Leu Val Glu Ser Gly Gly Gly Leu Val Gln Pro Gly Gly1 5 10 15Ser Leu Arg Leu Ser Cys Xaa Xaa Ser Gly Phe Thr Phe Ser Xaa Xaa 20 25 30Ser Met Asn Trp Val Arg Gln Ala Pro Gly Lys Gly Leu Glu Trp Val 35 40 45Ser Tyr Ile Ser Ser Xaa Ser Ser Thr Xaa Tyr Xaa Ala Asp Ser Val 50 55 60Lys Gly Arg Phe Thr Ile Ser Arg Asp Asn Ala Lys Asn Ser Leu Tyr65 70 75 80Leu Gln Met Asn Ser Leu Arg Asp Glu Asp Thr Ala Val Tyr Tyr Cys 85 90 95Ala Arg Arg Ile Ala Ala Ala Gly Xaa Xaa Xaa Tyr Tyr Tyr Ala Xaa 100 105 110Asp Val Trp Gly Gln Gly Thr Thr Val Thr Val Ser Ser 115 120 125142121PRTArtificialConsensus SequenceMISC_FEATURE(33)..(33)Xaa can be Gly or AlaMISC_FEATURE(48)..(48)Xaa can be Val or LeuMISC_FEATURE(49)..(49)Xaa can be Ala or SerMISC_FEATURE(53)..(53)Xaa can be Phe or HisMISC_FEATURE(57)..(57)Xaa can be Leu or Ile

142Gln Val Gln Leu Val Glu Ser Gly Gly Gly Val Val Gln Pro Gly Arg1 5 10 15Ser Leu Arg Leu Ser Cys
Ala Ala Ser Gly Phe Thr Phe Ser Ser Tyr 20 25 30Xaa Met His Trp Val Arg Gln Ala Pro Gly Lys Gly
Leu Glu Trp Xaa 35 40 45Xaa Val Ile Ser Xaa Asp Gly Ser Xaa Lys Tyr Tyr Ala Asp Ser Val 50 55
60Lys Gly Arg Phe Thr Ile Ser Arg Asp Asn Ser Lys Asn Thr Leu Tyr65 70 75 80Leu Gln Met Asn Ser
Leu Arg Ala Glu Asp Thr Ala Val Tyr Tyr Cys 85 90 95Ala Arg Glu Arg Thr Thr Leu Ser Gly Ser Tyr Phe
Asp Tyr Trp Gly 100 105 110Gln Gly Thr Leu Val Thr Val Ser Ser 115
120143124PRTArtificialConsensus SequenceMISC_FEATURE(58)..(58)Xaa can be Glu or
LysMISC_FEATURE(103)..(103)Xaa can be Thr or Ser 143Gln Val Gln Leu Val Glu Ser Gly Gly Gly
Val Val Gln Pro Gly Arg1 5 10 15Ser Leu Arg Leu Ser Cys Ala Ala Ser Gly Phe Thr Phe Ser Ser Tyr 20
25 30Gly Met His Trp Val Arg Gln Ala Pro Gly Lys Gly Leu Glu Trp Val 35 40 45Ala Val Ile Trp Tyr Asp
Gly Ser Asn Xaa Tyr Tyr Ala Asp Ser Val 50 55 60Lys Gly Arg Phe Thr Ile Ser Arg Asp Asn Ser Lys
Asn Thr Leu Tyr65 70 75 80Leu Gln Met Asn Ser Leu Arg Ala Glu Asp Thr Ala Val Tyr Tyr Cys 85 90
95Ala Arg Asp Arg Gly Tyr Xaa Ser Ser Trp Tyr Pro Asp Ala Phe Asp 100 105 110Ile Trp Gly Gln Gly
Thr Met Val Thr Val Ser Ser 115 1201441026DNAHomo sapiens 144aactcgggta acaactgagg
gaaccaaacc agagacgcg c tgaacagaga gaatcaggct 60caaagcaagt ggaagtgggc agagattcca ccaggactgg
tgcaaggcgc agagccagcc 120agatttgaga agaaggcaa aagatgctgg ggagcagagc tgtaatgctg ctgttgctgc
180tgccctggac agctcagggc agagctgtgc ctggggcag cagccctgcc tggactcagt 240gccagcagct ttacagaag
ctctgcacac tggcctggag tgcacatcca ctagtgggac 300acatggatct aagagaagag ggagatgaag agactacaaa
tgatgttccc catatccagt 360gtggagatgg ctgtgacccc caaggactca gggacaacag tcagttctgc ttgcaaagga
420tccaccaggg tctgatttt tatgagaagc tgctaggatc ggatatttc acaggggagc 480cttctctgct cctgatagc
cctgtgggccc agcttcatgc ctccctactg ggcctcagcc 540aactcctgca gcctgagggt caccactggg agactcagca
gattccaagc ctactgcca 600gccagccatg gcagcgtctc ctctccgct tcaaatcct tgcagcctc caggcctttg
660tggctgtagc cgcccgggctc ttgcccattg gagcagcaac cctgagtc taaaggcagc 720agctcaagga tggcactcag
atctccatgg cccagcaagg ccaagataaa tctaccacc 780caggcacctg tgagccaaca ggtaattag tccattaatt
ttagtgggac ctgcatatgt 840tgaaaattac caatactgac tgacatgtga tgctgacctg tgataaggtt gagtattat
900tagatgggaa gggaaatttg gggattatt atctcctgg ggacagtttg gggaggatta 960tttattgtat ttatattgaa ttatgtactt
tttcaataa agtcttatt ttgtggctaa 1020aaaaaa 1026145189PRT Homo sapiens 145Met Leu Gly Ser Arg
Ala Val Met Leu Leu Leu Leu Pro Trp Thr1 5 10 15Ala Gln Gly Arg Ala Val Pro Gly Gly Ser Ser
Pro Ala Trp Thr Gln 20 25 30Cys Gln Gln Leu Ser Gln Lys Leu Cys Thr Leu Ala Trp Ser Ala His 35 40
45Pro Leu Val Gly His Met Asp Leu Arg Glu Glu Gly Asp Glu Glu Thr 50 55 60Thr Asn Asp Val Pro His
Ile Gln Cys Gly Asp Gly Cys Asp Pro Gln65 70 75 80Gly Leu Arg Asp Asn Ser Gln Phe Cys Leu Gln
Arg Ile His Gln Gly 85 90 95Leu Ile Phe Tyr Glu Lys Leu Leu Gly Ser Asp Ile Phe Thr Gly Glu 100 105
110Pro Ser Leu Leu Pro Asp Ser Pro Val Gly Gln Leu His Ala Ser Leu 115 120 125Leu Gly Leu Ser
Gln Leu Leu Gln Pro Glu Gly His His Trp Glu Thr 130 135 140Gln Gln Ile Pro Ser Leu Ser Pro Ser Gln
Pro Trp Gln Arg Leu Leu145 150 155 160Leu Arg Phe Lys Ile Leu Arg Ser Leu Gln Ala Phe Val Ala Val
Ala 165 170 175Ala Arg Val Phe Ala His Gly Ala Ala Thr Leu Ser Pro 180 1851461399DNAHomo
sapiens 146ctgtttcagg gccattggac tctccgtcct gccagagca agatgtgtca ccagcagttg 60gtcatctctt ggtttccct
ggttttctg gcatctcccc tcgtggccat atgggaactg 120aagaaagatg ttatgtcgt agaattggat tggatccgg atgcccctgg
agaaatggtg 180gtcctcacct gtgacacccc tgaagaagat ggtatcacct ggacctgga ccagagcagc 240gaggcttag
gctctggcaa aaccctgacc atccaagtca aagagttgg agatgctggc 300cagtacacct gtcacaaagg aggcgaggtt
ctaagccatt cgctctgct gcttcacaaa 360aaggaagatg gaattggc cactgatatt taaaggacc agaaagaacc
caaaaataag 420accttctaa gatcgaggc caagaattat tctggacgtt tcacctgctg gtggctgacg 480acaatcagta
ctgattgac attcagtgct aaaagcagca gaggctctc tgaccccaa 540ggggtgacgt gcgagctgc tacactctc
gcagagagag tcagagggga caacaaggag 600tatgagtact cagtggagt ccaggaggac agtgctgcc cagctgctga
ggagagtctg 660cccattgagg tcatggtgga tgccgtcac aagctcaagt atgaaaacta caccagcagc 720ttctcatca
gggacatcat caaacctgac ccaccaaga actgacagc gaagccatta 780aagaattctc ggcaggtgga ggtcagctgg
gagtaccctg acacctggag tactccacat 840tctacttct cctgacatt ctgcgttcag gtccagggca agagcaagag
agaaaagaaa 900gatagagtct tcacggcaa gacctcagcc acggtcatct gccgcaaaaa tgccagcatt 960agcgtgcggg
cccaggaccg ctactatagc tcatcttggg gcaatggc atctgtgcc 1020tcagttagg ttctgatcca ggatgaaat
ttggaggaaa agtggaagat attaagcaaa 1080atgtttaaag acacaacgga atagaccaa aaagataatt tctatctgat
ttgctttaa 1140acgtttttt aggatcacia tgatatctt gctgtattg tatagtaga tgctaaatgc 1200tattgaaac
aatcagctaa ttatgtata gatttccag ctctcaagt gccatggcc 1260ttcatgctat taaatatt aagtaatta tgtattatt
agtatattac tgtatttaa 1320cgtttctg ccaggatgta tggatgtt cactactta tgacctgac catcaggatc
1380agtccctatt atgcaaat 1399147328PRT Homo sapiens 147Met Cys His Gln Gln Leu Val Ile Ser Trp
Phe Ser Leu Val Phe Leu1 5 10 15Ala Ser Pro Leu Val Ala Ile Trp Glu Leu Lys Lys Asp Val Tyr Val 20
25 30Val Glu Leu Asp Trp Tyr Pro Asp Ala Pro Gly Glu Met Val Val Leu 35 40 45Thr Cys Asp Thr Pro

Glu Glu Asp Gly Ile Thr Trp Thr Leu Asp Gln 50 55 60Ser Ser Glu Val Leu Gly Ser Gly Lys Thr Leu Thr
Ile Gln Val Lys65 70 75 80Glu Phe Gly Asp Ala Gly Gln Tyr Thr Cys His Lys Gly Gly Glu Val 85 90
95Leu Ser His Ser Leu Leu Leu Leu His Lys Lys Glu Asp Gly Ile Trp 100 105 110Ser Thr Asp Ile Leu
Lys Asp Gln Lys Glu Pro Lys Asn Lys Thr Phe 115 120 125Leu Arg Cys Glu Ala Lys Asn Tyr Ser Gly
Arg Phe Thr Cys Trp Trp 130 135 140Leu Thr Thr Ile Ser Thr Asp Leu Thr Phe Ser Val Lys Ser Ser
Arg145 150 155 160Gly Ser Ser Asp Pro Gln Gly Val Thr Cys Gly Ala Ala Thr Leu Ser 165 170 175Ala
Glu Arg Val Arg Gly Asp Asn Lys Glu Tyr Glu Tyr Ser Val Glu 180 185 190Cys Gln Glu Asp Ser Ala
Cys Pro Ala Ala Glu Glu Ser Leu Pro Ile 195 200 205Glu Val Met Val Asp Ala Val His Lys Leu Lys Tyr
Glu Asn Tyr Thr 210 215 220Ser Ser Phe Phe Ile Arg Asp Ile Ile Lys Pro Asp Pro Pro Lys Asn225 230
235 240Leu Gln Leu Lys Pro Leu Lys Asn Ser Arg Gln Val Glu Val Ser Trp 245 250 255Glu Tyr Pro
Asp Thr Trp Ser Thr Pro His Ser Tyr Phe Ser Leu Thr 260 265 270Phe Cys Val Gln Val Gln Gly Lys
Ser Lys Arg Glu Lys Lys Asp Arg 275 280 285Val Phe Thr Asp Lys Thr Ser Ala Thr Val Ile Cys Arg Lys
Asn Ala 290 295 300Ser Ile Ser Val Arg Ala Gln Asp Arg Tyr Tyr Ser Ser Ser Trp Ser305 310 315
320Glu Trp Ala Ser Val Pro Cys Ser 3251482826DNAHomo sapiens 148acaagggtgg cagcctggct
ctgaagtgga attatgtgct tcaaacaggt tgaagagagg 60aaacagtctt ttctgcttc cagacatgaa tcaggtcact
attcaatggg atgcagtaat 120agccctttac atactctca gctggtgtca tggaggaatt acaatataa actgctctgg
180ccacatctgg gtagaaccag ccacaatttt taagatgggt atgaatatct ctatatattg 240ccaagcagca attaagaact
gccaaccaag gaaacttcat tttataaaa atggcatcaa 300agaaagattt caaatcaca ggattaataa aacaacagct
cggctttggt ataaaaact 360tctggaacca catgcttcta tgtactgcac tgctgaatgt cccaacatt tcaagagac
420actgatatgt ggaaaagaca tttcttctgg atatccgcca gatattcctg atgaagtaac 480ctgtgtcatt tatgaatatt
caggcaacat gactgcacc tggaatgctg ggaagctcac 540ctacatagac acaaaatagc tggtagatgt gaagagtta
gagacagaag aagagcaaca 600gtatctcacc tcaagctata ttaacatctc cactgattca ttacaagggtg gcaagaagta
660ctfgtttgg gtccaagcag caaacgcact aggcattgaa gagtcaaaac aactgcaaat 720tcacctggat gatatagtga
taccttctgc agccgtcatt tccagggtc agactataaa 780tgctacagt cccaagacca taatttattg gtagatgcaa
acaacaattg aaaaggttc 840ctgtgaaatg agatacaagg ctacaacaaa ccaacttgg aatgttaaag aatttgacac
900caattttaca tatgtgcaac agtcagaatt ctactggag ccaaacatta agtacgtatt 960tcaagtgaga tgtcaagaa
caggcaaaag gtactggcag ccttgaggt cactgtttt 1020tcataaaaca cctgaaacag tccccaggt cacatcaaaa
gcattccaac atgacacatg 1080gaattctggg ctaacagttg ctccatctc tacagggcac ctacttctg acaacagagg
1140agacattgga cttttattg gaatgatcgt ctttgctgt atgttgcaa tttttctt 1200gattgggata ttaacagat cattccgaac
tgggattaaa agaaggatct tattgttaat 1260accaaagtg ctttatgaag atattcctaa tatgaaaaac agcaatgtg
tgaaaatgct 1320acaggaaaat agtgaactta tgaataataa ttccagttag caggctctat atgttgatcc 1380catgattaca
gagataaaag aatcttcat ccagaacac aagctacag actacaagaa 1440ggagaataca ggaccctgg agacaagaga
ctaccgcaa aactcgctat tgcacaatac 1500tacagttgta tatattctg atctcaacac tggatataaa cccaaattt caattttct
1560gcctgagggg agccatctca gcaataataa tgaattact tccttaacac ttaaaccacc 1620agttgattcc ttagactcag
gaaataatcc caggttaca aagcatccta atttgtctt 1680ttctgtttca agtgtgaatt cactaagcaa cacaatatt ctggagaat
taagcctcat 1740attaaatcaa ggagaatgca gttctctga catacaaac tcagtagagg aggaaaccac 1800catgctttg
gaaaatgatt caccagtg aactattcca gaacagacc tgcctctga 1860tgaattgtc tctgtttg gtagctgaa
tgaggagtg ccatctatta atacttatt 1920tccacaaaat atttggaaa gccactcaa taggattca ctcttgaaa agtagagctg
1980tgggtcaaa atcaatatga gaaagctgcc ttgcaatctg aactgggtt ttccctgcaa 2040tagaaattga atctgcctc
ttttgaaa aatgtattc acatacaaat ctccatgag 2100acacatggtt tcattccct tgataaata cctaggtagg ggattgctg
gccatgat 2160aagcatatgt ttcagttcta ccaatctgt ttccagagta gtgacattc tgtgctcta 2220ccatccat
gtaagaattc cggggagctc catgccttt taattttagc catttctg 2280ctcatttct taaattaga gaattaaggt cccgaagggtg
gaacatgctt catggtcaca 2340catacaggca caaaaacagc attatgtgga cgcctcatgt atttttata gagtcaacta
2400tttctctt atttccctc attgaaagat gcaaaacagc tctctattgt gtacagaaag 2460ggtaaataat gcaaaatacc
tggtagtaaa ataaatgctg aaaattttcc tttaaaatag 2520aatcattagg ccaggcgtgg tggctcatgc ttgtaatcc
agcactttgg taggctgagg 2580taggtggatc acctgaggtc aggagttcga gtccagcctg gccaatatgc tgaaacctg
2640tctactaa aattacaaaa atagccggc catggtggca ggtgcttga atcccagcta 2700ctggggaggc tgaggcagga
gaatcactg aaccaggaag gcagaggttg cactgagctg 2760agattgtgcc actgcactcc agcctgggca acaagagcaa
aactctgtct ggaaaaaaa 2820aaaaaa 2826149629PRTHomo sapiens 149Met Asn Gln Val Thr Ile Gln
Trp Asp Ala Val Ile Ala Leu Tyr Ile1 5 10 15Leu Phe Ser Trp Cys His Gly Gly Ile Thr Asn Ile Asn Cys
Ser Gly 20 25 30His Ile Trp Val Glu Pro Ala Thr Ile Phe Lys Met Gly Met Asn Ile 35 40 45Ser Ile Tyr
Cys Gln Ala Ala Ile Lys Asn Cys Gln Pro Arg Lys Leu 50 55 60His Phe Tyr Lys Asn Gly Ile Lys Glu Arg
Phe Gln Ile Thr Arg Ile65 70 75 80Asn Lys Thr Thr Ala Arg Leu Trp Tyr Lys Asn Phe Leu Glu Pro His
85 90 95Ala Ser Met Tyr Cys Thr Ala Glu Cys Pro Lys His Phe Gln Glu Thr 100 105 110Leu Ile Cys
Gly Lys Asp Ile Ser Ser Gly Tyr Pro Pro Asp Ile Pro 115 120 125Asp Glu Val Thr Cys Val Ile Tyr Glu
Tyr Ser Gly Asn Met Thr Cys 130 135 140Thr Trp Asn Ala Gly Lys Leu Thr Tyr Ile Asp Thr Lys Tyr Val

Val145 150 155 160His Val Lys Ser Leu Glu Thr Glu Glu Glu Gln Gln Tyr Leu Thr Ser 165 170 175Ser
Tyr Ile Asn Ile Ser Thr Asp Ser Leu Gln Gly Gly Lys Lys Tyr 180 185 190Leu Val Trp Val Gln Ala Ala
Asn Ala Leu Gly Met Glu Glu Ser Lys 195 200 205Gln Leu Gln Ile His Leu Asp Asp Ile Val Ile Pro Ser
Ala Ala Val 210 215 220Ile Ser Arg Ala Glu Thr Ile Asn Ala Thr Val Pro Lys Thr Ile Ile225 230 235
240Tyr Trp Asp Ser Gln Thr Thr Ile Glu Lys Val Ser Cys Glu Met Arg 245 250 255Tyr Lys Ala Thr Thr
Asn Gln Thr Trp Asn Val Lys Glu Phe Asp Thr 260 265 270Asn Phe Thr Tyr Val Gln Gln Ser Glu Phe
Tyr Leu Glu Pro Asn Ile 275 280 285Lys Tyr Val Phe Gln Val Arg Cys Gln Glu Thr Gly Lys Arg Tyr Trp
290 295 300Gln Pro Trp Ser Ser Leu Phe Phe His Lys Thr Pro Glu Thr Val Pro305 310 315 320Gln Val
Thr Ser Lys Ala Phe Gln His Asp Thr Trp Asn Ser Gly Leu 325 330 335Thr Val Ala Ser Ile Ser Thr Gly
His Leu Thr Ser Asp Asn Arg Gly 340 345 350Asp Ile Gly Leu Leu Leu Gly Met Ile Val Phe Ala Val Met
Leu Ser 355 360 365Ile Leu Ser Leu Ile Gly Ile Phe Asn Arg Ser Phe Arg Thr Gly Ile 370 375 380Lys
Arg Arg Ile Leu Leu Leu Ile Pro Lys Trp Leu Tyr Glu Asp Ile385 390 395 400Pro Asn Met Lys Asn Ser
Asn Val Val Lys Met Leu Gln Glu Asn Ser 405 410 415Glu Leu Met Asn Asn Asn Ser Ser Glu Gln Val
Leu Tyr Val Asp Pro 420 425 430Met Ile Thr Glu Ile Lys Glu Ile Phe Ile Pro Glu His Lys Pro Thr 435
440 445Asp Tyr Lys Lys Glu Asn Thr Gly Pro Leu Glu Thr Arg Asp Tyr Pro 450 455 460Gln Asn Ser
Leu Phe Asp Asn Thr Thr Val Val Tyr Ile Pro Asp Leu465 470 475 480Asn Thr Gly Tyr Lys Pro Gln Ile
Ser Asn Phe Leu Pro Glu Gly Ser 485 490 495His Leu Ser Asn Asn Asn Glu Ile Thr Ser Leu Thr Leu
Lys Pro Pro 500 505 510Val Asp Ser Leu Asp Ser Gly Asn Asn Pro Arg Leu Gln Lys His Pro 515 520
525Asn Phe Ala Phe Ser Val Ser Ser Val Asn Ser Leu Ser Asn Thr Ile 530 535 540Phe Leu Gly Glu
Leu Ser Leu Ile Leu Asn Gln Gly Glu Cys Ser Ser545 550 555 560Pro Asp Ile Gln Asn Ser Val Glu Glu
Glu Thr Thr Met Leu Leu Glu 565 570 575Asn Asp Ser Pro Ser Glu Thr Ile Pro Glu Gln Thr Leu Leu
Pro Asp 580 585 590Glu Phe Val Ser Cys Leu Gly Ile Val Asn Glu Glu Leu Pro Ser Ile 595 600
605Asn Thr Tyr Phe Pro Gln Asn Ile Leu Glu Ser His Phe Asn Arg Ile 610 615 620Ser Leu Leu Glu
Lys6251502100DNAHomo sapiens 150gggtggctgaa cctgcagcagg ggagagagg ctcccctggg gctgtggggc
tctacgtgga 60tccgatggag ccgctgggta cctgggtggg cccctctc ttctctcc tctgtccag 120gcagggcgct
gcctgcagaa ccagtgagtg ctgtttcag gacccgcat atccgatgc 180agactcaggc tcggcctcgg gcctagga
cctgagatgc taccgatat ccagtatcg 240ttacgagtg tctggcagt atgagggtcc cacagctggg gtcagccact
tctgcggtg 300ttgccttagc tccggcgct gctgctactt cccgcccgc tcagccacca ggctgcagtt 360ctccgaccag
gctgggggtg ctgtgctga cactgcaca ctctgggtg aatcctggc 420caggaaccag acagagaagt ctctgaggt
gaccctgcag ctctacaact cagttaaata 480tgagcctct ctgggagaca tcaaggtgct caagtggcc gggcagctgc
gtatggagtg 540ggagacccc gataaccagg ttgtgctga ggtgcagttc cggcaccgga caccagcag 600cccatggaag
ttgggcgact gcggacctca ggatgatgat actgagctct gcctctgcc 660cctggagatg aatgtggccc aggaattcca
gctccgacga cggcagctgg ggagccaagg 720aagttctgg agcaagtgga gcagccccgt gtgcgttccc cctgaaaacc
ccccacagcc 780tcaggtgaga ttctcgttg agcagctggg ccaggatggg aggaggcggc tgaccctgaa
840agagcagcca acccagctgg agcttcaga aggctgcaa gggctggcgc ctggcacgga 900ggtcacttac cgactacagc
tccacatgct gtctgccc tgtaaggcca agccaccag 960gaccctgcac ctgggaaga tgcctatct ctgggtgct
gcctacaacg tggtgcat 1020ctcctgaac caattggct ctggcctgaa ccagacgtgg cacattctg ccgacacca
1080cacagaacca gtggctctga atatcagcgt ccgaaccaac gggaccacca tgtattggcc 1140agccccggct
cagagcatga cgtattgcat tgaatggcag cctgtgggccc aggacggggg 1200ccttgccacc tgcagcctga ctgcgccga
agaccggat ccggctgaa tggcaacct 1260cagctggagt cgagagtctg gggcaatggg gcaggaaaag tttactaca
ttaccatct 1320tgctctgcg caccgaga agctcacctt gtgtctacg gtctgtcca cctaccactt 1380tgggggcaat
gcctcagcag ctgggacacc gcaccacgtc tcggtaaga atcatagctt 1440ggactctgtg tctgtggact gggcaccatc
cctgtgagc acctgtccc gctcctaaa 1500ggagatggt gtccgtgcc gagatgaaga cagcaaacag gtgtcagagc
atcccgtgca 1560gcccacagag acccaagta cctcagtg cctgcgggct ggtgtagcct acacggtgca
1620ggtgcgagca gacacagcgt ggctgagggg tctctggagc cagccccagc gttcagcat 1680cgaagtgcag gttctgatt
ggctcatct ctgcctcc ctgggagct tctgagcat 1740ccttctctg ggcgtcctg gctacctgg cctgaacagg
gccgcacggc acctgtgcc 1800gccgtgcc acaccctgtg ccagctccg cattgagttc cctggagga aggagactg
1860gcagtggatc aaccagtg actccagga agagcatcc ctgcaggagg cctggtgtg 1920agagatgctc
tgggacaaag gcgagaggac ttagcctctc gagaagacag agctacctga 1980gggtgccct gagctggccc tggatacaga
gtgtccttg gaggatggag acaggtgcaa 2040ggccaagatg tgatcgttga ggctcagaga ggggtgagtga ctgcgccgag
gctacgtagc 2100151662PRTHomo sapiens 151Met Glu Pro Leu Val Thr Trp Val Val Pro Leu Leu Phe
Leu Phe Leu1 5 10 15Leu Ser Arg Gln Gly Ala Ala Cys Arg Thr Ser Glu Cys Cys Phe Gln 20 25 30Asp
Pro Pro Tyr Pro Asp Ala Asp Ser Gly Ser Ala Ser Gly Pro Arg 35 40 45Asp Leu Arg Cys Tyr Arg Ile
Ser Ser Asp Arg Tyr Glu Cys Ser Trp 50 55 60Gln Tyr Glu Gly Pro Thr Ala Gly Val Ser His Phe Leu
Arg Cys Cys65 70 75 80Leu Ser Ser Gly Arg Cys Cys Tyr Phe Ala Ala Gly Ser Ala Thr Arg 85 90
95Leu Gln Phe Ser Asp Gln Ala Gly Val Ser Val Leu Tyr Thr Val Thr 100 105 110Leu Trp Val Glu Ser

Trp Ala Arg Asn Gln Thr Glu Lys Ser Pro Glu 115 120 125Val Thr Leu Gln Leu Tyr Asn Ser Val Lys Tyr
Glu Pro Pro Leu Gly 130 135 140Asp Ile Lys Val Ser Lys Leu Ala Gly Gln Leu Arg Met Glu Trp Glu145
150 155 160Thr Pro Asp Asn Gln Val Gly Ala Glu Val Gln Phe Arg His Arg Thr 165 170 175Pro Ser
Ser Pro Trp Lys Leu Gly Asp Cys Gly Pro Gln Asp Asp Asp 180 185 190Thr Glu Ser Cys Leu Cys Pro
Leu Glu Met Asn Val Ala Gln Glu Phe 195 200 205Gln Leu Arg Arg Arg Gln Leu Gly Ser Gln Gly Ser
Ser Trp Ser Lys 210 215 220Trp Ser Ser Pro Val Cys Val Pro Pro Glu Asn Pro Pro Gln Pro Gln225 230
235 240Val Arg Phe Ser Val Glu Gln Leu Gly Gln Asp Gly Arg Arg Arg Leu 245 250 255Thr Leu Lys
Glu Gln Pro Thr Gln Leu Glu Leu Pro Glu Gly Cys Gln 260 265 270Gly Leu Ala Pro Gly Thr Glu Val
Thr Tyr Arg Leu Gln Leu His Met 275 280 285Leu Ser Cys Pro Cys Lys Ala Lys Ala Thr Arg Thr Leu
His Leu Gly 290 295 300Lys Met Pro Tyr Leu Ser Gly Ala Ala Tyr Asn Val Ala Val Ile Ser305 310 315
320Ser Asn Gln Phe Gly Pro Gly Leu Asn Gln Thr Trp His Ile Pro Ala 325 330 335Asp Thr His Thr Glu
Pro Val Ala Leu Asn Ile Ser Val Gly Thr Asn 340 345 350Gly Thr Thr Met Tyr Trp Pro Ala Arg Ala Gln
Ser Met Thr Tyr Cys 355 360 365Ile Glu Trp Gln Pro Val Gly Gln Asp Gly Gly Leu Ala Thr Cys Ser 370
375 380Leu Thr Ala Pro Gln Asp Pro Asp Pro Ala Gly Met Ala Thr Tyr Ser385 390 395 400Trp Ser Arg
Glu Ser Gly Ala Met Gly Gln Glu Lys Cys Tyr Tyr Ile 405 410 415Thr Ile Phe Ala Ser Ala His Pro Glu
Lys Leu Thr Leu Trp Ser Thr 420 425 430Val Leu Ser Thr Tyr His Phe Gly Gly Asn Ala Ser Ala Ala Gly
Thr 435 440 445Pro His His Val Ser Val Lys Asn His Ser Leu Asp Ser Val Ser Val 450 455 460Asp Trp
Ala Pro Ser Leu Leu Ser Thr Cys Pro Gly Val Leu Lys Glu465 470 475 480Tyr Val Val Arg Cys Arg Asp
Glu Asp Ser Lys Gln Val Ser Glu His 485 490 495Pro Val Gln Pro Thr Glu Thr Gln Val Thr Leu Ser Gly
Leu Arg Ala 500 505 510Gly Val Ala Tyr Thr Val Gln Val Arg Ala Asp Thr Ala Trp Leu Arg 515 520
525Gly Val Trp Ser Gln Pro Gln Arg Phe Ser Ile Glu Val Gln Val Ser 530 535 540Asp Trp Leu Ile Phe
Phe Ala Ser Leu Gly Ser Phe Leu Ser Ile Leu545 550 555 560Leu Val Gly Val Leu Gly Tyr Leu Gly Leu
Asn Arg Ala Ala Arg His 565 570 575Leu Cys Pro Pro Leu Pro Thr Pro Cys Ala Ser Ser Ala Ile Glu
Phe 580 585 590Pro Gly Gly Lys Glu Thr Trp Gln Trp Ile Asn Pro Val Asp Phe Gln 595 600 605Glu
Glu Ala Ser Leu Gln Glu Ala Leu Val Val Glu Met Ser Trp Asp 610 615 620Lys Gly Glu Arg Thr Glu Pro
Leu Glu Lys Thr Glu Leu Pro Glu Gly625 630 635 640Ala Pro Glu Leu Ala Leu Asp Thr Glu Leu Ser
Leu Glu Asp Gly Asp 645 650 655Arg Cys Lys Ala Lys Met 660152360DNAHomo sapiens
152caggtgcagc tgcaggagtc gggcccagga ctggtgaagc cttcacagac cctgtccctc 60acctgcactg tctctggtgg
ctccatcagc agtgggtggtt actactggag ctggatccgc 120cagcaccag ggaagggcct ggagtggatt gggcacatcc
attacagtgg gaacacctac 180tacaacccgt cctcaagag tcgagttacc atatcagtag acacgtctaa gaatcagttc
240tccctgaaac tgagctctgt gactgcccgc gacacggccg tgtattactg tgcgcgaaat 300cgcggttct actacggtat
ggacgtctgg ggccaaggga ccacggtcac cgtctcctca 360153120PRTHomo sapiens 153Gln Val Gln Leu Gln
Glu Ser Gly Pro Gly Leu Val Lys Pro Ser Gln1 5 10 15Thr Leu Ser Leu Thr Cys Thr Val Ser Gly Gly
Ser Ile Ser Ser Gly 20 25 30Gly Tyr Tyr Trp Ser Trp Ile Arg Gln His Pro Gly Lys Gly Leu Glu 35 40
45Trp Ile Gly His Ile His Tyr Ser Gly Asn Thr Tyr Tyr Asn Pro Ser 50 55 60Leu Lys Ser Arg Val Thr Ile
Ser Val Asp Thr Ser Lys Asn Gln Phe65 70 75 80Ser Leu Lys Leu Ser Ser Val Thr Ala Ala Asp Thr Ala
Val Tyr Tyr 85 90 95Cys Ala Arg Asn Arg Gly Phe Tyr Tyr Gly Met Asp Val Trp Gly Gln 100 105 110Gly
Thr Thr Val Thr Val Ser Ser 115 12015423PRTArtificial SequenceHoneybee melittin signal 154Met Lys
Phe Leu Val Asn Val Ala Leu Val Phe Met Val Val Tyr Ile1 5 10 15Ser Tyr Ile Tyr Ala Ala Ala
201556PRTArtificial SequenceHis Tag 155His His His His His His1 5